Genus IBIS.

IBIS PAPILLOSA (Temm.). WARTY-HEADED IBIS.

This Ibis is more common than the Black-headed, and is fond of open places, as well as the sandy shores of the larger streams and rivers. They are seen in flocks in the open country, picking up insects; the stomach of one which I shot contained nothing but the heads, legs, and wing-cases of locusts; that of a second was full of large grasshoppers, and a lizard; that of a third was filled with the *chrysalides*? of butterflies. At the approach of evening the Warty Ibis retires to thick trees to roost, uttering its loud and discordant cry. It breeds during the months of February, March, April, May and June, laying as many as three, and probably four eggs, of a pale bluish-white, slightly streaked and spotted with pale brown, $2\frac{4}{10}$ in. in length, by nearly $1\frac{7}{10}$ in. in width. I found the nest of this Ibis built on the top of a peepul tree (a species of Banian), and containing three young birds, in the month of March.

April 24, 1855.

John Gould, Esq., F.R.S., in the Chair.

The following paper was read:—

1. ON SOME NEW OR LITTLE KNOWN SPECIES OF BIRDS IN THE
DERBY MUSEUM AT LIVERPOOL.

BY PHILIP LUTLEY SCLATER, M.A.

(Aves, Pl. LXXXV.—LXXXVIII.)

The zoological collection of the late Earl of Derby, now at Liverpool, contains one of the largest and finest series of birds at present in existence, many of the examples being valuable not only for their rarity, but also as types of species described long ago by Latham in his 'General History' and other works, and which are hardly to be recognized without examination of the original specimens.

Mr. Thomas Moore, the present Curator, is busily engaged in arranging this mass of materials, and affords every facility to those who are anxious to inspect any of the objects committed to his care. Among the birds I have had an opportunity of examining there during a recent short visit are the following, which I venture to characterize as new.

1. CONIROSTRUM FERRUGINEIVENTRE, Sclater. (Pl. LXXXV.)

C. cærulescenti-schistaceum, pileo et alis caudaque intus nigricantibus: superciliis latis et elongatis albis: subtus intense ferrugineum aut ferrugineo-rufum.

Long. tota 4·9, alæ 2·75, caudæ 2·0.

Hab. in Bolivia.

J. Wolf, lith

M & N. Hanhart, Imp^r

CONIROSTRUM FERRUGINEIVENTRE, *Sclater*.

J. Wolf, lith.

M & N Hanhart, Imp^r

RAMPHOCÆNUS CINEREIVENTRIS, *Sclater*

Wolf, del.

M. & N. Hanhart. imp.

CYPHORINUS ALBIGULARIS, *Sclater*

This is a typical *Conirostrum*, and quite distinct, I think, from any species hitherto described. The members of this genus with which I am at present acquainted are—

1. *CONIROSTRUM CINEREUM* (Lafr. and d'Orb.); D'Orb. Voy. Ois. pl. 59. fig. 1. From Tacna in Peru and Sicasica in Bolivia.

2. *CONIROSTRUM RUFUM*, Lafr. Mag. de Zool. 1843. *Dacnis rufo-cinerea*, Bp. Atti 6^{ta} Riun. Sc. It. 1845, p. 404, et Consp. p. 401. From Bogota.

3. *CONIROSTRUM SITTICOLOR*, Lafr. Rev. Zool. 1840, p. 102. *C. bicolor*, Less.; Gray's Gen. pl. 34. From Bogota.

4. *CONIROSTRUM ALBIFRONS*, Lafr. R. Z. 1842, p. 301; Mag. de Zool. 1843, Ois. t. 35 (from which, I believe, *C. atrocyaneum*, Lafr. R. Z. 1848, p. 9, and *C. cæruleifrons*, Lafr. R. Z. 1842, p. 302, only differ in age or sex). From Bogota.

Conirostrum superciliosum of Hartlaub, R. Z. 1844, p. 215, and Bp. Consp. p. 402. sp. 5, is a true *Sylvicola* of Swainson, the same as *Parula mexicana*, Bp. Consp. p. 310, and has of course nothing to do with these birds.

The only other species that have been referred to this genus, as far as I am aware, are *Conirostrum ornatum*, Townshend, Ann. Lyc. New York, 1851, p. 112, pl. 5. fig. 1, from Texas (the same as *Ægithalus flaviceps* of Sundeval, according to Dr. Hartlaub), and the *C. fuscum* and *colombianum* of Lesson, Descr. d. Mamm. et Ois. pp. 273 and 274, none of which I have as yet recognized.

2. *SYNALLAXIS ERYTHROTHORAX*, Sclater. (Pl. LXXXVI.)

S. fuscus, olivaceo-tinctus, capite obscuriore: gutture nigro-cinereo: alis extus, nisi parte apicali, tectricibus subalaribus et vitta lata pectorali rufo-castaneis: ventre medio cinerascete, lateribus brunnescenti-olivaceis: cauda brunnea: rostro nigro: pedibus brunneis.

Long. tota 5·3, alæ 2·3, caudæ 2·5.

Hab. in America Centrali; Coban et Honduras.

Of this *Synallaxis*, which seems different from all other members of the genus that I am acquainted with, there is a specimen in the Derby Museum procured at Coban by Delattre in 1843. The British Museum contains an example from Honduras, and I have a single skin in my own collection purchased in Paris, which I believe to be from the same locality. The occurrence of species of this group north of the Isthmus of Panama seems hitherto unnoticed, except by the Prince Charles Bonaparte, in a list of a Guatemala collection of birds in these Proceedings for 1837, p. 118, in which he includes the *Synallaxis cinerascens* of Temminck (Pl. Col. 227. fig. 3). But the characters there given do not at all agree with Temminck's bird, and would seem *more* applicable to the present species. Besides, the true *Synallaxis cinerascens* is said to be from Brazil, and is not likely to occur also in Guatemala.

In my specimen of the present species the throat is slightly speckled with whitish.

3. *RAMPHOCÆNUS CINEREIVENTRIS*, Sclater. (Pl. LXXXVII.)

R. olivaceo-brunneus; *capitis lateribus rufis, spatio postoculari nigro*: *alis extus brunnescentibus*: *gutturē albo, nigro-cinerascente striato*: *abdomine cinerascente, medialiter albescentiore, lateraliter autem olivascentiore*: *cauda nigricanti-fusca*: *rostri mandibula superiore nigrescente, hujus autem apice et mandibula inferiore albidis*.

Long. tota 4·0, alæ 2·0, caudæ 1·3.

Hab. in rep. Novæ Grenadæ, Pasto.

A third species of this peculiar genus, beautifully intermediate in colouring as in locality between the *Ramphocænus melanurus* of Brazil and the *rufiventris* of Central America. When I say a *third* species, I am perhaps doing an injustice to M. Lesson, who has already described a third and a fourth. But I have never seen the *Ramphocæni trinitatis** and *viridis*†, and indeed they are hardly likely to be recognized again from such meagre descriptions.

The *Ramphocænus cinereiventris*, of which there is only one example in the Derby Museum, was procured at Pasto, in the mountains of New Grenada, by the indefatigable Delattre. It is a rather shorter-billed bird than the other two to which I have compared it. Like *R. rufiventris*, it has the sides of the head rufous, but differs in showing a well-marked postocular spot. It is also striated on the throat like that species, but has no tinge of rufous on the abdomen, which is darkish cinereous. The tail of the specimen, I regret to say, is not quite perfect, but there is no appearance of the white markings which are the distinguishing characteristic of the Guatemalan bird.

The Derby Museum contains examples of *R. rufiventris* from Coban and Panama, and also specimens of *R. melanurus*. The latter species appears to extend from the Amazon, where Mr. Wallace collected specimens in the neighbourhood of Para, to South Brazil, where Prince Maximilian of Neuwied notices its occurrence under the name of *Troglodytes gladiator*, Beit. iii. p. 751.

4. *CYPHORINUS ALBIGULARIS*, Sclater. (Pl. LXXXVIII.)

C. intense rufo-brunneus: *alis extus obsolete nigro-fasciatis*: *cauda nigra, brunneo fasciata*: *capitis lateribus nigris*: *superciliis posticis et gutture toto pure albis*: *abdomine crissoque nigris, fasciis minutis albidis transvittatis*: *rostro nigro, tomis pallidis*: *pedibus nigris*.

Long. tota 5·75, alæ 2·7, caudæ 2·1.

Hab. in Isthmo Panama.

This fine large typical *Cyphorinus*, distinguishable by its pure white throat and dark closely-banded under plumage, is also due to the

* Lesson, Rev. Zool. 1839, p. 42. *R. pileo rufo*: *dorso et alis brunneo-rufis*: *corpore infra niveo, lateribus griseis*.

† Lesson, Traité d'Orn. p. 377. *Vert-olivâtre en dessus, jaune en dessous*!

researches of M. Delattre, by whom it was brought from the Isthmus of Panama. The only species I can find which resembles it in some degree is *Cyphorinus leucostictus*, Cab. Orn. Notiz. in Wiegmann's Archiv, 1844, p. 206; Schomb. Reise, iii. p. 673. sp. 37, from Mexico and Guiana; but that would appear to be a much smaller bird, and has the under parts from the chin to the belly white, with the sides and crissum reddish-brown.

Among the rare types in the Derby Museum is Mr. Eyton's *Dendrexetastes capitoides* (Cont. Orn. 1851, p. 76). This does not seem to me different from M. de Lafresnaye's *Dendrocolaptes temmincki* (Rev. et Mag. de Zool. 1851, p. 154. pl. 4), named about the same time, but I think the latter term has a slight precedence in point of date, and the bird will therefore stand as *Dendrexetastes temmincki*, if thought worthy of continuing to rank as a separate genus. The Derby Museum specimen is, to judge by its make, decidedly a Cayenne skin. The *Denrocolaptes temmincki* in the Leyden Museum is said to be from Bogota.

Upon examining Dr. Kaup's *Psaris fraseri*, of these Proceedings, 1851, p. 47, I found it the same as *Tityra albitorques*, Du Bus, Bull. Ac. Brux. 1847, xiv. pt. 2. p. 104; and his *Psaris parinus*, ib. p. 48, seems to me very closely allied to, if not identical with, *Pachyramphus atricapillus* (Gm.), Pl. Enl. 687. fig. 1.

I can also confirm what Dr. Hartlaub has said in Wiegmann's Archiv, 1854, that *Todirostrum pectorale*, Kp., of the same paper is *T. granadense*, Hartl., *T. ruficeps*, Kp. = *T. multicolor*, Strickl., and *Setophaga flammea*, Kp. = *S. intermedia*, Hartl. R. Z. 1853, p. 5. But in the two latter cases Dr. Kaup's names were first given, though from the long delay in publishing the Proceedings the others were first published.

When criticising other writers, it is proper also to mention my own mistakes; and I take this opportunity therefore of stating, that my *Tænioptera striaticollis* of these Proceedings, 1851, p. 193 (of which the Derby Museum contains examples), has been long ago named and figured in D'Orbigny's Voyage as *Tyrannus rufiventris*, p. 312. pl. 32. fig. 2.

May 8, 1855.

G. R. Waterhouse, Esq., in the Chair.

Mr. Gould exhibited a portion of a collection of birds formed by Mr. Hauxwell in a district lying on the eastern side of the Peruvian Andes, in the neighbourhood of the River Ucayali, one of the tributaries of the Upper Amazon. Mr. Gould observed, that the exploration of this particular district had been one of the earliest