This article was downloaded by: [OARE Consortium]
On: 04 July 2015, At: 06:03
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: 5 Howick Place, London, SW1P 1WG

Annals and Magazine of Natural History: Series 1

Publication details, including instructions for
authors and subscription information:

<http://www.tandfonline.com/loi/tnah07>

XL.—Contributions to the fauna of Ireland

Frederick McCoy Esq. M.G.S.D.

Published online: 23 Dec 2009.

To cite this article: Frederick McCoy Esq. M.G.S.D. (1845) XL.—Contributions to the fauna of Ireland , Annals and Magazine of Natural History: Series 1, 15:98, 270-274, DOI: [10.1080/037454809495316](https://doi.org/10.1080/037454809495316)

To link to this article: <http://dx.doi.org/10.1080/037454809495316>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs, expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

22. *Sorex? siccatus*, mihi.—Size and proportions of the last nearly, but distinguished by its feet being clad in fur down to the nails, and by its depressed head and tumid bulging cheeks (mystaceal region). Ears large and exposed as in the first two species (*murinus* and *pygmaeus*), and like them having a rounded tapering tail, but somewhat longer in proportion than theirs. Colour a uniform sordid or brownish slaty blue, extending to the clad extremities. Size nearly of the last. Snout to rump $3\frac{1}{2}$ inches; tail $2\frac{2}{3}$; head $1\frac{7}{16}$; palma $\frac{1}{2}$; planta $\frac{13}{16}$; weight $\frac{3}{4}$ oz.

This animal was caught in a wood plentifully watered, but not near the water. I never saw nor heard of it as a tenant of houses, any more than the last-named. It had no musky smell when brought to me dead. I did not examine its glands.

P.S.—The above paper completes the notices of Nepalese Mammals published in India, and, like the papers written there, has been composed without the aid of library or museum.

XL.—*Contributions to the Fauna of Ireland.* By FREDERICK MCCOY, Esq., M.G.S.D.

[With a Plate.]

NOTICES of some of the following animals were read to the Natural-History Society of Dublin at the June meeting for 1844. The Invertebrata are a portion of a large series collected by Mr. McCalla on the west coast of Ireland, and placed in my hands for examination and description by my valued friend Dr. Scouler, whose kindness I have before had to acknowledge for the liberal manner in which he has always allowed me to examine and describe any specimens of interest either in the museum of the Royal Dublin Society or in his own private collection.

MAMMALIA.

Vespertilio Nattereri (Kuhl).—The singularly small number of bats found in Ireland only amounting at present to three, of which one (*V. Daubentonii*) has occurred but in a single instance, renders any addition to their number of peculiar interest to the Irish naturalist, or to those who take an interest in comparing the fauna of Ireland with that of England. The present species, the reddish gray bat of British authors, has not I believe been hitherto recorded in Ireland; a specimen however was brought to me last summer by G. Mangan, Esq., and is now in the museum of the Natural-History Society of Dublin; he killed it near that city, and was of opinion that it was the common species in his neighbourhood: whether the pipistrelle (the most common Irish bat) might have been confounded with it on the wing, or whether it

is really a common species, I cannot say; I can only state the occurrence of the species, and so add it to the list of Irish animals.

AVES.

Tringa rufescens, Buff-breasted Sandpiper.—A specimen of this rare bird, shot by J. Hill, Esq. near the Pigeon House, Dublin, is preserved in the Museum of the Natural-History Society of Dublin. The specimen is in the same plumage as those described by Mr. Yarrell; the markings on the under side of the wings are very beautiful. The specimen noticed is the only Irish one I am aware of.

Sterna leucoptera (Temm.).—A specimen of this beautiful tern was shot by J. Hill, Esq. on the Shannon in 1841, in company with the black tern (*S. nigra*), with which it was confounded; the specimen was sent along with one of the latter species to the Natural-History Society of Dublin, in whose museum they are now preserved; but from their general resemblance to each other, the present species has remained there undistinguished till the present time. As I believe this to be the first record of the occurrence of this bird in Britain, I subjoin a short description of the specimen and a sketch half the natural size, to assist in drawing the

attention of British naturalists to it, as it will probably be found not unfrequent. *Measurement*—Total length, to extremity of tail, about 8 inches; bill from base $10\frac{1}{2}$ lines, from rictus 1 inch 5 lines; greatest depth of both mandibles $2\frac{1}{2}$ lines; middle toe, including the claw, 11 lines in length; tarsus 8 lines; naked portion of the tibia 4 lines. *Colours*—Legs and feet in the preserved specimen pale yellow, probably red in the living bird, claws darker; bill dark blackish brown; head, neck, breast, abdomen,

under wing-coverts and back deep glossy black ; small wing-coverts, tail, and upper and under tail-coverts pure white ; first three quill-feathers black, with a broad longitudinal band of white on their inner webs ; remainder of the wings ash-gray, becoming darker towards the body.

The form, proportions and size of this species are very nearly those of the black tern (*S. nigra*) ; like that species too the webs of the toes are very deeply indented, being reduced to a mere rudiment between the middle and inner toes. The two species are however easily distinguished,—the under wing-coverts of the *S. nigra* are white, of the *S. leucoptera* black ; the tail of the former is dark gray, of the latter pure white ; in the *S. nigra* the throat is white, breast and abdomen dark gray, and the back lead-colour, while in the *S. leucoptera* all those parts are black. I speak of both species in their perfect plumage.

ECHINODERMATA.

*Priapul*us ——— ? Pl. XVI. fig. 1.—A species of *Priapul*us is very common in the sand at low water mark in Connemara, and occasionally found on the east coast : as it does not agree with the figures of Müller or Prof. E. Forbes of the *P. caudatus*, nor agree exactly with their descriptions, I have made a drawing of a moderate-sized specimen for comparison. In a great number of specimens I have examined, the characters seemed to present no variation ; in all of them the trunk is transversely striated, the general proportions thicker and more robust than in *P. caudatus* ; the body uniformly cylindrical, not dilated at the extremities ; the posterior extremity, instead of being dilated and open, is narrowed and closed ; there is no longitudinal striation in any of the specimens at either end of the body ; the posterior extremity is irregularly tuberculated ; the caudal appendage is much thicker and more bushy than in the *P. caudatus*. As I have not seen living specimens I have left the species an open question ; if it should prove distinct from the species described by Müller and Prof. Forbes, it might perhaps be called *P. hibernicus*. Specimens are in the museum of the Natural-History Society of Dublin, and a large series of them, of every age and size, in the museum of the Royal Dublin Society.

*Syrinx granul*osus (McCoy), Pl. XVI. fig. 2.—*Sp. Ch.* Body nearly smooth, very minutely and uniformly granulated, and obsoletely striated transversely ; basal half of the proboscis and the posterior extremity of the body roughened by deep concentric rugæ and small crowded tubercles ; proboscis from the anal pore to the mouth one-fourth the length of the body, posterior extremity mucronate.

This fine species is intermediate in character between the *S. pa-*

pilosus and *S. Harveii*, the body being entirely without the distinct distant papillæ of the former species; while on the other hand, instead of being smooth as in the *S. Harveii*, it is minutely but regularly granulated, the granules being close together and uniformly distributed over the body, except at the posterior extremity and basal half of the trunk, where the surface is rendered rugged by larger tubercles and deep transverse irregular rugæ. Colour usually a uniform dull brown. Length about 7 inches; diameter of body about 9 lines. Very common at Roundstone bay in certain localities.

Syrinx Forbesii (McCoy), Pl. XVI. fig. 3.—*Sp. Ch.* Posterior half of the body suddenly attenuated; extremity dilated, club-shaped; trunk slender, dilated towards the extremity, granulated its entire length to the base of the tentacula, remainder of the body perfectly smooth.

The form and proportions of this species are nearly those of the *S. nudus*, from which it is distinguished by its smaller size, perfectly smooth body, and in having the proboscis granulated to the base of the tentacula. The extremity of the trunk is conical, narrow, surrounded by a circle of short digitate tentacula, behind which the trunk is rather inflated for a short distance and marked with minute granules disposed in circular lines; the remainder of the proboscis, forming about one-fourth of the entire length of the animal, is of a considerably smaller diameter and covered with a minute crowded granulation; with the exception of the trunk, the entire animal is perfectly smooth: where the granular portion ceases the body is abruptly dilated to nearly four times the diameter of the trunk, and continues cylindrical for about the same length, when it again diminishes to about one-third its diameter, and so continues to the posterior extremity, which is dilated into a pear-shaped knob. The anal pore is situated on the dilated smooth portion of the body, a little below its junction with the trunk.

Not uncommon at Roundstone bay, Connemara, from whence there are specimens in the museum of the Royal Dublin Society.

Syrinx tenuicinctus (McCoy), Pl. XIV. fig. 4.—*Sp. Ch.* Cylindrical, diameter of body from the base of the trunk to the posterior extremity perfectly uniform; posterior extremity obtuse, slightly mucronate at the tip; trunk about one-seventh of the entire length; tip obtuse, surrounded by a few rows of minute puncta; entire surface of the animal sharply striated concentrically: with a lens the trunk and a small portion of the posterior extremity are seen to be also marked with numerous faint, short, rather distant, longitudinal striæ.

It is only after much hesitation that I have ventured to cha-

racterize this pretty species. I had long been struck by its peculiar uniform, cylindrical body, short proboscis, blunt posterior end, and tough, finely striated, glossy integument, the species being very common on the west coast of Ireland. I was uncertain whether the species might prove to be the young of the *S. granulosus* which I have just described; I have however recently been able to examine so large a number of the latter species at all ages, that my doubts are entirely cleared up. The adult length of the *S. tenuicinctus* seems to be about three and a half inches: young specimens of the *S. granulosus* of the same size may be readily distinguished by their granulated body (which is proportionably more distinct in the young than in the old individuals), and their posterior extremity being more gradually pointed and remarkably roughened both by transverse wrinkles and granules. From the *S. Harveii* it is distinguished by its tough striated integument, blunt posterior end, the faint and peculiar character of the reticulation at the two ends of the body, and in the entire of the trunk being marked in the same manner as its base. The colour is usually a clear, uniform hair-brown; length about 3 inches; diameter of body 3 lines; diameter of trunk 1 line. The figure given by Pennant of the *Siphunculus nudus* seems to belong rather to this species; the *Phascolosoma carneum* figured by Ruppell from the Red Sea is also similar, but is distinguished by its pink colour, perfectly smooth body and longer proboscis.

Specimens are in the collection of the Royal Dublin Society from the west coast.

BIBLIOGRAPHICAL NOTICES.

Illustrations of Indian Ornithology. By T. C. Jerdon. No. 1. 8vo. Madras, 1843. London, Richardson.

THIS work is intended, when complete, to contain fifty coloured plates of new or unfigured birds from the south of India. The present number comprises twelve of these plates, lithographed and coloured by native Indian artists, and in a style which does them great credit. With a few exceptions the designs are well-drawn, and the colouring is executed with a far higher degree of finish than European artists can generally afford to give to their publications. We understand however that Mr. Jerdon, anxious to make his work as perfect as possible, has made arrangements for the lithographs in the succeeding numbers to be executed in England, though we think that with the resources which the native talent of India can supply, this step was hardly necessary.

The letter-press is drawn up by Mr. Jerdon with much care and attention, comprising ample descriptions of the species figured, with