EXPLANATION OF ILLUSTRATION.

- a.* Twig in fruit, natural size, received from Mr. S. R. Webb.
 - b.* Flower cut open, received from Dr. Schinz as No. 5240 Schlechter.
 - c.* Calyx separated, showing position of double row of immersed oil glands.
 - d.* Showing disc and hair-tipped carpels, and glabrous style.
 - e.* Petal.
 - f.* Fertile stamen.
 - g.* Staminode.
 - h.* Fruit.
 - i.* Seed.
- The figures *b* to *h* are magnified.

A NEW JAPANESE GRATELOUPIA. By E. M. HOLMES,
F.L.S. (Plate III.)

Amongst some marine algæ collected in Japan by Mr. S. Okubo, and brought to me for identification, I noticed one which I was unable to match either at the Kew Herbarium or at South Kensington. Although bearing some resemblance to *Grateloupia filicina*, it differs so much in habit and colour that I have decided to describe it as a new species.

Grateloupia subpectinata, n. sp. — Fronde compressa, plana, ad 15 cm. longa, et 3 mm. lata, pinnatim ramosa, ramis a basi angustiore, longe subuliformibus, inferioribus longioribus, infra medium latere inferiori ramellis brevibus subpectinatis, latere superiori dentibus paucis præditis, ramis supremis simplicibus, brevibus. Color pulchre roseus.

Hab.: Japan, S. Okubo, 1912.

The nearest approach to this species, in the mode of branching, is *Grateloupia Pennatula*, Kuetzing, a native of Cuba ("Tab. Phyc.," vol. xvii., tab. 27, *a*, *b*.), which has similar ramelli, but both the ramuli and ramelli are lanceolate-linear rather than subuliform, and are much shorter in proportion. The rose colour, so far as I know, is never found in forms of *Grateloupia filicina*, although I have seen many forms referred to this species. The structure is, however, typical of *Grateloupia*, and not of loose texture as in the section *Gloiogenia*, to which a rose-coloured species, *G. acuminata*, from Japan, previously described by me, belongs.

Grateloupia subpectinata, nov. sp.