nological division of the newer Tertiaries into older Pliocene, newer Pliocene or Pleistocene, and Post-pliocene is untenable; too much stress having been laid by authors upon the shell-evidence on this point. At the same time, it is not meant to be implied that all the species of the fauna ranged everywhere throughout the area: some in all probability were peculiar to the south, and others to the north.

The presence of *Hippopotamus major* in the pliocene deposits was pointed out as being of great importance in indicating the character of the pliocene land, which, extending between England and the Continent, must have afforded a great system of rivers and lakes, and probably had a comparatively warm temperature, as late as the deposition of the Grays beds, where also (as is well known) occur some southern freshwater shells, now extinct in England.

After some remarks on the negative evidence afforded by this mammalian fauna with regard to the supposed refrigeration of the land during the Pliocene period, Dr. Falconer reviewed the opinions of some English geologists on the physical conditions and faunæ of this region during the newer Tertiary epoch, especially the views of Mr. S. Wood, Mr. Prestwich, and Mr. Trimmer; and concluded with a few remarks on the occurrence of *E. antiquus* in the Cefn and Kirkdale Caves, and of *E. primigenius* in Kent's Hole, and on the non-existence of *E. primigenius* south of the Alps, and its restriction in the United States of America to the Northern and Central States. In the Southern States and in Mexico a distinct fossil species, *E. (Euelephas) Columbi*, hitherto undescribed, occurs along with remains of *Mastodon*, *Myiodon*, *Megatherium*, Horse, &c.

JUNE 17, 1857.

Charles Preston Molony, Esq., Capt. Madras Army, Holles Street, Dublin, and George Robbins, Esq., Grosvenor Place, Bath, were elected Fellows.

The following communications were read:—

1. *On some COMPARATIVE SECTIONS in the OOLITIC and IRONSTONE SERIES of YORKSHIRE.* By JOHN PHILLIPS, M.A., LL.D., F.R.S., F.G.S., Reader in Geology in the University of Oxford.

[Plate VI.]

Introduction.—A geologist well versed in the Oolitic groups of the South of England or the North-west of France finds himself perplexed by the first aspect of the coeval strata in the north of Yorkshire, which are so rich in bands of ironstone, layers of coal, and hundreds of beds of gritstone and shale, as to resemble a tract of old carboniferous, rather than a terrace of oolitic rocks. Nor are the features and physical geography more similar in the two districts—broad mountainous moor-lands in one, rich corn-laden hills in the other. On further research, the strong contrasts which thus appear between the series of the north and south of England are found to

be most remarkable in the group of the Bath or Lower Oolite, though some peculiarities worthy of notice occur in almost every part of the section between the Chalk and the Triassic formations, and even extend to the Chalk itself. The author proposes, on a future occasion, to present a memoir on the Lower Cretaceous and Upper Oolitic deposits, including the Speeton Clay; but his remarks are now limited upwards by the Coralline Oolite and downwards by the Lias, both included. And they relate to the deposition and succession of strata, with some references to the distribution of organic life.

It is now above thirty years since, following the steps of William Smith, I began to draw the parallels of geological time from the south of England into the north-eastern district of Yorkshire. Separated as these districts are by the nonconformity under the Yorkshire Wolds, and still more alienated by modifications affecting more or less all the strata, it was necessary to use great caution in selecting the principal lines of contemporaneity. Guided by the evidence of organic remains and the succession of strata, we traced in the first place the continuous area of the Lias, and marked the place and value of the Marlstone, which Smith, first of all observers, had characterized near Bath. We then defined the Kimmeridge Clay, Coralline Oolite, Calcareous Grit, and Oxford Clay, and beheld with surprise a rock 60 or even 90 feet thick replacing the weak sands of Christian Malford, and holding in abundance the peculiar fossils of Kelloway Bridge. Even this was not achieved without some difficulty; for the Coralline Oolite of Malton contains several fossils identical with, and others not easily separated from, frequent residents in the Inferior Oolite.

Beyond this every fresh step was difficult at that time, and requires wary movements at the present moment.

In my work on the Yorkshire Coast, published in 1829, the section of the cliffs was taken as a type, and the results then affirmed, after three complete surveys of the whole coast, have met with general acceptance. In one important point they have, indeed, been employed by some of my friends with quite as high a trust in their contents as I ever felt,—for I have always regarded as a bold decision my reference of the grey limestone series of White Nab to the Forest Marble and Great Oolite of Bath; and the shelly sandstones of Blue Wick to the Inferior Oolite of Somersetshire (see edition 1, 1829).

On another point, the fixing of the Brandsby slate on a parallel above the assumed Great Oolite, though no doubt has been publicly expressed, my friends can hardly have avoided feeling it; for this rock is analogous to the Stonesfield slate, and that lies under the Oolite of Bath. By three separate investigations I have assured myself of the accuracy of my first determination; and thus, if the place of the Great Oolite be rightly given in the sections of Yorkshire, the Brandsby stone is an equivalent of Hinton Sandstone or Forest Marble.

It is not, therefore, for the purpose of a formal correction, or of an implied amendment of my published sections, that I offer this paper; but for the purpose of recording variations in the deposits,

within the limited area chosen, which may throw light on the history of this peculiar type of the Oolites. This history has now acquired other points of interest than those long known to belong to it. The bands of ironstone which were noticed in several parts of my sections, and which, gathered in fragments on the beach, yielded a few hundred tons of iron in a year, when transported to Newcastle, are now so much explored in many of the interior hills and vales, as to feed long ranges of blast-furnaces on the Tees, and send half a million of tons of ore in a year to the great iron-works in the west of Durham, by railways which bring back in return 100,000 tons of iron. In a few years this vast "yield" will be doubled or trebled, for the iron-ore is inexhaustible, and what is now worked is much surpassed in richness by some as yet only beginning to be known.

It must not be supposed that in every part of the district the Lower Oolites of Yorkshire fit themselves to any one section assumed as a type of their development in the north-east of the county: on the contrary, there occur great variations. Neither can the Bath type of the Oolites be adapted to the Midland districts of England, without some important changes. I propose to show, in the first place, what are the principal differences in Yorkshire, and, by comparing these with the sections in North Lincolnshire and Oxfordshire, to show how real is the connexion and how great is the variation of the several parts of this large Oolitic field.

The Yorkshire coast furnishes one complete section of this series, measurable in every part, which has again and again been measured and remeasured by myself, both before and since the year 1829, the year of the publication of the 'Illustrations of the Geology of Yorkshire.' I have no material change to make in this section; but in regard of the beds at Gristhorpe, which correspond to the Bath Oolite and the strata between it and Cornbrash, some details will be useful which were not needed in that era of geology. Prof. W. C. Williamson * and Prof. J. Morris † have already contributed some observations on this remarkable section, but they do not render it undesirable for me to give measures in detail.

Coast Sections near Scarborough.—A vertical section of the beds on the Yorkshire Coast from the base of the Kimmeridge Clay to the top of the Lias, taken by measure from the face of the cliffs, with only the addition of the Upper Calcareous Grit in Silpho Brow, which is near the coast, is about 1120 feet in thickness, of which 420 feet belong to the Middle Oolitic formation in the subjoined proportions:—

		feet.
Middle Oolite formation.	Upper Calcareous Grit (not seen on the coast, but observed at Silpho, Pickering, Sinnington, &c.)	60 + —
	Coralline Oolite.....	60
	Lower Calcareous Grit.....	80
	Oxford Clay	150
	Kelloway Rock.....	50 to 70
(It is thicker in other places.)		

* Trans. Geol. Soc. 2 ser. vol. v. p. 223; and vol. vi. p. 143; Proc. vol. ii. p. 429.

† Quart. Journ. Geol. Soc. vol. ix. p. 317.

On comparing this with the corresponding section of the south of England, it may be remarked, that only near Weymouth is the Upper Calcareous Grit well exhibited, and that it is there associated with Kimmeridge clay, and contains, like it, *Ostrea deltoidea*. I have seen this once in the Vale of Pickering in Yorkshire. Again we remark the extraordinary development of Kelloway rock in Yorkshire, to ten times its dimensions in Wiltshire; while, on the other hand, the Oxford clay is reduced to one-fourth of the thickness which it has in the Midland Counties, and is a much more sandy and much less fossiliferous deposit. We now proceed to consider the Lower Oolitic formations. (See Sections 1 and 2, Pl. VI.)

Section No. 1 is the well-known general section of the Lower Oolitic series near Bath, as it was understood and named by Smith. Section No. 2 is the corresponding series in Yorkshire.

No. 1.	No. 2.
<p><i>f.</i> Cornbrash.</p> <p><i>e.</i> { 3. Hinton Sandstone. 2. Forest Marble. 1. Bradford Clay.</p> <p><i>d.</i> Great Oolite.</p> <p><i>c.</i> { Upper Fuller's-earth clays. Fuller's-earth rock. Lower Fuller's-earth clays.</p> <p><i>b.</i> Inferior Oolite.</p> <p><i>a.</i> Sands over the Lias.</p>	<p><i>f.</i> Cornbrash.</p> <p>3. Shales, sandstones, ironstones; plants.</p> <p><i>e.</i> { 2. Shelly oolitic beds, ironstone; &c.</p> <p>1. Shales, sandstones, ironstones; plants, coal, freshwater shells.</p> <p><i>d.</i> Oolite of Gristhorp.</p> <p>Shales, sandstones, ironstones.</p> <p><i>c.</i> { Beds of coal, plants sometimes erect.</p> <p><i>b.</i> Dogger and oolitic ironstone.</p> <p><i>a.</i> Sands of Blue Wick.</p>

The two sections being drawn to occupy the same vertical space, the enormous extension of the sandy and the great contraction of the calcareous portions in Yorkshire, corresponding to the more littoral character of that region, is very apparent. There is also very little "clay," as the term is used in the south of England, in this section in Yorkshire; it is usually more shaly, often more sandy or even streaked with sandstone laminæ, so as to resemble what is called "lin and woon" in Lancashire. The following observations supply a few details regarding the several groups *f* to *a*.

f. Cornbrash.—This is often not above 2 feet 6 inches thick, and in this small thickness is a parting. The top is ferruginous, and mostly very shelly; the lower part is shelly, and often contains root-like bodies at the bottom. Pale blue clay 4 or 5 feet below.

In the strata (*e* 3) immediately below Cornbrash, clays and shales predominate over sandstones for about 80 or 100 feet; then we have for the base of this series about 20 or 30 feet of sandstone, resting on pale clays, full of oblique lamination, often containing fragments of wood. The following is a summary of the beds as they appear in Gristhorp Bay*, south of the island, the thicknesses being sometimes measured, sometimes estimated. Total 121 feet 10 inches.

* In one of my later examinations of the Gristhorp Section, I received no small help from Mr. Peter Cullen.

		ft. in.
Sandy and slaty beds below Cornbrash : about 121 feet 10 inches.	<i>f.</i> Cornbrash above.	
	Pale clay, with some root-like masses resting on dark shaly clay	5 0
	Thin skerry sandstone	0 4
	Pale clays	3 ft. to 6 6
	Skerry sandstone and shale, changing to solid sandstone	1 ft. to 6 0
	Shale, darkens at the bottom	14 0
	White sandstone—"pipy"—with two thin partings...	5 6
	Shale, growing <i>dark</i> at the bottom	6 0
	<i>e 3.</i> Pale blue shale and laminated white sandstone (in one place ironstone).....	8 0
	Alternations of pale bluish and purplish shaly clay ...	9 0
	Thin sandstone	
	Series of pale bluish and purplish shaly clays, with bands of nodules of <i>granular iron-ore</i>	29 6
	Variable grits and shales	8 0
	Variable sandstones, with layers of wood, the top very irony ; it is <i>unconformed</i> on the beds below...12 ft. to 20	0
	Pale clays, with a band of jet and some lumps of ironstone at the bottom	4 0

The thicknesses are all variable in a short distance, and the beds of sandstone vary in quality in a few yards.

On the north side of the island, the lower part of this series, below the granular iron-ore bands, appears as under :—

	ft. in.
Sandstone	4 0
Shale	5 0
Sandstones, with shale partings	5 0
Sandstones and shales, in wedge-like alternations; the lowest shale very dark	16 ft. to 21 0
Irregular sandstone, with partings of shale and layers of wood, in the state of jet or coal	12 0

This rests unconformably on the subjacent series, *e 2*.

In the whole of this series, no special plant-bed has been recognized in the Gristhorpe section, nor have any shells been seen there. North of Scarborough, Cycadaceous leaflets occur in a brown ironstone about 40 feet below Cornbrash, and at Scalby Beck is a layer of the leaves which I called *Sphaenopteris latifolia* (Geol. of Yorkshire, ed. 1. p. 148, pl. 7. fig. 18).

e 2.—The beds thus designated are only seen in a complete state on the south side of the island at Gristhorpe; on the north side of the island the top was removed by denudation before the lowest beds of *e 3* were deposited.

Section of the series of subcalcareous and ferruginous beds (*e 2*), some of them yielding marine shells, in Gristhorpe Bay, south of the island :—

		ft. in.
Fossiliferous Beds. <i>e 2.</i>	Hard sandy bed, irony at top, with decomposing bisulphuret of iron and some jet.....	1 3
	Grey streaky sandstone, with some decomposing bisulphuret	2 0
	Brown laminated, partly ferruginous, bed	2 0
	Grey sandy shales, sulphureous	2 6
	Ironstone-balls in a band (<i>Avicula Braamburiensis</i>).....	1 0
	Subcalcareous, partly shelly, bed	2 0
	Ironstone-balls in a band.....	0 6

1857.]

PHILLIPS—OOLITE AND LIAS, YORKSHIRE.

89

	ft	in.
Fossiliferous Beds. $\epsilon 2$ {		
Calcareous bed, with <i>Belemnites Aalenensis</i> at the top, <i>Ostrea Marshii</i> , &c.	2	4
Sulphureous laminated shale, with nodular ferruginous top	1	8

On the north side of the island, as already observed, the upper part is wanting—the uppermost bed seen is the band of ironstone-balls with *Avicula*.

The next series of beds, marked $\epsilon 1$, is found nearly alike on both sides of the island in the upper members, but very unlike in the lower members.

Commencing in the south, we have, mostly of freshwater or estuary origin,—

	Hard sharp laminated sandstone, with thin lines of shale and distinct vertical joints	ft. in.
	Shale, dark and "coaly" at top and bottom, the middle occasionally resembles "seat earth"	5 0
	"Pipy" white sandstone, passing downward to "seat earth," and still lower to shale with ironstone pins	5 0
	Lumpy sandstone, partly dotted with ironstone, jet, and wood. Forms resembling <i>Lutreria</i> , <i>Modiola</i> , but not known for certain to be organic	2 ft. 6 in. to 4 0
Coal Series. {	Dark laminated curled COAL shale, with many plants, and especially <i>Equisetites columnaris</i> . Here is a "sulphur* line"	1 8
	Sandy grey laminated shales, or sandstone, a little pipy ...	2 0
	Shales, with a thin sandstone layer and dark COAL bands, many plants	8 ft. to 9 0
	Sandstone laminated with shale lines and lumps—thin coaly layers, <i>Equisetites</i> , and other plants	4 0
	Sandstone with thin partings, bands, and lumps of iron-ore, and plants,—very variable in thickness and composition.	20 0

Below is the Oolite with *Cricopora*, marked d .

In the cliffs on the north side of the island we have, wholly or mostly of freshwater or estuary origin,—

	Brown sharp laminated sandstone, with fine shale ...	ft. in.
	Pale shale	5 0
	Dark shale	4 ft. } 6 0
	Soft pipy sandstone	2 ft. 6 in. }
	Pale pipy shale	1 ft. 6 in. }
	Dark shale	1 ft. 3 in. }
	White pipy sandstone	3 3
	Dark shale and COAL in waved laminæ (<i>Equisetites</i>)	6 in. }
	Pipy pale sandstone	10 in. }
	Coaly shale	6 in. }
	Laminated shaly sandstone or sandy shale	1 10
Coal Series. {	Pale laminated sandstone and shale	1 ft. 6 in. }
	Dark laminated shale	1 ft. 3 in. }
	Lumpy white sandstone	1 ft. 9 in. }
	Pale shale, with very fine and numerous plants and ironstone pins,—ANODON	1 ft. 8 in. }
	Shale with COAL streaks (pyritous layer at top)	1 ft. 3 in. }
	COAL bed, crossed by "cleat," and of marine or mostly marine origin	1 ft. }

* Such yellow lines are marked by efflorescent sulphate of iron and some free sulphur, derived from decomposed bisulphuret of iron.

	Sandstone with partings of shale, the top "pipy," the bottom laminated;— <i>Tancredia</i> , <i>Modiola</i> , &c.....	ft. in. 4 6
	Soft clay	6 in. }
	Pale shale with <i>Brachyphylla</i> and other plants. 1 ft. }	1 6
	Ironstone in shale, with <i>Trigonia</i> , <i>Cardinia</i> , &c., in situ	1 0
Sandy bed, 5 ft., including	Ironstone top, containing <i>Trigonia</i> , <i>Pinna</i> , <i>Gervillia</i> , and <i>Pecten</i>	1 0
	Sandstone with ironstone pins and small shells	2 6
	Grey sandy part.....	1 6
	Nodular ironstone in laminated shale, with <i>Modiola</i> , &c. Here sulphuret of zinc occurs in nodules and in shells	5 0
	Sandy bed, with ironstones and many shells of <i>Pholadomya acuticosta</i> , <i>Lima gibbosa</i> , &c.	1 0
	Shale with <i>Pholadomya</i> , <i>Ostrea</i> , <i>Avicula</i> , <i>Natica</i> , &c.	1 3
	Skerry sandstones and shale with scattered plants;— <i>Pholadomya</i>	4 0
	Shales with some ironstone in the lower part, scattered traces of plants;— <i>Pentacrinus</i>	12 0
	Below is the Oolite <i>d</i> .	

d.—The oolite with its upper laminæ of *Cricopora* has been estimated at 15 feet. It seems not necessary to enter into details regarding it at present. It may be recognized at the point north of Cayton Bay and at Cloughton, though the superincumbent strata are widely different at these two points. Either this or the beds marked *e 2* appear in all the region to the north as far as Whitby, to the north-west as far as Ingleby, and to the west as far as Thirsk, from whence it can be traced to the southward.

c.—The series below these oolitic beds has been sufficiently detailed in the 'Illustrations of the Geology of Yorkshire.' It consists, from Cloughton northward, in the upper part, of a thick series of sandstones and shales; the sandstones are regular, solid, partially laminated, some containing jet, some waved, some containing branching forms like fucoids. The series changes below to the more usual aspect of shales and sandstones which accompany coal, several rich plant-layers occur, and at about 100 feet deep in it coal is observed.

Thick sandstones succeed, 60 feet.

A series of sandstones and shales, 200 feet. Vertical *Equisetites* in places. COAL.

Grit rocks and thin shales, with ironstone bands and *Cycaduceæ*, 60 feet.

b.—Inferior oolite.

This is represented on the coast by about 70 feet of sandstone—of which

30 feet are fine-grained yellow, micaceous, irony sandstones, in large blocks variously bedded and jointed, containing several layers of pebbles and shells; the top is very irony, but without fossils. *Nerinaea*, *Actæon*, *Trigonia*, *Astarte*, and many other shells.

20 feet are fine-grained yellow micaceous sandstone, with nests or irony masses of *Serpulæ*, *Lingulæ*, &c. (*Lingula Beanii*).

1857.]

PHILLIPS—OOLITE AND LIAS, YORKSHIRE.

91

20 feet are grey and micaceous, soft and argillaceous sandstone, mostly laminated and fissile, gradually passing into the Lias shale below. Irony nests of *Serpula*, *Belemnites*, *Avicula*, *Pinnæ*, &c. (*Facies liassic*.)

Lias shales below.

Inland Section, Hambleton Hills.—We may now refer to a corresponding section within the same geological limits, taken from the ancient sea-cliff—now a bold inland escarpment—of Hambleton, within a small distance of Thirsk. The total thickness of the oolitic series above the Lias is 859 feet.* The Middle Oolite formation consists of

	feet.
Upper Calcareous Grit, as in Wass Bank.....	60
Coralline Oolite	60
Lower Calcareous Grit	100
Oxford Clay	30
Kelloway Rock, including argillaceous beds at bottom	100 or 120

Section No. 3, Pl. VI. represents the Lower Oolite formation in this district. The Cornbrash is not seen here; its place is assumed, with great probability, at about 120 feet below the top of the Kelloway rock.

	f. (Cornbrash, place of.)	ft. in.
Sandstone and Shales, 182 feet.	Purplish and grey shales, yielding springs	80 0
	Yellowish sandstone	
	Grey shales and white sandstone.....	
	Fire-clay. Plants.....	2 0
	Shales and sandstone. Plants.....	20 0
	Sandstone, white, yellow, fine-grained	35 0
	Grey shales	15 0
	Brown iron-ore (rich)	
	Ferruginous sandstone, including towards the base lenticular aggregations of shells (<i>Avicula</i>).....	30 0
	Sandy laminated shale, with shells at bottom	6 0
Calcareous group, 37 ft. 6 in.	Rough nodular shales, with <i>Ostrea Marshii</i> at bottom ...	4 0
	Shales.....	2 0
	Solid gritty calcareous beds ("glance").....	8 0
	Irony argillaceous purple bed	0 9
	Brown laminated sands	0 3
	Blue solid calcareous bed.....	4 0
	Shale and layers of <i>Ostrea crassa</i>	3 0
	Shale	6 0
	Shale and laminated sandstone	3 0
	White soft sandstone	32 0
Grit, Shale, and Coal Series, 122 ft.	Light-coloured thin sandy laminæ	3 0
	Carbonaceous shale or fire-clay (elsewhere COAL in this stage).....	4 0
	White gritstone.....	40 0
	Yellow sandstone	25 0
	Rough nodular calcareous bed ("glance"), with <i>Avicula</i> ..	2 0
	Rubbly sandstone. (<i>Nucleolites</i>).....	18 0

* I have had the advantage of being accompanied in my late surveys of this tract by the Rev. C. Johnstone, Dr. Verity, and Mr. C. Strickland.

		ft.	in.
Ironstone Series, 147 feet.	Grey shale	10	0
	Shale and small pale ironstone pins, with one thin layer of <i>shelly</i> sandstone	12	0
	Cement-stone — pale-coloured limestone in cuboidal blocks. (<i>Glyphia</i>)	3	0
	Dark shale, with ironstone balls. (<i>Pecten</i>)	12	0
	Iron-rock or compacted balls	3	0
	Shale, yellowish-grey and dark	20	0
	Sandstone (Pearson's quarry)	12	0
	Shales	20	0
	Cement-stone nodules	1	0
	Pale clay	1	0
	{ Sandstone, brown, yellow, &c.	40	0
	{ The lower part sometimes very ferruginous	3	0
	Shale and ironstone balls	3	0
	Ferruginous and shelly oolite, resting on upper lias	7	0

The total thickness of the Lower Oolitic Series is thus found to be
489 ft. in this combination.

If we now compare these sections measured on the actual coast and against the ancient sea-cliffs, we find in regard to the Middle Oolite formation only differences of thickness; the Oxford Clay being remarkably thinner near Thirsk, but the Kelloway rock thicker. But in all the strata below the Cornbrash the differences between the two sections are very great.

The Cornbrash, as seen on the Yorkshire coast, is never above 5 feet thick. It contains several distinguishable parts—a thin part at the top rather laminated, and a part at the bottom, argillaceous, or actual clay; while in the middle is one thick mass, or two beds, of shelly and somewhat ferruginous limestone, hardly oolitic. It contains *Nucleolites orbicularis*, *Myacites securiformis*, *Ammonites Herveyi*, and many other shells, but hardly a trace of *Belemnites*. Near Thirsk, and for a great breadth of the moorlands eastward of Hambleton, I have not clearly seen it. It is, however, conspicuous in Newton Dale, and thicker than on the coast.

The series of sandstones and shales between the Cornbrash or Kelloway rocks and the calcareous beds which first succeed below, yields on the coast two bands of ironstone, and contains *Cycadaceæ*; and one band (rich brown iron-ore) appears near Thirsk.

The calcareous shelly beds below, which occur on the coast, and seem to lie in the place of the Forest Marble group of the south of England, appear under Hambleton at the base of the carbonaceous sandstone, and may perhaps be regarded as including the shelly nests in the lower part of the sandstone. A bed of *Ostrea Marshii* occurs in both situations, near the middle part of the shelly series; a bed of *Ostrea crassa* lies towards the lower part at Hambleton; but *Belemnites* and *Terebratulæ* have not been seen there, as at Gristhorpe and Cloughton.

Below this point there is great difficulty in tracing lines of contemporaneity across the district from the coast to the inland cliff. If we suppose, in conformity with many observations, the Gristhorpe plant-bearing series to be merely a local deposit, not seen at any

point north of Haiburn Wyke, nor in Staintondale Cliff, the grit, shale, and coal of Hambleton, 164 feet thick, will be below the whole of the Bath Oolite, and correspond to the Haiburn coal series.

What in this case is the rough nodular "glance" oolite (2 feet) with *Avicula*, resting on rubbly sandstone, with *Nucleolites*, 18 feet? Is it of the date of Stonesfield slate?

The Cricopora-beds of Gristhorp, below the coal series there, lose their distinctness in going northward and westward, so that they become untraceable; on the other hand, we find the Belemnitic beds to fail entirely toward the southward and westward. On a line which runs north and south between the coast and the Hambleton Cliff, and which thus partakes of both these negative geographical influences, neither of them is yet certainly known, though we may expect to find the upper set of beds; nor have we on that line any trace of the Gristhorp plants. The rough nodular "glance" bed is recognized, as I think, on the line of the Whitby railway, about 100 feet below the shelly beds of the Bath Oolite.

Still less obvious are the lines of contemporaneity lower down in the thick series above the Inferior Oolite, when compared on the two parallels from north to south. On the coast are sandstones and shales, with zones of plants, but no calcareous bands, and no remarkable ironstone; under Hambleton, with a smaller thickness of arenaceous and argillaceous sediments, we have one or even two groups of white cement-nodules (compact argillaceous limestone), one containing *Glyphia*, a group of valuable ironstone-nodules, and even a 3-foot band of ironstone. These deposits are only seen in the western range of the Oolites.

The lowest part of the Bath Oolite formation on the Yorkshire coast consists of the ferruginous "dogger," with two bands of shells at the Peak; under it is a group of grey, micaceous, soft, sandy rocks, apparently the "Gold Cap Sandstone" of Lyme Regis, "Sand of the Inferior Oolite" of Bath, and of the Cliff-hill of Lincolnshire. Under Hambleton these are represented by ferruginous oolites, containing *Ammonites*, *Hyboclypeus*, *Trigonia*, &c.; but this character of the rock, by which it approximates to the lower part of the sections of Inferior Oolite in Gloucestershire, ceases as we proceed northward, and is modified as we proceed southward.

On a line drawn from north to south between the coast and the Hambleton, we find the oolitic band at the top of the Lias very rich in iron. In a part of Rosedale, especially, it is an *oolitic iron-rock*, the central part of a great mass on one side of the valley being very attractable by the magnet, and yields, at a maximum, about 50 per cent. of iron.

Other Inland Sections.—We may now proceed to trace briefly the variations which take place in proceeding southward from the country of Thirsk toward the Humber and Lincolnshire, where something nearer the normal type of the Bath Oolites is recognized.

Kilburn.—In the vicinity of Hood Grange, Hoodhill, and Kilburn,

the oolitic limestone (*d*) is frequently quarried; and below it, especially on the southern slope of Rowleston Scar, coal has been obtained within the memory of man only a few feet below the oolite.

About Carlton, Coxwold, Newborough Park, and Owlton, the limestone is frequently exposed and dug in quarries; and here also coal has been found below it. At Newborough Park it was about 50 feet below the oolite, which is dug and burnt to lime.

Brandsby.—Proceeding southward from the country of Thirsk to the parallel of Easingwold, in the Howardian Hills, about Brandsby and Gilling, we find the remains of old coal-works, ranges of limestone, and quarries of roadstone, presenting on the whole the following section:—

Sandy series, yellowish	ft. in.
Roadstone, 2 or 3, or 4 or 5 feet; laminated sandy glance-lime- stone, often full of shells, often in broad flat nodules, lying in soft yellowish sand	50 0
Springy sandy clay below	
Sandy series	
Soft shelly sandstone (at Brandsby)	30 0
Sandy series	
Hard laminated sandy glance-limestone	4 6
Pale clay	1 3
Ironstone	0 2
Pale clay	3 0
Solid limestone beds, 2 ft. 6 in. each	9 ft. or less.
Springs and clay below	
Shale and carbonaceous grits. Signs of COAL	50 0

On Gilling Moor a section is reported by Mr. Winch, 'Geological Transactions,' First Series, vol. v. p. 545, &c., to the depth of 179 ft. 6 in., commencing with—

	ft. in.
Limestone	9 0
Grits and plates	27 0
Hard blue "Calliard"	2 0
Coal (a trace)	
Plate and soft grit	30 0
Blue limestone	4 0
Grits and plates	99 0
Coal	0 6
Shale	2 0

At no point further to the south-eastward has the coal-seam been found worthy of opening,—it has indeed scarcely been traced; but the oolitic limestone, a few yards or fathoms below which it lies, continues distinct, and is quarried almost in a connected line, by Colton, Dawby, and Terrington, to Bulmer and Crambe. The roadstone is found above it distinctly only a short distance, and can scarcely be identified beyond Terrington and Wiganthorpe. At Welburn, Bulmer, and Crambe, white cement-stones occur below the Oolite, as near Thirsk; and below these are oolitic ironstone and ferruginous sands, corresponding to the Dogger-series of Thirsk and the coast; but the strong white gritstones and most of the shales which lie between the Oolite and the Lias about Brandsby, Kilburn, and Boltby are almost absent from the banks of the Derwent.

Crossing the Derwent, we trace in like manner the Oolite by

Westow to its disappearance (through unconformity) below the chalk at Kirby-under-dale. The rock emerges again from its cretaceous covering at Sancton, near Market Wroughton, and is traced to Brough-on-Humber, where it is quarried above grey partly micaceous sand and sandstone, and is evidently equivalent to the Oolite of Lincolnshire, which appears on the opposite shore of the Humber, and rests on sands, grey, red, brown or white, as these do on the Upper Lias.

The only example which I propose now to present of the Lincolnshire sections is that of Harpswell Hill, north of Lincoln, which was traced by myself in 1821, from the actual road-cutting*. The series begins with—

Bath Oolite Series.	{	White oolitic limestone	30 feet.
		Pale clay and sand	12
		White sand	3
		Sandy with iron-balls	10
		White micaceous sand	3
		Brown sand	4
		Clay parting	
		Brown sandstones and shells	40
		Blue Lias clays, succeeded by marlstone and lower lias.....	exposed 20

Above the white oolite are clays and thin stony beds of the Forest Marble series and Cornbrash. These have been fully illustrated in South Lincolnshire by Prof. Morris†, and are found again with similar characters at many localities in Oxfordshire. This white oolite is perhaps generally admitted to be the Great Oolite of Bath,—a point on which I reserve my opinion; it is certainly represented in Yorkshire by that of Cave, Westow, and Crambe.

The Lias Formation.—The Lias varies not so much as the Oolites in a given geographical area, because for the most part its materials were accumulated from long and widely suspended fine argillaceous sediment. The main differences observable in Yorkshire relate to the upper part of this great series, and may be sufficiently illustrated by three actual sections, representing the succession of beds seen in the cliffs about Whitby,—measured in the works at Eston Nab,—and explored in a boring at Feliskirk, near Thirsk.

The Lias of the Yorkshire coast is exhibited in a complete and magnificent section through all the upper and middle parts of the lower groups for above 500 feet in thickness; about 300 feet of beds still lower being imperfectly traced in the interior. In general terms the series‡ stands thus (see Section No. 4, Pl. VI.) :—

Upper Lias Shales, 200 feet.	{	Alum-shale (80+feet), including cement-nodes and ironstone-nodes. In the upper parts, gradually diminishing downwards, are <i>Discina reflexa</i> , <i>Nu- cula ovum</i> , <i>Ammonites bifrons</i> , <i>A. heterophyllus</i> , and <i>Belemnites tubularis</i> . Abundance of Enalio- saurians.
------------------------------------	---	--

* See Memoirs of W. Smith, p. 97.

† Quart. Journ. Geol. Soc. vol. ix. p. 317.

‡ See Illustr. of Geol. of Yorkshire. Also a section of the Upper Lias and Marlstone, communicated to the Geological Society, by Mr. L. Hutton, 25th May, 1836; Geol. Proc. vol. ii. p. 41. This is an elaborate series of the Lias in Rockcliff.

		feet.
Upper Lias	Shale with nodules	60 to 80
Shales,	Jet-rock—shale with jet and pyrites	20
200 feet.	Hard sandy shale	20 to 40
Ironstone	Ironstone-band, or ironstone-rock, with irregular	
and	partings of shale. (<i>Avicula cygnipes</i> , <i>Pecten æqui-</i>	
Marlstone	<i>valvis</i>)	15 to 25
Series,	Shales and ironstone-bands	50 to 70
150 feet.	Sandstone and shale, with <i>Ophiuride</i> , <i>Cardium trun-</i>	
	<i>catum</i> , and many other fossils	40
	Sandstone and shale	30
	Lower Lias Shale below.	

The details of the great ironstone-bands above the marlstone appear in the following section, measured from the workings at Eston Nab*, and communicated to me by Mr. John Murley, C.E. These beds yield about 50,000 tons to the acre.

	ft. in.	ft. in.	ft. in.
Ironstone: top block left as roof	0 11		
(parting)			
Ironstone: second block	2 3		
(parting)			
Ironstone: main block	12 0		
(parting)			
Ironstone: bottom block (variable)	1 10		
Shale	7 0		
Ironstone-band	1 8		
Shale	6 0		
Ironstone-band	0 10		
		17 0	
		15 6	32 6

At Grosmont, near Whitby, the same series stands in this altered and divided state, with about 8 feet of ironstone, yielding about 20,000 tons to the acre.

	ft. in.
Shale	
Ironstone, " <i>Pecten</i> -seam" in two bands, separated by	
1 ft. 6 in. of shale	4 0
Shale	4 0
Ironstone: good	1 0
Shale	7 0
Ironstone: good	1 6
Shale	18 0
Ironstone: " <i>Avicula</i> -seam"	4 0
	38 6

At Feliskirk, in the vicinity of Thirsk, a trial-bore was made, by my advice, through the upper lias, to the ironstone and marlstone, with the following result. The upper bed mentioned is the Inferior Oolite capping the Lias, as at Thirsk, here very ferruginous (see Section No. 5, Pl. VI.):—

	ft. in.
The oolitic iron-rock of Mount St. John	6 to 7 0
Shale, in the upper part cement stones	116 0

* An analysis of the Cleveland iron-ore from Eston is given in Mr. Dick's paper, Quart. Journ. Geol. Soc. vol. xii. p. 357. See also Mr. Crowder's papers, Edinb. New Phil. Journ. New Ser. vol. iii. p. 286, and vol. iv. p. 49.

	ft.	in.
Upper nodular band of ironstone	0	7
Shale	3	0
Lower nodular band of ironstone	0	6
Shale	7	6
Marlstone	1	9
Sandy shale	20	0

Farther to the south this poor representation of the great Ironstone and Marlstone series of the Yorkshire coast is traceable; until in Lincolnshire, Rutland, and Northamptonshire it resumes much of its thickness and importance, though not as yet its commercial value.

Conclusion.—On reviewing the facts here selected to illustrate and exemplify the variations of the middle and lower Oolites, and upper and middle portions of the Lias of Yorkshire, we find sufficient grounds to mark out approximately the geographical ranges of some remarkable mineral conditions—and accumulations of organic remains,—indicative of peculiarities in the depth and currents of the sea, the direction and proximity of land, and other great characteristics of the mesozoic period in this part of the earth's surface.

Reserving for another communication the development of this subject, which cannot be properly examined without additional details, I desire to call attention in the mean time to a few prominent data. (See Section No. 6, Pl. VI.)

The whole series of strata from the Cornbrash to the Marlstone, inclusive, grows thinner towards the south, and in a less degree towards the west; so that from the sea-coast near Whitby to the hills near Thirsk the thickness decreases from 1000 to 590 feet. This appears in each of the main groups. Near Whitby and Guisborough the marlstone and ironstone occupy 150 feet of the cliffs; but near Thirsk only a few feet. The Upper Lias measures 200 feet near Whitby, but near Thirsk less than 120 feet; and in this general diminution each part partakes,—though perhaps the upper part, which is the favourite seat of *Nucula ovum* and *Ammonites bifrons*, has been most severely truncated.

The great series of sandstones, shales, coal-plants, and ironstones, lying above the Lias and below the Oolite of Gristhorp, 500 feet thick in the Peak and the grand range of cliffs at Staintondale, is only 270 feet thick near Thirsk, and is further reduced near Brandsby, until on the banks of the Derwent it is scarcely traceable. The upper carbonaceous series of the coast is of nearly the same thickness at Thirsk, on a line from east to west; but the Oxford Clay diminishes from above 100 to less than 30 feet.

Lines may be drawn in directions not deviating much from E.N.E. to W.S.W. which shall coincide with bands of equal deposition, or equal disappearance, of particular mineral sediments, and particular distributions of organic life. For these lines—as defining equality of earthy sediments,—and thus sometimes marking out similarity of sea-depth and current-action—the author proposes the term “isochthonal,” and is of opinion that the tracing of them will hereafter

tend much to increase our knowledge of the physical conditions of definite epochs in geology. In the present case the normal to these lines, directed to the N.N.W., appears to the author to indicate the existence of land in that quarter during the whole of the period now under consideration,—land at no great distance, because of the proportion of many of the plants,—the occurrence of *Anodon* and other inhabitants of fresh or brackish water,—and the occasional vertical position of *Equisetites*. These latter facts, indeed, may require the admission of the presence of low marshy land within the area in question; but the author is more impressed by the probability of high and extensive land, which might yield to rivers, of some magnitude, the enormous mass of quartz-sand and mixed ferruginous sediments which abound between the truly marine limestones and subcalcareous shelly bands. It seems necessary to suppose some old palæozoic land towards the north, not eminently granitic, but rather of the argillaceous, arenaceous, and quartzose type, with intermixed trap-rocks, such as the Lammermuir Range, which is now so abruptly truncated on its eastern side.

If an isochthonal line be drawn from near Thirsk to Robin Hood's Bay, it will be found that on the north-western side the ironstone band at the base of the Upper Lias is valuable for working—its value growing greater to the north-west.

Nearly on this line the Inferior Oolite receives its maximum dose of iron, growing more and more sandy to the north-west, and losing its thickness and character to the south-east.

If a line be drawn from near Coxwold to Scarborough, parallel to the preceding, it will nearly coincide with the north-western limit of really oolitic character in the calcareous beds of Westow and Gristhorp, and with the south-eastern limit of the coal and freshwater deposits of the district. For a limited breadth north-west of this line, *Belemnites Aulensis* is found in the upper calcareous beds of the Gristhorp series—a position much above what is usual for this shell,—and it is associated at White Nab with *Ammonites Blagdeni*, equally a species more common in the Inferior Oolite of the South of England.

The further development of this subject, the physical condition and depth of the sea-bed, the alternate influence of sea- and fresh-water in the same basin, and the relation of these rocks to those in Lincolnshire and Oxfordshire, must be left for another communication.

-
2. *On the OOLITE ROCKS of GLOUCESTERSHIRE and NORTH WILTS.* By JAMES BUCKMAN, F.G.S., F.L.S., F.S.A., Professor of Natural History, Royal Agricultural College, Cirencester.

[Plate VII.]

Prefatory Remarks.—The object of the present paper is to point out the general geological characteristics of the different members of the Oolite rocks, as they occur in the Cotteswold Hills,

SECTION FROM BIRDLIP TO SWINDON.

Horizontal Scale 7/8 Chaining to 1 Inch

Vertical Scale 1600 Feet to 1 Inch