lation there has always been small and compact. There are no densely settled places, nor are there any sparsely settled ones. The country is so remote that commerce and intercourse with other countries have been very much restricted. There are a large number of Americans living in close proximity to the natives, and the country has a larger proportion of physicians than such small countries usually have—and a good many very able ones, too. Fifty years ago there were no lepers there. Thirty years ago one in each thirty Hawaiians was a leper. There is no absolute record with regard to the first case among them, but there were a number of Chinese imported, and they undoubtedly brought the disease. There was no word in the native language for leprosy until the disease appeared, and it is now called "mai pake," meaning "Chinese sickness." There is some dispute as to the time and place when that name was first given to it, but it is universally used by the natives, and I judge that the disease comes by the name honestly, because, as I say, a lot of Chinese were imported to work on the sugar plantations, and, of course, they brought the leprosy with them. About thirty-five years ago, fifteen years after these Chinese came with their disease, lepers were isolated on the island of Molokai, and a colony established there. We noticed that the disease kept on spreading after the establishment of that colony, until it reached its maximum, about thirty years ago. Since that time it is claimed that the disease has diminished, but my opinion is that it has probably remained about the same, because the population has been reduced, and while there are about 800 lepers in this colony, there are about 400 at large, making an aggregate of 1,200. Therefore, the effect of segregation has evidently been to check any spread. The reason for not isolating cases in the early stage was simply for fear that a correct diagnosis had not been made. When cases were sent to the board of health for examination they were not willing to segregate them, so they were pronounced leper suspects, and sent back to their districts to be kept under surveillance and make periodical reports to the local officers. In that way they were allowed to linger in the district a long time, and conditions got worse. I think that is the way the disease has been allowed to keep on and smolder as it has.

DR. ISADORE DYER—Referring to the point made by Dr. Swarts, I had a case of a girl who had been treated by Dr. Blanc some seven years previous to her calling on me. She bore no evidence of leprosy beyond a true trophic ulcer on the ball of the big toe. I did not examine it bacteriologically. The father of the girl was a typical tubercular leper and a recent case. It was a matter of history that the girl had been infected at least seven years before she came to me. The father was an Irishman who had lived here for years, and the girl was about 22 years old when I saw her.

DR. T. T. FRENCH—As to leprosy being hereditary, there is a home in Honolulu for the non-leprosy children of lepers. The parents often object to their children being taken away, but there is generally an average of twenty or twenty-five girls kept there until they are 18 or 20 years of age. Only one case of leprosy has ever developed there, though we have many children of two leprosy parents. After they leave the home you can not tell anything about them, but when you consider that they stay there until they are about 20 it seems to me that is a pretty strong argument against leprosy being hereditary.

Hemostatic Action of Ovarian Extract.—The tendency to hemophilia seems to belong exclusively to the male sex. It may be transmitted from grandfather to grandson, remaining latent in the mother. This fact and the frequent observation of severe epistaxis in young girls approaching puberty, when the ovarian functions are not yet fully established, suggested to Dr. S. Zavadier of Fouriesburg, southern Africa, that the ovary might possibly be the organ that inhibits the tendency to hemophilia and thus renders women exempt although able to transmit the tendency to their offspring. In a recent severe case of hemophilia, rebellious to all other known measures, he finally administered ovarian extract and with the most encouraging results. The *Semaine Médicale* mentions in quoting his communication that Dr. E. Fuller of New York recently proposed thyroid treatment for the same purpose.

FIXATION OF THE PROLAPSED KIDNEY.*

AUGUSTIN H. GOELET, M.D.

Professor of Gynecology in the New York School of Clinical Medicine; Gynecologic Surgeon to the Metropolitan Hospital.

NEW YORK.

That so many modifications of this operation have been made since it was originally attempted by Hahn in 1881 is the best evidence of the necessity of correcting prolapse of the kidney by permanent fixation, despite the opposition to operative interference in some quarters by those who favor belts and corsets. It suggests also the possible inefficiency of many of these procedures. Almost every well-known surgeon has some special technic of his own, but few of them have many followers. It is my purpose in submitting here a technic of my own, which I have employed now for nearly three years and

Fig. 1.—Showing the relation and attachment of the colon to the kidney, and the relative location of the muscles encountered in reaching the kidney through the vertical lumbar incision.

have found uniformly successful, to point out some of the causes of failure with this operation and show how it may be made successful and satisfactory.

The inadequacy of belts or corsets or any other form of external support when the kidney has descended below the border of the last rib in front so its upper pole may be palpated, is more readily appreciated if we consider what such support does. It is only a substitute for relaxed abdominal walls and the pendant abdominal organs, but in only a small proportion of these cases is the abdominal wall relaxed. The kidney is normally a movable organ, its normal range of mobility downward being from an inch to an inch and a half on deep in-

* Read at the Fifty-fourth Annual Session of the American Medical Association, in the Section on Obstetrics and Diseases of Women, and approved for publication by the Executive Committee: Drs. A. Palmer Dudley, H. P. Newman and J. H. Carstens.

spiration, and it is restrained from further descent only by its surrounding fat and the vessels entering the hilum. To some slight degree additional support is afforded at times by the intestines—the colon in particular, to which it is attached on its anterior surface. This attachment of the colon, however, is one of the chief reasons for its displacement, for when the colon becomes distended it drags on the kidney and the traction is invariably downward when the waist is constricted by corsets or clothing, the distended bowel, as is well known, being crowded below the waist line.

Belts and corsets are not, however, entirely without value, and are useful in the earlier stages of mobility before the kidney has descended below the rib in front, to prevent or retard further descent, and also after operation to support the abdominal organs when necessary and give the patient a feeling of greater security. I believe, however, they are of little or no value in preventing recurrence of the prolapse when the operation for securing fixation of the organ has been properly done. On the other hand, belts and corsets, beside being inadequate, are positively injurious when adjusted to support the kidney after it has descended below the border of the rib because of the irritation to the kidney from the pressure exerted on it. My views on this point correspond with those of Henry Morris of London, who has also had abundant opportunity for careful observation.

The disadvantage, then, of belts or corsets adjusted to support the kidney when it is prolapsed below the ribs in front, is that the pressure necessary to afford effective support will certainly provoke undesirable irritation and eventually lead to serious inflammation in or about the kidney.

INDICATIONS FOR OPERATION.

Prolapse of the kidney to the third degree or beyond, that is, when it has descended below the border of the last rib in front and its upper pole can be palpated when the patient is in the erect position, may be considered sufficient indication for operation because:

1. Prolapse of the organ to that degree permits abnormal mobility and is not conducive to health or comfort though it may not produce intolerable symptoms at first.
2. The position of the kidney when the patient is in a sitting or the erect position interferes with the draining off of the excreted urine, either because of flexure of the ureter or compression of the ureter by the kidney which overlaps it and presses backward on it when the abdomen is constricted by the corsets or clothing.
3. Its circulation and function are seriously impaired by the displacement, and inflammation of its structure frequently results.
4. The kidney when prolapsed to this extent overlaps the ovarian vein as it ascends along the spine, compressing it and retarding the return circulation from the pelvis, thus being a frequent cause of pelvic disease.

If the opposite kidney is sufficiently movable to descend on deep inspiration so that half of the organ can be palpated, it should be fixed at the same time as its fellow to avoid the necessity of subjecting the patient to a second operation, which would be inevitable, because when the kidney is movable to that degree the mobility invariably increases, the degree of descent being progressive. I have several times been obliged to repeat the operation on the left kidney a year or two after fixing the right when either the patient or her physician has opposed a double operation or when the degree of descent of the left kidney at the time of the

first operation was not considered by me sufficient to warrant operation.

Nephropexy is justifiable in hysterical and neurasthenic patients because it is an anomalous condition that should not exist; it is a constant source of irritation; it impairs the health of the patient and may operate as a cause of these nervous disorders by acting as an internal traumatism.¹

PREPARATION OF THE PATIENT FOR OPERATION.

In addition to the usual preparation of the field of operation to secure asepsis, there is some general preparation of the patient that is essential for success. It is useless to delay operation in an attempt to build up a patient whose health is impaired in consequence of this condition, but very much can be accomplished that will be of material aid at the time of operation and during convalescence by correcting digestive disturbances beforehand and securing normal functional activity of the liver, and it is well worth the delay required.

It is very necessary to avoid excessive vomiting following anesthesia because of the strain it puts on the freshly attached kidney, and this can only be obviated by relieving the digestive disturbances beforehand by careful dieting and medication and restoring normal activity of the liver. At least a week should be consumed with such

Fig. 2.—Showing position of the patient on the table, ready for operation, with the air pillow in position under the abdomen.

preparation and in cases of long standing from two to three weeks' preparation will insure a smoother convalescence.

OBJECTS TO BE OBTAINED IN NEPHROPEXY.

1. Permanent fixation of the kidney in its normal position.
2. Complete separation of the kidney from its attachment to the colon and duodenum in front to obviate subsequent dragging on it.
3. Avoidance of injury or mutilation of the kidney.
4. Avoidance of mutilation of the patient by a disfiguring scar or injury of the structures penetrated in gaining access to the kidney.
5. Cure of the symptoms or conditions caused by the prolapse.

TECHNIC OF THE OPERATION.

The patient when anesthetized is turned over on the abdomen and a roll or air-pillow eight inches in diameter is placed under the abdomen which flexes the spine, increases the costo-iliac space and forces the kidney up into position. A vertical incision is made through the skin and fat down to the first layer of muscles beginning just below the last rib and extending downward along the outer border of the erector spinæ muscle for about three and a half inches. This will be about three to four fingers' breadth from the spine according to the

1. Albarran, quoted by Henry Morris.

width of the back. The sheath of the erector spinae should not be opened, but the fibers of the latissimus dorsi are drawn aside with blunt hooks and the lumbar aponeurosis is divided with scissors to the limit of the wound above and below, exposing the quadratus lumborum muscle beneath. Bleeding points should be caught with hemostatic forceps and tied before proceeding.

Fig. 3.—Showing location of the vertical lumbar incision on both sides.

Broad retractors are now inserted and the quadratus lumborum muscle is drawn to one side toward the spine, exposing beneath the transversalis fascia. This fascia is divided with scissors to the extent of the wound above and below, exposing the fatty capsule immediately beneath it. In dividing this fascia it is important to avoid the iliohypogastric nerve, which crosses obliquely in

Fig. 4.—Author's blunt hooks (reduced in size) employed for separating muscles in the direction of their fibers in gaining access to the kidney after incision through the integument and fat.

front of the quadratus lumborum muscle and to include it in the retractor on the spinal side of the incision.

The kidney is now located by inserting the finger and sweeping it around over the surface of its fatty investment. The fatty capsule is then opened with scissors well up toward the upper angle of the incision and well over toward the spine to avoid opening into the peri-

Fig. 5.—Author's broad universal retractors (reduced in size).

toneal cavity or wounding the intestine which bulges up at the lower angle of the wound, and the incision is extended in a vertical direction downward on the finger as a guide. The margin of the fatty capsule on each side is seized with T forceps and traction is made on them to bring the kidney up into the wound and the work of detaching the fatty capsule from the kidney is begun. In

adjusting the T forceps to the outer margin of the fatty capsule care must be taken not to include the intestine or peritoneum.

If the peritoneum is inadvertently opened in exposing the kidney no harm will result if it is immediately closed by a running suture of fine catgut. The incision then to expose the kidney is to be made further over toward the spine. This accident should not occur if the kidney is in position and is properly located before the incision into the capsule is made.

In detaching the fatty capsule from the kidney the finger should be used, aided by blunt scissors where particularly resisting bands of adhesion are encountered. The lower pole of the kidney is to be freed first, then traction on the detached fatty capsule will bring it up into the wound. If the kidney can not be delivered through the incision by traction on the capsule the patient is rolled downward on the air-pillow under the abdomen, which brings its pressure more directly on the border of the ribs and forces the kidney up into the wound.

Traction now downward on the kidney grasped with a gauze pad held between the fingers and traction on the fatty capsule at the same time will succeed in delivering the kidney through the incision on the surface of the back.

The fatty capsule is then completely detached from the kidney around both poles and on each side back to

Fig. 6.—Author's kidney needle (actual size) for inserting the sustaining sutures under the fibrous capsule.

the hilum. Complete detachment of the colon and duodenum in front is thus secured, and behind the kidney is left bare of its fatty investment for subsequent adhesion to the solid structures of the back.

It is not always necessary to deliver the kidney through the incision in order to detach the fatty capsule and insert the sustaining sutures, and it consumes more time. When, therefore, complete detachment can be accomplished without doing so this part of the technic may be omitted.

The pelvis and ureter are now carefully explored with the finger for stone, which should be removed if found before proceeding with the operation.

The fibrous capsule is not detached from the kidney nor is its surface scarified. Firm adhesion of the kidney with its fibrous capsule intact can be secured invariably if the kidney is held immovably in contact with the opposing structures for a sufficient length of time. To accomplish this, however, the suture material employed should be one that does not stretch or become absorbed, and it must be inserted and tied so that the suture loop does not become loosened by cutting into the tissues. We have in silkworm gut a material that will meet these requirements, and to render it pliable it is boiled and preserved in a 3 per cent. solution of lysol. My record of 159 nephropexies by this method without a failure to secure permanent fixation shows conclusively that stripping off the fibrous capsule or further mutilation of the kidney as by sutures which transfix the parenchyma is quite unnecessary.

The sustaining sutures, two in number, are inserted on the posterior aspect of the free border of the kidney. One, the lower, is inserted at about the junction of the middle and lower thirds of the organ and the other at about its center or midway between the poles. The

suture is threaded into small, full-curved needles one inch long, and is bent sharply at the eye of the needle and twisted tightly to avoid tearing the capsule as it is drawn through.

The needle is inserted first from above downward in a direction somewhat obliquely inward to the long axis of the kidney, and penetrates only beneath the fibrous capsule for about half an inch in length and the suture is drawn through; then it is inserted transversely a little below the point of exit of the first insertion, penetrates beneath the fibrous capsule only for a distance of about half an inch and the suture is drawn through, and then again inserted obliquely from below upward at a point opposite the first insertion about an inch from it and the suture is carefully drawn through. The needle is removed and the two ends are clasped temporarily with a hemostatic forceps. The second suture is inserted in a similar manner just above the first. (See Fig. 7.)

Inserted in this manner the suture will resist very considerable strain on it without cutting out—very many

Fig. 7.—Author's sutures for fixing the kidney. Showing their location and method of insertion under the fibrous capsule, which is left intact.

times more strain than the transverse suture, which penetrates the parenchyma, or any suture inserted through the detached fibrous capsule, and it does not mutilate. The stability of this suture seems to be due partly to the exposed portion of the loop left on the outer surface of the capsule and partly to the direction of the traction which is borne mainly by the transverse insertion of the loop and which is in a direction parallel to the capsule or surface of the kidney and not at right angle to it, as is the case with all other sutures employed for fixing the kidney.

Before returning the kidney into its normal position the redundant fatty capsule behind is trimmed off with scissors or torn off, and that in front also, but here care is necessary to avoid encroaching on the peritoneum or bowel. The fat about the lower pole is not removed.

If the kidney has been delivered through the incision it is now replaced and the ends of the sutures are carried through all the structures of the back and brought out on

the surface at the upper angle of the incision—a long, curved perineal needle with eye near the point being utilized for this purpose. The sutures are tied over a fold of several layers of gauze placed lengthwise to the incision. This prevents the suture from cutting through the skin and loosening of the loop. When tied these sutures bring the kidney up into its normal position with its upper pole under the ribs.

A gauze drain is now packed around and under the lower pole of the kidney, forcing back the fat that has not been removed and the bowel which bulges up below the kidney, and the end is brought out at the lower angle of the wound.

The wound is closed by uniting the divided superficial fascia with a running catgut suture and the skin with the subcuticular suture. A bountiful supply of loose aseptic gauze is applied next the wound and over this is placed a thick pad of absorbent cotton covered with gauze, and this is held in place by adhesive strips and a binder encircling the body.

AFTER-CARE OF THE WOUND AND OF THE PATIENT.

Twenty-four hours after the operation the dressing which will be found soaked with drainage from the wound should be removed and renewed, but the wound is

Fig. 8.—Showing position of the gauze pads at the upper angle of the incision, with the sustaining sutures tied over them.

not otherwise disturbed. At the end of forty-eight hours the gauze drain is removed and the dressing renewed. After this the dressing is removed whenever necessary.

The sustaining sutures are not removed until the twentieth day, the day before the patient gets out of bed. The exposed portion of the loop of each suture on one side of the gauze cushion is cut with sterile scissors and the suture pulls out readily, adhering to the gauze when it is lifted off. A simple dressing of gauze is applied afterward.

When renewing the dressings the patient should be turned on the side operated on, and if both kidneys have been fixed at the same time she should be turned on the side that the displacement was greatest.

During the first twenty-four or forty-eight hours the patient will complain of considerable pain in the back and the temperature may rise to 101 or 102 degrees, but both pain and temperature will subside when the drain is removed. An eighth of a grain of morphin may be given hypodermically during the first twelve hours if the pain is sufficiently severe to require it, but it should be avoided when possible. In those cases where drain-

age has not been employed the pain has been quite as severe, so it does not seem to be due altogether to the presence of the gauze, but rather to the manipulation of the structures of the back at the time of the operation.

Vomiting should be avoided if possible, and for this purpose nothing acts so well as oxalate of cerium, in doses of from seven to ten grains, repeated in three to four hours. The stomach should be kept empty for the first twenty-four hours, nothing being given during this time but hot water in small quantity at a time. If thirst is annoying, an enema of normal saline solution should be given. The bowels should be moved on the second day after operation and every day thereafter. Careful dieting will be necessary in some of these cases to keep in abeyance digestive disturbances which may continue after operation, and functional activity of the liver should be preserved throughout convalescence.

During the period of confinement to bed, which is three weeks, the patient should be kept strictly on her back if both kidneys have been operated on. But if only one has been fixed she may turn on the side of the affected kidney. The reason for this is obvious, as it prevents the kidney from dragging on the sutures.

When the patient gets up she should wear an elastic abdominal belt which reaches from the pubes to the level of the lower border of the ribs. This should be worn for at least six months and she should avoid exertion of any kind that would put unusual strain on the kidney. If she has worn corsets before they may be resumed after a month, but the belt must be worn as well, and they must be adjusted so they will be snug over the lower abdomen and hips and loose at the waist and above.

One thing that must not be lost sight of is the digestive disturbances or other disordered conditions that the displaced kidney may have caused. These may not subside unaided after fixation of the kidney, and the patient should be kept under observation and treatment until a fair state of health has been restored. Removal of the cause does not always cure immediately disturbances that have become chronic. Some of these patients show improvement from the beginning of convalescence and require no further attention after they are able to get about again, but others may require attention for some time after.

THE CAUSES OF FAILURE AFTER NEPHROPEXY.

There are many reasons why this operation may prove either a failure or may fail to afford anticipated relief, and to make them more comprehensive they are arranged as follows under separate heads, viz.:

1. Delay in operating until the kidney has become seriously disabled or an incurable nephritis has developed.

2. Delay in operating until the health of the patient and digestive apparatus are seriously impaired or incurable.

3. Unusual and irreparable ptosis of the abdominal organs as a complication of nephroptosis. In these cases, however, since the kidney is many times the most important element of the enteroptosis, fixation of this organ and support of the others by belt places the patient in a comparatively comfortable condition.

4. Failure to prepare the patient properly for the operation.

5. Failure to detach completely the fatty capsule from the kidney—that on the posterior surface to permit the kidney to come into contact with the structures to which it must become adherent, and that on the anterior surface so as to separate the attachment of the colon and

duodenum on the right side and colon and pancreas on the left. Unless this detachment of the bowel from the kidney is complete there will still be dragging on the kidney, which, if it does not pull the kidney away from its attachment, will give rise to annoying pain and other symptoms.

6. Improper insertion of the sustaining sutures whereby they either tear out completely or cut through the fibrous capsule and permit mobility of the kidney before it can become permanently attached.

7. Failure to immobilize the kidney until it can become permanently attached by adhesive inflammation, either by improper tying of the sutures or the use of suture material that stretches when it becomes softened after being buried in the deep tissues of the back or absorbs too quickly, or by fastening the retention sutures to the soft structure only which yields to the constriction of the suture allowing the loop to loosen.

8. Too early removal of the retention sutures.

9. If the remaining fatty capsule is permitted to become interposed between the kidney and the structure to which it should adhere, which may occur either at the time the wound is closed or shortly after, the stability of the attached kidney is greatly lessened. This may be obviated by employing a gauze drain packed around and under the lower pole. An additional advantage of this gauze drain is that it lessens the strain on the retention sutures and prevents the accumulation of blood and serum about the kidney after the operation.

10. Delaying the removal of the gauze drain too long after operation, which would permit accumulation of the discharge back of the gauze.

11. Excessive vomiting or coughing shortly after operation may loosen the kidney from its fresh attachment.

12. Permitting the patient to turn on the opposite side too soon after operation may cause detachment.

13. Permitting the patient to assume the erect position too soon after operation.

14. Neglect to adjust proper support for lax abdominal walls after operation when the patient gets out of bed.

15. Too early resumption of the corsets or their improper adjustment.

16. Permitting too early such exertions as would put unusual strain on the kidney.

17. Attachment of the kidney too low down in an abnormal position whereby it would be constricted and irritated by the pressure of the clothing at the waist or by the corsets.

18. Neglect to correct by remedial measures after operation derangement of the digestive apparatus and other conditions caused by the prolapse.

RESULTS.

When the kidney is properly and permanently fixed in its normal position and the conditions caused by the displacement have been corrected, the result of the operation will be satisfactory. We must not, however, lose sight of the many possible contingencies that may interfere with a final successful issue and lay undue stress on the correction of the displacement alone.

Objection to the operation on the score of its mortality does not hold, for I have done 159 nephropexies on 126 patients, in 33 of these fixing both kidneys at the same time, without a death and without a failure to secure permanent fixation. Cure has resulted in all cases that I have been able to trace in from two to twelve months after operation. In one case only was the operation followed by an unpleasant complication—iodoform pois-

oning resulting from the use of 10 per cent. gauze freshly prepared with ether and alcohol. The patient, however, made a good recovery and was out of bed at the usual time.

It has been objected that the risk of an operation is not warranted in these cases when the symptoms may be relieved by a properly adjusted belt or corset. Estimated from the above report the operation has no mortality, hence there is no risk entailed, and temporary relief of symptoms does not mean a cure. Neither the corset nor the most perfectly adjusted belt will replace the kidney or hold it in its normal position. Nor will it cure the prolapse. But they may and do often provoke serious irritation of the kidney by the pressure they exert which will eventually lead to inflammation in or about the organ.

2030 Broadway.

DISCUSSION.

DR. CARL WAGNER, Chicago—In every operation on so delicate an organ as the kidney cryoscopic examination is a most important preliminary step. This most scientific means of diagnosis of disease of the kidney has been greatly perfected and used extensively with great advantage and satisfaction by the eminent workers, Von Koranyi, Leopold Casper, Friedrich Richter, Kummel, Beckman and others. It is in use all over Europe, while we hear little of it in America as yet. By the author's new suture, which places the kidney in an absolutely physiologic position, is secured a more perfect result than we have ever achieved before. I have used a similar method¹ on the stumps of ablated uteri. I called the suture a guy-rope, and it is intended to hold the stump in position against the parietal wall of the peritoneum. It is fastened over a Maydl bridge and keeps the stump of the uterus in that position until the external wound is closed. The guy-rope is a silk suture carried through the stump and knotted at its end, so that it can easily be removed when the granulating wound is almost healed shut. I have done the same thing on the kidney by putting two lateral sutures through the capsule and fastening over a Maydl's bridge, left *in situ* for twelve days with great satisfaction. The strip of gauze which Dr. Goelet inserts under the lower pole of the kidney for drainage corresponds to some extent to Dr. Senn's method, with the only difference that the gauze strip is used by the latter in larger size and serves as a rest and means of suspension for the lower part of the kidney. I would leave this strip in about six days, so that it not only drains but also favors the development of granulations, as another reliable means for fixation of this organ.

DR. L. H. DUNNING, Indianapolis—I believe we have not yet found the ideal method of operating on the kidney, although 159 successes ought to commend any method. Dr. Goelet's success, however, I think, is due quite as much to the gauze as to the method of placing the stitches. I long ago gave up the idea of attempting to retain the kidney with stitches without opening the fibrous capsule, which, if allowed to remain intact, serves as an obstacle to the formation of firm adhesions unless you get them from the perinephritic fat. I believe that if Dr. Goelet would leave out his gauze and use only his method of suturing he would have an occasional failure. I can not see the advantage of leaving the fibrous capsule intact. I have never observed any ill effects from opening it; I have opened it at least 40 times. The objection that he advanced, that the kidney has been anchored too low by former methods, is true. If the kidney is anchored as indicated in the drawing, through a vertical incision, even this method will leave the kidney lower than normal. We often find that we can palpate only the lower pole of the kidney even when we incise close to the twelfth rib. We must pull the kidney down into the wound. The kidney is about two-thirds of an inch above the twelfth rib in many cases, if not in the average case, so that even in these cases we anchor it lower than normal. But I do not believe that that

cuts any figure whatever in the results. I do believe that the opening of the capsule is very beneficial in overcoming the attendant pain and in helping to establish stronger adhesions to the contiguous parts. The after-treatment is very essential. Constipation and vomiting being among the most frequent causes of poor success in the anchoring of the kidney, I look very carefully after the bowels. I do not give any food that is likely to cause fermentation, and whenever there is a tendency toward, or a desire for, a bowel movement the patient should be assisted in every possible way. This is so important that it is advisable for the operator to look after it himself.

DR. EDWIN RICKETTS, Cincinnati—A record of 159 operations in three years with no failures is the best record in this country or Europe. I want to commend the silkworm gut suggested, and also the practice of bringing the kidney higher up without splitting the capsule; and yet, possibly, there might be one thing to take into consideration, and that is, if in the future that kidney might be affected with Bright's disease, such a splitting of the capsule, if that is a cure of Bright's disease, could have prevented the occurrence of that disease. I agree with Dr. Dunning that without the use of the gauze Dr. Goelet would have had some failures. The gauze must of necessity bring the kidney higher up than any other operation that has been devised. I do not think that it is wise to slit the capsule, because this is a healthy kidney and the slitting opens the door for infection. The operation is to be commended from that standpoint and also because of its simplicity.

DR. JOHN G. CLARK, Philadelphia—What constitutes a movable kidney? In every abdominal section during the last two years where the incision was of sufficient size, I have introduced my hand, passed it backwards into the loin and, with the patient in the Trendelenburg posture, letting the tips of the fingers lightly rest on the ventral aspect of the kidney, gently drawn it downward to its lowest possible point. During this maneuver an assistant measures the distance traversed by the arm to estimate the range of the kidney. I am certain that too little stress has been laid on the fact that under normal conditions the right kidney may be a very movable organ. I have always looked with considerable scepticism on a simple movable kidney as a factor in the exaggerated symptoms which are sometimes attributed to it. If the mobility is slight and the symptoms are markedly of the gastro-intestinal type, a general enteroptosis is usually the principal factor. In the measurements of the range of mobility of the right kidney I have found that it was seldom under 2 cm., and many times from 3 to 4 cm., and yet the history of the case showed no symptom whatever that could be ascribed to this apparently undue excursion of the kidney. The left kidney seldom has a mobility of more than 2 cm. With these definite anatomic facts before us, it would be well to study a case very closely before we determine that a kidney, the lower pole of which may be distinctly felt below the borders of the floating rib, is a distinct pathologic entity. We are frequently told that movable kidneys are not relieved by abdominal binders. This, I believe, is in a large measure true, but in many instances of so-called movable kidneys, where binders or properly fitted corsets are applied, the patient experiences a considerable if not total relief. This is an apparent paradox, but apparent only, for such cases come under the classification of enteroptosis rather than of true movable kidney. I believe that the classical symptoms of marked gastric indigestion more or less dull or acute pain on the right side, marked respiratory mobility of the kidney, which may be so exaggerated under bimanual palpation as to force the kidney well down toward the pelvic brim or Dietel's crises, must be looked on as coming directly from the undue range of mobility, and that such cases may be considered as favorable for symptomatic relief in the event of a successful suspension or fixation of the kidney. In my opinion there are but three methods of fixation which should be considered: 1. By suture, very much as Dr. Goelet has described. 2. By splitting the capsule, turning it back and using it as a suspensory medium. 3. The use of a rubber tube or gauze as a sling to lightly

1. Chicago Med. Recorder and Jour. of Obstet.

draw the kidney up into the wound and hold it there until adhesion has taken place sufficient to retain it in an unmovable position. For more than a year I employed the stitch suggested by Brödel from his anatomic studies, which is similar to that described in Dr. Goelet's paper. The principle of this suture is that by its triangular introduction it snares, as it were, the connective tissue stroma of the kidney and thus bears more strain than does the simple suture passed directly into the cortex of this organ. I do not believe this suture can be improved on. The only question is the serious one, is it advisable to introduce directly into the pulp of the kidney any suture? I know well that, as a rule, no serious complication follows this procedure, but I believe it would be better and more rational to employ the method suggested by Edebohls. For the last year I have constantly employed the latter, and am so far entirely satisfied, although I have not had sufficient experience to justify me in fully commending this method. The third method has not appealed to me as a satisfactory surgical procedure, although Beyer, Penrose and others have had success with this method. Until I have further evidence my choice of method will be that described by Edebohls.

DR. C. S. BACON, Chicago—Does Dr. Goelet use the subcuticular suture and also drainage?

DR. GOELET—I use the subcuticular suture for closing the skin only.

DR. C. L. BONIFIELD, Cincinnati—Many movable kidneys do not really require an operation. A movable kidney is somewhat like a hernia. The vast majority of hernias in the adult require operation, but I have known cases of inguinal and femoral hernia to disappear when the patient became robust and took on additional flesh, and to recur during or after a long spell of illness. I think the amount of mobility of a kidney depends largely on the amount of perirenal fat. Many of the cases operated on for movable kidney do not require surgical treatment at all, although I do not mean to question Dr. Goelet's cases. I would advise the general practitioner to treat the neurasthenia, the indigestion and the general health of the patient.

DR. D. H. CRAIG, Boston—I have seen a number of cases of movable kidney associated with splanchnoptosis that have suffered materially through the perfectly honest and conscientious efforts of general practitioners who, in trying hard to keep abreast of the times, have prescribed abdominal belts. The belt supports temporarily and possibly benefits the patient for the time being, but only in a certain degree. After the belt has served this purpose the patient is left in a worse condition for operative procedure than before.

DR. A. H. GOELET, New York—One reason why this suture is as strong as it is, and why it will pull the kidney up into its normal position under the rib without tearing out, is that no matter which way you pull on it the traction is in an upward direction parallel to the surface of the kidney and its fibrous capsule and not at a right angle to it. There is one objection to leaving the drain in for more than twenty-four hours. I find that when I remove the drain there is a great deal of accumulated discharge behind it, and if this discharge is allowed to remain there it will get in between the kidney and the structures of the back to which the kidney has been attached and thus interfere with the formation of adhesions. That is, the gauze ceases to be a drain after that time. I laid particular stress on the importance of stripping the colon from the kidney. I believe that is one of the most important points in making a success of the operation, and one of the most frequent causes of failure when it is neglected. Where it is left attached to the kidney, if it does not pull the kidney loose it will drag on it and cause pain and other disagreeable symptoms. I have left the gauze drain out in a number of cases and have had success in every instance. Sometimes I think it is not necessary. When there is very little fatty capsule to get out of the way I frequently leave out the gauze drain and my results have been just as good. As to stripping the fibrous capsule from the kidney, I see no reason for mutilating the kidney in this manner when we can secure fixation just as well and better without doing so. If the detached

fibrous capsule is utilized for fixing the kidney it is invariably fixed too low down. I have had a number of patients come to me after operation by that method by other operators, and they were still suffering, though the kidney was fixed, even worse than before the operation. I therefore oppose that method of securing fixation of the kidney. I see no reason for considering this a serious operation, as Dr. Clark contends it is. If I can do 159 nephropexies without a death—and others may have the same results if they adopt the same method—it can not be a very serious operation. My sutures do not penetrate the parenchyma. The needle may wound it in penetrating beneath the fibrous capsule, but that is insignificant and does no harm. Formerly, when a patient would not consent to an operation I yielded and ordered a belt. But now I never prescribe a belt before operation if I can avoid it. In the first place it delays the operation because the patient gets temporary relief, but she invariably comes back worse than before, and the operation becomes a necessity sooner or later. Two years ago I was consulted by a lady with a prolapsed kidney that was causing symptoms. She felt obliged to postpone operation and I prescribed a belt. She returned a little latter saying that she felt so well that she would not have an operation done. Two years afterward she came back to me for another trouble. I asked her to let me examine the kidney, which she said was giving her no inconvenience. I found it fixed in an abnormal position as the result of perinephritic inflammation caused by irritation from the belt, and examination of the urine showed a pyelonephritis, which did not exist at the time of her first visit. This is an illustration of what the belt will do, and emphasizes the necessity of early operation to correct the malposition. The corset, or any other form of external support, is to be placed in the same category with belts and is open to the same objection.

INHIBITED TUBAL PERISTALSIS: A CAUSE OF ECTOPIC GESTATION.*

O. G. PFAFF, M.D.
INDIANAPOLIS.

In submitting this contribution for your consideration, its brevity may be urged as a virtue. A great deal has been written concerning the etiology of ectopic gestation, some of which has approached demonstration, more has been speculative, but certainly much has been done toward the solution of many of the attending phenomena.

The function of the ciliated epithelium of the oviducts has been made much of as the chief, if not the sole, motive force in the transportation of the ovum into the uterine cavity. Building on this assumption Tait and others have attempted to show that in the destruction of the cilia by the inflammatory diseases of the oviducts, the function of the tubes is destroyed and the resulting inertia is to be regarded as the prime factor in the production of tubal pregnancy.

According to Lilienthal, Williams and others, the most frequent site of the chorionic attachment is at the isthmus, near the uterine horn, which would make it appear that if the cilia are really the essential factors of the transportation apparatus the destructive force of the inflammation must have spent itself uniformly in the uterine end of the tube, which is not in accordance with the observed facts in the history of salpingitis. There are many instances of tubal pregnancy in which the cilia appear to be present in normal condition and abundance. In some cases anatomic peculiarities, such as sacculations

* Read at the Fifty-fourth Annual Session of the American Medical Association, in the Section on Obstetrics and Diseases of Women, and approved for publication by the Executive Committee: Drs. A. Palmer Dudley, H. P. Newman and J. H. Carstens.