velopment we find ophthalmology and otology associated both in practice and in teaching and the dissociation is of so recent a date as to be still in progress. Up to thirty-five years ago the appointment of an ophthalmic surgeon to the Massachusetts Charitable Eye and Ear Infirmary carried with it the obligation to receive, and to treat, such cases of disease of the ear as might present themselves in the course of the service and it was not until 1870 that in the distinctive post of aural surgeon was established a step in advance, which was shortly coupled with another, the formal recognition, on the part of the board of managers, of medical teaching as one of the vital functions of an institution for the dispensation of medical charity, and it is only within a year that the joint chair of ophthalmology and otology in one of our larger schools, the University of Michigan, has been divided, and two separate professorships established.

CONCLUSION.

The relationship between otology and laryngology has as little a physiologic foundation as that above mentioned, but between otology and rhinology there is both a physiologic and a practical basis for collaboration, which is however especially concerned with that portion of the organ of hearing devoted to sound transmission and finds its own limitation of relationship in this respect. The service rendered to otology by the later advances in rhinology while computable as great, still remains to be definitely determined, because the end has not yet been reached, and is to be attained only by carefully compared coöperative observation, and it is the necessity for this which makes the true relationship of these specialties. To attain the best results it is important that the two should not be too much merged, since each offers a broad field for study, in its particular region, which it is advisable to conserve, and each has individual points of departure along lines of its own association. To study rhinology from the otologic standpoint alone is greatly to circumscribe rhinology, while to study otology from the rhinologic standpoint only is to leave out of consideration much that is of importance to an organ of special sense which has varied relationships with the functions of other organs and with the well being of the human economy as a whole.

The association of otology with ophthalmology was one of economic origin and without scientific reason, the association with laryngology has always been one of uncertain tenure, and it is the association with rhinology alone that has any substantial ground for persistence, and that to a limited extent, for, inevitably, as rhinology grows, and there could be no better evidence of growth than the number of valuable contributions, both in original research and in clinical observation which mark the program of this and of similar meetings, it will become more and more a distinctive field, capable of commanding the individual devotion of the scientific investigator and earnest practitioners, as otology has done from its beginning.

DISCUSSION.

DR. R. G. SHAMBAUGH, Chicago, said that the very intimate relations which exist between what have sometimes been considered two separate specialties—rhinology and otology—can not be too often or too forcibly emphasized. He thinks that otologists are willing to admit that the most important chapter in recent years, aside from aural surgery, has been added to our knowledge of otology through the study of rhinology.

He fears that some as yet have a very poor appreciation of just what this relationship between rhinology and otology means, and so long as the subject is still of a more or less

indefinite nature, some perhaps are inclined to think that relationships are seen which do not exist. Every rhinologist has had the experience of examining patients who had previously consulted other physicians who had advised operations on the nose, perhaps on the throat, when there was no reason whatever for operative interference. It seems to be not an uncommon thing to advise a patient suffering from chronic catarrh of the middle ear to have any little anatomic variation in the nasal chamber such, for example, as a small crest on the septum, removed for the purpose of benefiting the ear. All sorts of impossible imaginary relations are believed by some to exist between nose and ear trouble. Important relations do exist between middle-ear disease and conditions in the nose and throat, but it does not follow that every case of chronic middle-ear catarrh requires an operation on the nose or throat. The most important relation between the nose and throat and ear trouble is that found when the pharyngeal tonsil is enlarged. Most of the patients in middle life who present themselves for treatment of chronic middle-ear catarrh could have been greatly benefited had the proper attention been given to the adenoids in early life. When these patients are seen in middle life, however, the period when the ear disease is being influenced by the condition in the nose or throat has usually long since passed, and the only interference in the nose or throat that is justifiable is such as may be indicated to relieve local trouble in these parts and no effect on the ear trouble from such treatment can be expected.

DR. OTTO T. FREER, Chicago, referred to Dr. Blake's statement concerning the number of specialties which may be associated by a man wishing to keep his work up to the highest standard of excellence and progress. He agreed with Dr. Blake that this number is diminishing and will in time be limited to one. Twenty years ago the eye, ear, nose and throat physician could keep up his knowledge, technic and instrumentarium, in all his fields. It is impossible for him to do so to-day, when each branch includes a literature so vast and manipulations so difficult that only constant diligence makes it possible to keep up with progress. Inevitably, therefore, the man who assumes so many specialties will slight one or more of them. In this way many eye, ear, nose and throat specialists are to-day making the noble sciences of laryngology and rhinology mere appendages to eye and ear work, timidly and ineffectively entering into the treatment of the diseases of the nose and throat. While in smaller towns it will always be necessary to practice several specialties, in a few years he who does so in the larger cities will be looked on as representing a commercial rather than a scientific class of workers.

MULTIPLE NEUROMATA OF THE SKIN.*

WILLIAM EDGAR DARNALL, A.M., M.D.
ATLANTIC CITY, N. J.

This subject is one that is noticed in very few textbooks of medicine and very scant attention is given to it in those that do refer to it. From the quotations given below it may be seen that the affection is not common, even in the broadest application of the term. I have collected three cases, reported herewith.

This form of tumor presents itself in the form of multiple hard spindle-shaped masses, varying in size from that of a small pea to that of a walnut, and distributed along the course of a single nerve, or of all the nerves of an extremity, or even the whole body. It originates from the connective tissue of the nerve sheaths and usually contains within its substance a certain number of nerve fibers included by the tumor in the course of its development.

"Painful neuroma of the skin is an exceedingly rare form of the affection, only two cases in which the skin was primarily

* Read in the Section on Diseases of Children of the American Medical Association at the Fifty-seventh Annual Session, June, 1906.

affected being on record. Duhring's case¹ was the first noted and was followed by a report of a case by Kosinski.² Both cases occurred in men aged seventy and thirty, respectively. In Duhring's case the tumor had been developing for ten years and in Kosinski's for fourteen. The tumors, varying in size from a pin head to a filbert, confluent and disseminated, were thickly studded over the areas affected. In the first case they extended from the scapula over the shoulder, down the arm to the elbow, occupying principally the area of distribution of the circumflex nerve; and in the second case, over the buttocks and upper part of the thigh corresponding to the area supplied by the small sciatic nerve and the external cutaneous. The lesions were arranged irregularly, not corresponding exactly to the course of the nerves mentioned, and formed firm, flat or elastic nodules oval in shape, fixed in and extending below the skin, and movable only with it. The integument between the nodules was normal, the nodules being purplish or pink in color. In Duhring's case the skin over the nodule was slightly scaly. At the outset pain was variable but later it became excruciating and occurred in paroxysms lasting an hour or more. Since they are non-malignant these growths are best not interfered with unless great distress is caused by the pain. In both of the cases referred to above resort was had to the removal of a portion of the nerve supply with resulting quick relief from pain and a gradual subsidence of the nodules. The new growth is composed of firm connective tissue and non-medullated nerve fibers."³

"The multiple neuromata in some cases number thousands and vary in size from small peas to masses as large as the fist. They are located in three general ways: 1. They may appear along a single nerve, or along several nerve trunks at somewhat regular intervals like beads. 2. They may appear only on a single nerve or its branches. 3. They may invade nearly every nerve in the body, including the sympathetic. In the subcutaneous situations they appear like vascular cirroid dilations, but present a different consistency and anatomic position, and are not modified by pressure on venous or arterial channels. They are most frequently acellular and devoid of myelin. While they may be sensitive and attended by neuralgic pains they are usually unmarked by any sensory disturbance. These cases are sometimes of an hereditary nature."⁴

"The tumor takes its origin either from the endoneurium of a peripheral nerve, pushing aside the nerve fibers until they become widely separated, and even perish from pressure or disappear; or from the perineurium or epineurium, growing laterally from the original nerve trunk, sometimes surrounding it without disrupting the natural arrangement of the nerve fibers. It is, therefore, clear why a histologic study of the neurofibromas, as found in the skin and along the course of peripheral nerves should show gradations relative to the nerve fibers, which are sometimes abundant, again isolated or even absent, and why the fibers vary from apparently normal medullated fibers to naked axis cylinders, or axis cylinders variously formed."⁵

"The term neuroma has been loosely applied to all new growths containing nerve fibers or nerve cells in a fibromatous matrix. Strictly speaking, a neuroma is a tumor composed of newly proliferated proper nerve elements, either neurons or neuraxons, or both. It now appears that certain recently studied neoplasms can be included under this head, e. g., the multiple dermal neuromas found by Knauss in children. They contain both branching ganglion cells resembling those of the sympathetic ganglia, and numerous medullated and non-medullated fibers showing no anatomic connection with the nerves in the neighborhood, and apparently only to be explained on the basis of a proliferation of original nervous elements."⁶

"Neuroma of the skin is an exceedingly rare disease and even in the broadest application of the term it is not common. According to Duhring the subcutaneous nodule of Wood, resembling neuroma closely, differs in being situated in the

subcutaneous tissues, is always freely movable under the skin and is solitary. This and other cases of subcutaneous neuromas of the skin are not, strictly speaking, cutaneous growths, although they are usually so considered in the descriptions of the disease. Nor is Rump's case, sometimes quoted, a clear example of the malady. It consisted essentially of a fibroid tumor of the nerve, and was not accompanied by pain. Histologically, true neuroma of the skin is found seated in the corium extending into the deeper structures. On examination it is found to be made up of firm connective tissues containing non-medullated fibers running up as high as the papillary layer of the corium. They are in reality fibro-neuromata."⁷

CASE REPORTS.

The following case occurred in my practice:

CASE 1.—H. B. M., aged 19.

History.—Patient works at clerical work in an office. He has always been a rather delicate boy and unable to do hard work or engage in rough play like other boys. He has no bad

habits and no venereal history, does not use tobacco or alcohol. Development is poor, as may be seen from the accompanying pictures and his general appearance would indicate a rachitic history. Although 19 years of age he shows no signs of puberty, penis is undeveloped and in size corresponds to the penis of the average ten or twelve-year-old boy. He has no pubic or axillary hair. Some eight years ago his mother noticed that his back was not straight and since then he has developed a pronounced curvature of the spine (scoliosis) in the mid-dorsal region. He is somewhat anemic.

Mentally he is bright and intelligent. He has had no previous illnesses.

About six years ago a tumor about the size of an egg was removed from over his left eye, the nature of which was not recognized at the time.

Three years ago or more he noticed the enlargements here shown but as they gave him no discomfort he paid no especial attention to them. Sometimes he feels a numbness in certain parts of his body affected by the tumors. Pressure on the tumors causes pain which radiates up and down the course of the nerve. He has full use of his arms and legs. He com-

1. Am. Jour. Med. Sciences, October, 1873.

2. Centrbl. f. Chir., No. 16, 1874.

3. "Reference Handbook of the Medical Sciences," p. 259.

4. "Nervous and Mental Diseases," Church and Peterson, p. 279.

5. "American Text-Book of Pathology," p. 175.

6. Ibid., p. 194.

7. "Diseases of the Skin," Stelwagon, pp. 663-664.

plaints of pain in the precordial region and over the sternum, as well as a certain amount of dyspnea at times, all of which can be attributed to the deformity of his chest by reason of the curvature of the spine, and the consequent pressure thus exerted on his lungs and heart. His abdomen is protuberant. The functions of his bowels and bladder are normal and his appetite and digestion good.

Examination.—The movements of the facial muscles are normal. He closes the eyelids firmly, masseters contract firmly, tongue protruded straight and is not atrophied nor does it show fibrillary tremors; extraocular muscles are normal as are also his sensations of touch and pain in the face and over the trunk and extremities.

Movements of his limbs, upper and lower, are not impaired, there is no local atrophy, though he is generally thin and rather wasted. His hand grasp is normal, as are also the Argyll-Robertson reflex, the patellar reflex, and ankle clonus. Babinski reflex is present. His gait is the gait usually seen in sufferers from a curvature of the spine, but shows no other peculiarities. The distribution of the tumors is generally over the scalp, the shoulders and the back, the pectoral regions, and around the loins, as well as up and down the arms, in bead-like chains corresponding nearly to the course of the nerves supplying the various parts. They are elongated pea-like bodies freely movable under the skin rather than attached to it, and vary in size from the porcelain head of a shawl pin to the size of the end of one's thumb. There is no pain associated with them unless they are firmly grasped between the fingers.

Pathologic Examination.—Two of these tumors were excised under cocain anesthesia and submitted for a pathologic examination to Dr. Harry A. Duncan, pathologist to the Samaritan Hospital, Philadelphia. He reports:

"Macroscopically, the tumors are smoothly ovoid in shape, the size of a bean, uniformly firm, whitish gray in color and surface smooth. The smooth surface appears to be a white glistening capsule loosely adherent. To one side of the tumor examined is attached a bit of the external sheath of the nerve.

"Microscopically, the capsule consists of fibrous and loose areolar tissue, and in places is characterized by leucocytic invasion pointing to inflammatory changes. The epineurium or general nerve sheath shows a loose areolar tissue. Clumps of fat cells are numerous. Between the fat cells an infiltration suggests inflammatory changes. The tumor substance is made up of wavy fibrous tissue, although areas of it are undoubtedly myxomatous. Some nerve tissue is present. Cross sections show a separation of the nerve from the perineurium, probably due to inflammatory exudate. The perineurium has undergone a fibrous proliferation. Changes of a similar nature are progressing in the endoneurium and causing a separation or dissociation of the individual nerve fibers from one another. A transverse section of a nerve shows such decided fibrous change in the endoneurium that almost all signs of true nerve tissue are destroyed. The tumor is undoubtedly a neurofibroma, and is doubtless only the localized evidence of a diffused fibromatosis along the nerves."

I am also indebted to Dr. Duncan for the notes of the following case:

CASE 2.—H. C., aged 22, clothing cutter by trade, German by birth.

History.—Family history negative except that one sister suffered from similar growths. The patient had measles when a child, no other diseases, and no bad habits. No venereal history. The tumors are shown all over the body, following the course of the nerves, and were first noticed on his arm when he was 10 years old. For a time their growth was slow, but recently it has been very rapid. The tumors are hard and fusiform in shape, and freely movable under the skin.

Operations.—One tumor was opened at St. Joseph's Hospital before the patient came under observation by me, but was not removed on account of its size. Following this operation he was affected with foot drop. His pulse was 96 to 118, temperature 99 F. Respirations 20 to 24. Urine normal. Blood showed moderate anemia. Many of the tumors were removed. They varied in size from a peanut to a goose-egg. General health improved and he left the hospital. The growths con-

tinued to increase in size and I next learned of his presence in the Philadelphia Hospital. At this institution, because of a very large and painful sarcoma in the course of the sciatic nerve, a hip joint amputation was considered, but before any radical treatment was instituted he died from general exhaustion.

This patient also had a sister, aged 23 years, a tailor-ess, who had all over her body small elongated tumors of a similar nature, along the course of the nerves. They appeared when she was about 16 years old. Several of them had been removed. In both her case and her brother's microscopic examinations were made and the tumors were found to be neurofibromata. One tumor in the brother showed on examination a distinct tendency to sarcomatous change.

A case has been reported by Bland-Sutton, and Thompson,⁸ of Edinburg, refers to some interesting complications in his monograph on the subject. In reporting Keen's case Spiller has searched the literature and sifted it well, and has written a good report, though it is not all pathologic. Every article lays stress on the hereditary tendency of the malady, the painlessness of the tumors, and the long period of development, except when associated with sarcoma. Thompson thinks surgical interference often causes sarcomatous changes to set in.

Through the courtesy of Dr. William G. Spiller of Philadelphia I add the following case, the notes of which he has kindly sent me:

CASE 3.—W. R., aged 33, was admitted to the University Hospital May 12, 1905.

Special History and Habits.—Patient is a farmer and lumberman. Work is very heavy and hard; has not been able to work for about six months; uses alcohol to moderation, never to excess; tobacco, rather freely. Denies venereal history.

Family History.—Father, mother, three brothers and four sisters living and well. Patient is the oldest of the family. One brother dead-born. Has wife and two children, one 3 years old, one 14 months old, and in good health.

History of Previous Illness.—Had ordinary diseases of childhood. Six years ago had one tumor the size of a hazelnut removed from inner surface of left elbow; one from right axilla, one from outer surface of left leg about midway between knee and ankle, one just below internal malleolus of left leg, one just below external malleolus of right foot. The one on the left elbow had appeared about five years before removal, the others two to three years before removal. They were increasing very slowly in size and were very tender to touch or to any unusual strain put on the part. There was usually some spontaneous pain after he had retired at night. Following the operation for removal of these tumors patient had a mild ether pneumonia from which he recovered without complication. He was otherwise in good health to about a year ago, when present trouble began.

History Present Illness.—About eleven months ago patient began to have a feeling of mild pain and numbness in his hands, especially in the morning. This usually improved after he had worked a little. There was no weakness in the hands at this time. The pain and numbness increased in severity and extended up the arm and seemed to affect especially the thumbs and middle line of the arms. The right hand and arm have been more affected than the left hand and arm from the first. In June, 1904, he consulted a physician and was treated for rheumatism, without relief. In September of the same year his hand had become so weak that he had to give up most of his work.

During all of this time, that is, from the beginning of his trouble, he has had a feeling of constriction about his body from the axilla to the crest of the pelvis, more marked on the right side, and seeming to come on in attacks. At times he would seem to be free from this feeling entirely. About five months ago, in December, 1904, he noticed that his legs were

8. "Neuroma and Neurofibroma," Alexis Thompson.

becoming weak and he began to have pain in his toes and from the knees to the hips. This pain he describes as a "dead" or "numb pain." This was more noticeable on the right side. About this time the right hand and arm had become so weak that he was unable to use them at all. For the past four months the weakness in all his limbs has increased and the pain has become more severe, being confined largely to the extremities and being worse at night. He has been losing weight, and for about the last six months has had some difficulty in starting his water.

Examination.—Movements of the facial muscles normal; closes eyelids firmly, wrinkles forehead; shows teeth well, masseters contract firmly, tongue protruded straight, not atrophied, no fibrillary tumors. Extra-ocular muscles normal. Sensations for touch and pain in the face normal.

Upper Limbs: Movements much impaired; resistance to passive movement weak; no local atrophy anywhere, though the man is generally wasted and probably more so in the forearms than in the upper arms. Grasp of each hand is very feeble. Biceps tendon reflex, triceps tendon reflex very prompt on each side. Sensations for touch and pain are preserved in each upper limb, even in the hands, but these sensations are not so sharp as in the face. A small tumor is found in front of the left elbow about the size of a hazelnut, and is painful to touch. Right upper limb is weaker than the left.

Lower Limbs: Voluntary movement and resistance to passive movements weak in lower limbs. Sensations for touch and pain are preserved in the lower limbs, but are probably more impaired in the right lower limb than they are in the face. Patellar reflex is much exaggerated on each side. Ankle clonus present on each side, more distinct on the right. Babinski reflex is distinct and typical on each side. Sensations to touch and pain may be a little more acute in the left limb.

There is no deformity of the vertebral column. No tenderness to pressure along the spinal column. Patient can not feed nor dress himself. Cremaster reflex is present, but feeble on each side. He sways a little with eyes closed. Gait is somewhat staggering and he drags the right toes on the floor like in a hemiplegia. In walking, the right upper limb is hanging slightly flexed at the elbow. A small tumor is seen between the first and second toe on the right foot, not painful to the touch. Later other small painful tumors were found in the limbs.

Diagnosis.—Neurofibromatosis, probably with pressure on the cord by tumor of the nerve roots.

TYPES OF CONGENITAL SYMBOL AMBLYOPIA.*

J. HERBERT CLAIBORNE, M.D.
NEW YORK.

W. Pringle Morgan,¹ of Seaford, Sussex, England, was the first to point out that, in addition to the word-blindness which is the result of acquired cerebral lesion, there is a word-blindness congenital in nature which he has suggested is due to defective development of the word memory centers of the cerebral cortex.

This communication was followed by those of Hinshelwood² and Nettleship,³ Wernicke,⁴ of Buenos Ayres, and others⁵ the experience of all who have observed and studied these cases simply justifies the initial and simple conclusions of Morgan. The first American observer

who has recorded any observations on this subject is A. Schapringer, of New York, who read a paper on the subject before the Section of Ophthalmology of the New York Academy of Medicine, Feb. 19, 1906. His paper on that occasion was followed by one by myself in which I referred to two cases. That paper is the basis of this one.

It was my purpose to present an extended study of two boys whom I have recently seen and who have shown themselves so backward in learning to read that at the first blush it excited in my mind a suspicion of congenital word-blindness, owing to the shyness of both of them, I have not been able to study their cases as carefully as I wish, and am compelled to present my views after a more or less imperfect investigation. I do not doubt, however, that in these two boys there is a congenital deficiency, or at least a tardy development of the word-memory centers; and the class of cases to which they belong is to be clearly and definitely differentiated from those cases of word-blindness which occur in cerebral degeneration, intracranial tumors or lesions and injuries. Congenital word-blindness appears to me to be a department of pediatrics and neurology which has not received the attention it deserves.

CASE 1.—A boy, aged 10, was sent by his teacher to my clinic at the New Amsterdam Eye and Ear Hospital on account of an injury to his left eye. In my opinion, the injury has no bearings on the symptoms which I have observed, and it is sufficient to say that he received a blow from a small shot that was thrown or fired at him by a comrade. He said the shot lodged under his upper lid and fell out on examination of it by his sister.

Examination.—He has a vitreous opacity, a ruptured iris, and greatly diminished vision in his injured eye. It was, however, fairly free from irritation.

While examining him I was struck by his slowness and peculiarity, and asked his mother what his status at school was. She said he had never yet been able to learn how to read, but could write a few simple words. I tested him on his alphabet. At times he named some letters correctly, and later on failed to name the same ones correctly. I then asked him to make several letters of the alphabet, when he made repeated errors. When asked to write capital K he wrote capital T; for capital T he wrote little t, and he could not write u at all. Although he had written a K for a T, when asked to write K he could not make it, but when asked to write his name he did so with readiness. His name is James Kennedy and he did not halt at the K. I believe his name was written automatically. The letter E seemed to have great difficulty for him. He called it S at times, at other times correctly, and finally returned to calling it S. He wrote his name James correctly but he could not call the letter E after he had written it. He kept calling it S, although it was pointed out to him and he himself pointed it out clearly. His refraction was about emmetropic but I could get no accurate test of vision since he was never certain about his letters. His field of vision was normal, and he was right-handed. In other respects he was fairly bright, seemed to understand words and commands, knew the meaning of ordinary words, recognized objects and their uses, and according to his mother was talkative, communicative, even garrulous at times, and played with other boys in the street in a normal manner.

I had a letter from his teacher, and she sent me a few tests which she made of his writing at dictation and at copying. He writes very well, particularly his name, and I repeat that this act appeared to me when I examined him to be done automatically. As stated, he was right-handed and evidently copied correctly.

I myself made some tests of his spontaneous writing, which I have unfortunately lost, and which exhibited the earmarks of classical motor aphasia, although it must be remembered that motor aphasics, as a rule,

* Read in the Section on Diseases of Children of the American Medical Association at the Fifty-seventh Annual Session, June, 1906.

1. Morgan, W. Pringle: Brit. Med. Jour., Nov. 1896.

2. Hinshelwood, James: Lancet, 1900, vol. 1, p. 1506; also Ophthalmic Review, March, 1901.

3. Nettleship, E.: Ophthalmic Review, March, 1901.

4. Wernicke, Otto (Buenos Ayres: "Angeborene Wortblindheit," Centralbl. f. pr. August, 1903, p. 264).

5. Thomas (London) The Ophthalmoscope, August, 1905, (reports 8 cases); also Stephenson, Sidney: "Congenital Word-Blindness," Lancet, September 17, 1904, referred to in Centralbl. f. pr. Augenh., 1906, p. 59, in the abstract of Thomas' article; Lechner, C. S.: Weekblad voor Geneeskuunde, August 1, 1904.