J. Warren Howard, M.D., died at his home in St. Paul, Ind., August 29, from heart disease, aged 63.

Thomas C. Osborn, M.D., died at his home in Cleburne, Texas, August 20, aged 85.

Queries and Minor Notes.

ANONYMOUS COMMUNICATIONS will not be noticed. Queries for this column must be accompanied by the writer's name and address, but the request of the writer not to publish his name will be faithfully observed.

COINCIDENCE—ITS PROPER VALUE IN DIAGNOSIS.

St. Louis, Aug. 25, 1902.

To the Editor:—I think it would serve a useful purpose if you invited contributions to your columns concerning coincidences, and to express more clearly what I mean I will cite two cases from my own practice:

A man came to me complaining of pain in the right testis of two days' duration which he attributed to the testes having been suddenly squeezed between the left thigh and a bunch of keys in his right trouser pocket in the act of crossing his legs. No pain had ever been experienced in the part before. Examination disclosed advanced tubercular epididymitis.

A boy was brought to me supposed to be suffering from nervous shock, having been knocked over some days previously by a drunken man in the street. Routine examination commencing with the throat discovered diphtheria.

Here is a factor, then, which, as far as I know, has never been touched on in the treatises on diagnosis—coincidence. There is no escape from being obliged to give it due consideration in our mental summing-up of the case unless we examine our patients from a veterinary point of view—perhaps a better way—finding out what we can by our own unaided industry. The consideration of the particular value to be attached to coincidence would lead me further than it is my intention to trespass on your valuable space. Recitation of cases from others would not only be highly instructive but also most interesting and would furnish a moral to adorn many a clinical tale.

Yours faithfully,

J. R. CLEMENS.

ETHICS IN SUCCEEDING ANOTHER PHYSICIAN.

Lawrence, Mass., Aug. 28, 1902.

To the Editor:—Kindly decide this question of ethics, and oblige two members of the profession in good standing and others who may meet with a parallel case: A physician has been in attendance on a patient ten days (after miscarriage) and has made morning visit and is to call during afternoon. Meanwhile the family desires to dispense with his services—he has been paid in full for services rendered—they telephone for another physician who has no knowledge of the circumstances until he arrives at the bedside. When he is informed that a fellow practitioner is expected to call he refuses to accept the case or give an opinion and rises to depart. The patient and her husband state that they desire his services and shall not further employ the other physician. He explains his position and ends by refusing to have anything to do with the case unless in consultation with the other physician. The family refuse to have the other physician. "Then, you must notify your attending physician not to call any more," said the second physician, "before I shall be at liberty to attend this patient." The husband goes to the physician's office and leaves notice that his services are no longer required. The husband returns and states that he was unable to see the physician personally but left word at his office for him not to call. Meanwhile, the patient suffers considerable pain. The second then accepts the case, makes examination, gives opinion and outlines treatment. He in no way solicits patronage, and neither criticised nor commented on the previous conduct of the case. Has he (the second physician) broken a rule of ethics or been guilty of conduct in any way discourteous to his fellow practitioner? Very truly yours,

JOHN T. CAHILL.

Ans.—As stated there seems to be no violation whatever of the Code of Ethics. See Art. V, Sec. 4.

MALARIA WITH ANASARCA.

Mosher, Ark., Aug. 27, 1902.

To the Editor:—Since locating in Arkansas I have encountered a peculiar condition that the text-books fail to describe. It is an anasarca associated with or dependent on acute malarial infection. It commences with a chill and is followed by fever and sweating. About the second or third day the swelling begins and involves the whole body. The fever is now slight or wanting. The thoracic and abdominal viscera are apparently normal except a slightly enlarged spleen. Urine analysis: sp. gr. 1020, acid, amber, no sugar,

no albumin, no casts. Pot. ferio, cyanid, heat and nitric acid tests for albumin were carefully used. Blood examination revealed many plasmodia malaria—crescents in all cases. After giving quinin for a week or nine days a speedy convalescence ensued. The patients ranged in age from four to fifty-two years and none had had malaria for years. I would be gratified for any information as to where I can find literature on the disease or of anyone with like experience.

F. L. MORSE, M.D.

WOUNDS OF THE ABDOMEN.

ALBERT, W. VA., Sept. 4, 1902.

To the Editor:—Kindly answer the following questions through THE JOURNAL: 1. Given a patient with one or several stab wounds in the abdomen, what, in your opinion, is the very shortest time in which death could result from peritonitis? 2. What is the latest treatment advised for gunshot wounds of the abdomen—wounds inflicted with the ordinary weapons?

A. P. BUTT.

Ans.—1. Shock and hemorrhage can not be excluded, but when they are not dominant factors, peritonitis may result in death in 24 to 30 hours. 2. For the expert: if there is presumptive evidence of involvement of abdominal viscera, immediate operation, repair of damage and washing out of the abdominal cavity. For the ordinary practitioner: if there is strong evidence of visceral involvement, immediate operation is indicated; if not, expectant treatment.

Correspondence.

Concerning Indexes of Medical Journals.

Dillon, Mont., Sept. 4, 1902.

To the Editor:—Regarding the editorial note "Concerning Indexes of Medical Journals" in THE JOURNAL of August 30: The index furnished by THE JOURNAL is indeed a valuable addition to our comfort, but even such an index can not attain perfection, from its necessarily discontinuous publication.

I would suggest my plan of using card index. It is this: Soon after receipt and reading of each periodical, its contents are indexed on cards and a check mark placed on the journal to show that it is ready to be filed away until the volume is complete. Each periodical is given a number, which accompanies the volume numbers, for example: THE JOURNAL A. M. A. is No. 1. Then an article appearing in vol 39, page 496, on "Indexes of Medical Journals" is placed on the card devoted to "Indexing" as "1, 39-496, medical journals," meaning: THE JOURNAL A. M. A., vol. 39, page 496, indexing of medical journals.

I subscribe to only six journals, but in six years this amounts to 72 volumes and everything considered worth preserving is indexed at one place (in two drawers of my office desk). Thus, in a very few moments I am able to find every article on a given subject in any of the 72 volumes, and without rising from my desk. The work of indexing amounts to only a few minutes each week and the short time spent on it I consider most profitably employed.

M. A. WALKER, M.D.

Priority in Perineum Repair Technic.

ATLANTA, GA., Aug. 19, 1902.

To the Editor:—In THE JOURNAL of the 16th inst. is a communication from Dr. Marcy of Boston in which he calls attention to these articles of his, viz.: Transactions of the American Association of Obstetricians and Gynecologists, September 1888, "The Perineum; Its Anatomy, Physiology and Methods of Restoration After Injury"; also "The Reconstruction of the Pelvic Structures in Woman," American Association of Obstetricians and Gynecologists, 1892; also "The Reconstruction of the Pelvic Structures Incident to Lesions of Perineum," THE JOURNAL American Medical Association, October 21, 1899. He then says: "Here will be found all the essentials of Dr. Noble's interesting article on 'A New Operation for Complete Laceration of the Perineum.'"

In reply to this most extraordinary claim I will state that the chief feature of my operation is *elimination of the danger of infection from the rectum* by dissecting the lower anterior portion of the rectum free from the vagina and drawing the uninjured portion of the anterior wall of the rectum down through and external to the anus, making an unbroken septum between the rectum and vagina, virtually converting the complete tear into an incomplete laceration. Dr. Marcy does not attempt to do this; his operation is the Tait operation modified