A low tracheotomy was done and an ordinary large canula was introduced and secured with tape. The chloroform was then administered through a rubber tube connected with chloroform apparatus and the canula. The thyroid cartilage was then split (with difficulty owing to calcification) in the median line with bone forceps and the larynx laid wide open and examined with a strong head light. The interior of the larynx was then painted with cocain and adrenalin. The disease was found to be confined to the left half. The left half of the thyroid and cricoid cartilages were then removed by freeing the surrounding tissues and pushing aside the perichondrium. Great care was necessary in separating the laryngeal tissues from the esophagus. Several strips of iodoform gauze were placed in the laryngeal cavity and brought out through the lower part of the wound and secured. The upper part of the larynx was closed with sutures and the dressings placed.

Later History.—The subsequent treatment required close and careful attention. The tracheal tube was kept in two days. It was several days before fluids or solids could be swallowed without passing out the wound, so he was nourished through a tube. He made a good recovery, and now, at the end of four years, is 76 years old, in good health and able to earn his living (Fig. 1).

CASE 2.—Patient.—J. S. F., age 52, contractor, had an excellent tenor voice and often sang in choirs. He used tobacco excessively and at one period of his life used alcohol. His father, a soldier, died of dysentery during the war, at the age of 40; his mother died at 65 from an accidental fall. His six brothers and sisters were all living and healthy. There was no history of malignant disease or syphilis. The patient was an excellent specimen of manhood and had always enjoyed good health with the exception of chronic bronchitis.

Course of Disease.—A persistent hoarseness set in during the spring of 1904, following an acute laryngitis. During this attack a small papillomatous-looking growth appeared growing from below the cords in the anterior commissure. During the months of May and June applications of argyrol were made to the larynx with other local treatment and a saturated solution of the iodids administered internally. The progress of the growth was daily watched until about the first of July when sections were removed and microscopically examined. Each specimen was reported to be unmistakably a flat-celled carcinoma. After these examinations the patient took several weeks rest in the woods of Wisconsin. When he returned I advised an immediate and radical operation, and at that time a thyrotomy would have been, in my opinion, sufficient. After this advice (September 1) the patient was not seen again until December 20, when he called at my office and said that he was ready to have the radical operation performed, if I thought best. In the meantime, he said, he had consulted Dr. E. Fletcher Ingals of Chicago, who said that the growth was suspicious but that he hoped it was only a large papilloma. He removed the tumor intralaryngeally, and sent the specimen to Professor Hektoen for examination. His report was: "The specimen which you sent me is a flat-celled carcinoma." This was followed by local treatment for a few days and then the patient was permitted to return home. He again visited Dr. Ingals two months later, who found that the growth had returned. The patient was then advised to undergo a radical operation. After he had given me this history, an examination of the larynx was made and it was found that the disease had made rapid progress, but was as yet apparently confined to the left half of the larynx.

Operation.—After explaining to the patient his chances for recovery, and that a complete laryngectomy would probably be necessary, it was decided to have the operation performed at once, Dec. 26, 1905, at the Deaconess Hospital. The teeth and upper respiratory tract were rendered as nearly aseptic as possible. The trachea was prepared by frequent injections of antiseptic oils and sprays. The patient was placed on the table in the Trendelenburg-Rose position and the chloroform was administered first by mouth and then by tube and canula. The different steps of the operation were about the same as detailed in Case 1. Extensive removal, including a part of the right thyroid cartilage and vocal cord, in addition to the entire left half of the larynx and the adjacent lymphatic

glands, was found necessary. In separating the cartilages from the esophagus, the esophageal tube was introduced and found to be of decided help.

Later History.—In the after-treatment of this case there were a number of complications. The old bronchitis was troublesome and at one time pneumonia was threatened. A very rapid pulse and irregular and disturbed respiration, due probably to interference of functions of the pneumogastric, were finally corrected with digitalis and strychnia. After the wound healed the x-ray was used with apparent good effect in preventing a return. Three and one-half years have elapsed, the allotted time, the laryngeal cavity looks exceedingly healthy, and the patient has been perfectly well since. (Fig. 2).

The Hahn canula was not found necessary in either of these cases.

30 Willoughby Building.

DOUBLE-HEADED MONSTROSITY.

GEORGE E. OREBAUGH, M.D.

HIGHLAND, OHIO.

Parents.—Mrs. C. L. M., aged 25, weight 108 pounds; had been in good health for several years, except for a leucorrhœal discharge. She is perfectly developed. Her husband has no bad habits and no deformities. The family history on both sides is good, there being no abnormal births or twins in either family.

History.—I was called to see the woman during the latter months of pregnancy and found her suffering from persistent vomiting and from intense pain in the right side. On March 6, 1900, I was again called to see her. After protracted labor she gave birth to a female child which died in a few minutes. The breech presented, making the delivery difficult, and in addition the placenta was adherent. The mother is now in good health and has since given birth to a normal child.

Description of Child.—The child had one body, perfect in form from shoulders to feet. There were two legs and two arms, three spinal columns and two perfectly formed heads and necks, each separate and distinct. Each head measured fourteen inches in circumference. Other measurements were as follows: Across the shoulders, 8 inches; around thorax, 15½ inches; length of arm, 8½ inches; length of legs, 11 inches; length of body, 20 inches; weight, 14 pounds.

A CASE OF SUPPURATIVE PYELONEPHRITIS, WITH SPECIMEN.

G. SHEARMAN PETERKIN, M.D.

SEATTLE, WASH.

This case and specimen is presented as graphically illustrating possible sequelæ of gonorrhea and teaching (1) the necessity of care in pronouncing gonorrhea cured, i. e., that there will be no sequelæ; (2) the necessity of gentleness in genitourinary surgery; (3) the rapidity with which infection in the genitourinary tract may prove fatal.

The specimen was obtained in the following case:

History.—M., male, aged 36, occupation, logger; had gonorrhea in 1895. During the last two years he had had some trouble in voiding urine, but only when drinking heavily. At this time he had also a slight urethral discharge; otherwise, no symptoms. He was never incapacitated from work. On June 6, 1907, he had been drinking heavily, but had attended to his duties. On this date he had acute retention and the sound was used to relieve same. Twenty-four hours later, June 7, he came to Seattle and presented himself for treatment at the local hospital. His condition was referred to the visiting physician as that of a case of gonorrhea, with retention, complicated by orchitis because of the swollen scrotum. He was catheterized with difficulty and two or three ounces of bloody urine obtained; catheterization was repeated after four hours. Twenty-four hours later the case was referred to me.

Examination.—There was evidence of commencing uremic coma; extravasation of urine, extending over the perineum, scrotum and half way up to umbilicus.

Operation.—A suprapubic cystotomy was done to relieve the retention. No attempt was made toward diagnosis. One quart of reddish-black, grumous, ammoniacal urine was evacuated and the bladder mucous membrane came away in large shreds. The patient died in eight hours without recovering consciousness.

Autopsy.—Postmortem findings were as follows: At the bulbous portion of urethra, was a stricture (Fig. 1, A). Along side of this, but slightly posterior, was found a urethral calculus (Fig. 1, B); size, $\frac{5}{8}$ by $\frac{5}{16}$ inches, or 1.5 by 8.5 cm.; weight, 8.5 gms., or a little over 2 drams; composition, phosphatic calcium magnesium. A cavity was plainly visible where the stone in the urethra lodged (Fig. 1, C), and a piece of catheter has been passed through the opening caused by the stone (Fig. 1, D). The prostate was found to be the seat of

Fig. 1.—The urethra laid open; A, point of stricture; B, urethral calculus; C, cavity in which calculus lodged; D, piece of catheter passed through opening caused by stone; E, seat of abscess in prostate; F, bladder denuded of mucous membrane.

an abscess (Fig. 1, E) and the bladder was absolutely denuded of all mucous membrane (Fig. 1, F).

Left Kidney.—Macroscopic Findings: Size, 6 by 10 cm. ($2\frac{1}{4}$ by 4 in.); weight, 139.3 gms. ($4\frac{1}{2}$ oz.); surface irregular, lobulated and knotty; capsule, thin and tight; strips with ease; in cortex, near surface, abscess (Fig. 2A), 1 by 1.5 cm. ($\frac{3}{8}$ by $\frac{5}{8}$ in.) in size, filled with thick cheesy pus. Pelvis and calices enlarged; membrane, thick and rough; ureter, distended and thickened, coated with flakes of fibrin.

Microscopic Findings: Micro-organisms *Bacillus coli communis*, *Staphylococcus pyogenes albus*. Cortex, glomeruli congested, showing marked diapedesis of the red blood cells, a thinning distention and rupture of the capsular epithelium and membrane; in other places, the epithelial cells are lifted and separated from the basement membrane. The appearance of the tufts is irregular, having lost their definition and con-

volute arrangement. In the areas of interstitial degeneration, the tufts are either partially or totally destroyed. The convoluted and straight tubules show a mild degree of swelling, with a paling and disappearing of the nuclei; they give the appearance of moderate amyloid degeneration. The lumens of the tubules are not distended, but flat and normal in appearance; they contain no casts or moulds. The interstitial tissue, shows perhaps, the greatest change of all. There is present a

Fig. 2.—Left kidney split open showing an abscess (A) in cortex.

Fig. 3.—Left kidney showing abscesses A, B, C, D, seen only from the external surface.

marked active congestion, leucocytes often predominating the field. In other places, there is a wide and free dissection of the interstitial tissue, with rupture of glomeruli and tubules; many of these are communication cyst cavities. Between the tubules are small collections of pus. The dissection has separated the straight tubules and extends clear to the renal capsule—in many places resembling the appearance of emphysematous lung tissue, being a fine, irregular network of stroma tissue, creating spaces unlined with epithelium or cells. In Figure 3—left kidney—the white pieces of paper show other abscesses seen only from external surface.

Right Kidney.—Macroscopic Findings: Size, 8 by 11 cm. (3 by 4½ in.); weight, 217.70 gm. (7 ounces); heart-shaped; surface, slightly lobulated, smooth, capsule, adherent; cortex, 1.5 to 2.5 cm. (½ to 1 in.) thick, firm, no abscesses; pelvis large, irregular; ureter, large thickened walls; pelvis dilated, mucous membrane thickened, congested, and coated with patches of fibrin.

Microscopic Findings.—These show a similar picture to that of the left kidney, with the active congestion well marked and hemorrhagic infiltration of the tufts. The blood vessels show no characteristic changes, except the venous system, which is slightly dilated, with distortion of the capillaries.

TWO ADENOMATA OF THE ISLANDS OF LANGERHANS.

MARY ELIZABETH MORSE, M.D.

Pathologist to the New England Hospital for Women and Children.
BOSTON.

Adenomata of the islands of Langerhans have seldom been described in the literature, probably because they are accidental findings in the routine examination of autopsy material rather than because they are actually very rare. Nichols,¹ in 1902, reported one which he believed was the first of the kind on record, and Helmholtz,² in 1907, added a second case. The reader is referred to their articles for a discussion of several other tumors described as arising from the islands of Langerhans, but apparently of doubtful origin. A search through the literature reveals no additional cases. Recently, however, two tumors of this kind have come under my observation. The first was found at the New England Hospital in the service of Dr. Clara J. Alexander; the second at the Boston City Hospital by Dr. O. R. Mabee, through whose courtesy I am enabled to report it. Both were discovered at autopsy, and have no bearing on the clinical histories or the other anatomic changes.

The first growth was from the pancreas of a woman, aged 44, dying of peritonitis. The gland appeared normal macroscopically, and the tumor was found in the specimen from the splenic end saved for microscopic examination.

Examination of First Tumor.—It forms a sharply defined encapsulated nodule, 5 mm. (¼ in.) in diameter, situated in the midst of the pancreatic tissue. It is composed of interlacing strands of cells supported by a connective tissue framework carrying blood vessels. The columns are in general two or three cells broad, but they often spread out into irregular masses. The cells are uniform in appearance throughout the tumor. They are indistinct in outline, polyhedral or cuboid in form, with a large vesicular nucleus and a moderate amount of finely granular cytoplasm,

which stains faintly with eosin. The nucleus is usually centrally situated, and contains one or more prominent chromatin masses. In short, the cells correspond in appearance and arrangement to those of the islands of Langerhans. No mitotic figures are found. The connective tissue framework is rather heavy. The blood vessels are dilated, frequently forming large spaces. They are intimately related to the cell columns, the latter resting directly on the capillary walls. The capsule is composed of coarse fibrillated connective tissue. Groups of tumor cells are included in it at several points, but do not penetrate it.

Islands of Langerhans are abundant in the pancreatic tissue surrounding the tumor. Some of them are unusually large, but otherwise they show nothing noteworthy. The pancreas contains a slight excess of connective tissue. The acini are small, the gland cells are shrunken, stain deeply and show no signs of secretion. The entire pancreas was cut in cross section at intervals of about 5 mm. (¼ in.), but no other tumor nodules were seen. Sections from various parts of the gland show numerous enlarged islands, some three or four times the usual size.

The second adenoma, which was found in the pancreas of a man, aged 46, dying of cerebral hemorrhage, arteriosclerosis and chronic nephritis, closely resembled the first.

Examination of Second Tumor.—It forms an encapsulated nodule 3 mm. (⅛ in.) in diameter. It is, however, more cellular than the first tumor, and the cells show greater variation in size and appearance. Some are three or four times as large as others; still others have a coarsely reticulated cytoplasm which scarcely stains.

The relation of adenomata of the islands of Langerhans to malignant tumors of the pancreas is unknown; but it seems doubtful whether they have any importance

Adenoma of the islands of Langerhans. The tumor here shown is 3 mm. (about ¼ inch) in diameter and situated in the midst of pancreatic tissue. The tumor cells are indistinct in outline, polyhedral or cuboid in form and each contains a large vesicular nucleus and finely granular cytoplasm. The capsule is composed of coarse fibrillated connective tissue.

1. Jour. Med. Research, May, 1902, iii, 385.
2. Johns Hopkins Hosp. Bull., 1907, xviii