of Vienna, who has contributed a short historical preface. Mr. Steinhäuser, who is himself the author of several logarithmic tables, encouraged Mr. Blater in his work, and rendered him great assistance throughout. The actual calculation occupied eighteen months.

With respect to the general employment of Mr. Blater's table for the performance of multiplications, is to be feared that its utility has been jeopardized by the size of page adopted. Any one who has occasion to make constant use of tables knows the enormous advantage of the octavo form over the quarto. The book is placed to the left of the computer, and the effort of carrying by the eye a series of figures from the left hand page of a quarto table to the paper—a distance of 18 in. to 2 ft.—is but ill compensated for by large figures or fewer pages. Handsome as the book is to look at, it seems to us that the table would have had much more chance of bringing the method into general use if it had been of octavo form. It is greatly to be regretted that it was not printed on 400 octavo instead of 200 quarto pages, which would have been quite possible with the existing arrangement of the table. If this had been done, and if the type had been somewhat smaller, a neat and handy volume might have been produced.

The mode of entering the table is very insufficiently explained in the introduction. This is unfortunate, as the mode of entry by the last figures is so unusual in tables that it should have been explicitly mentioned. Also the translation into English is so very unsatisfactory as to be obscure in places. These, however, are minor blemishes which would have but slight influence on the general utility of the table, if only the form were convenient.

The question of how far the method of quarter squares is likely to come into use is of some interest. Hitherto the method has been very little known, and so far as we know, it has never been used in practice on any extended scale. The mere fact that it has been so constantly discovered anew is sufficient evidence of the slight attention that it has received. Still there ought to be room for a table that gives, to the last figure, the products of any two five-figure numbers with only two entries. A seven-figure table of logarithms is inadequate for this purpose, for, besides requiring three entries, it only gives the first seven figures of the result.

On the other hand, it may be said that in ordinary calculations seven figures are as many as are required, and that logarithms possess the advantage of being equally convenient for divisions and multiplications. It must be admitted that a five-figure quarter square table is appropriate to only a limited class of calculations: it applies only to multiplications, and the number of figures in each of the two numbers must not be greater than five. These conditions are of a somewhat special kind. In recent years when heavy multiplications have been required it has become the custom to make use of Thomas de Colmar's arithmometer; and probably, at the present time, nearly all systematic work of this character is carried out either by Crelle's tables or by the arithmometer.

Passing now to the general question of multiplication by means of a table of single entry, we have the two methods of quarter squares and logarithms, each possessing its special advantages. There is also an older method which passed out of notice with the invention of logarithms. This method was called "prosthaphæresis," and depended on the formula

$$\sin a \sin b = \frac{1}{2} [\sin \{90^\circ - (a-b)\} - \sin \{90^\circ - (a+b)\}]$$

A table of natural sines could therefore be used as a multiplication table, four entries being required. This method is due to Wittich, of Breslau, who was assistant for a short time to Tycho Brahe, and it was used by them in their calculations in 1582. It is referred to by Raymarus Ursus, Clavius, and Longomontanus; and it seems to have been used for performing multiplications not only when the numbers occurred as sines, but also in the case of ordinary numbers.

The method of quarter squares depends upon so simple a formula that it is strange that the first table should not have appeared until 1817. There seems no reason why it should not have been employed before the invention of logarithms, when it would have been a most valuable aid to calculation. The geometrical theorem, which is equivalent to the algebraical identity $(a+b)^2 - (a-b)^2 = 4ab$, on which the method depends, forms Prop. viii. of the second book of Euclid; and one would think that the application of the geometrical or algebraical theorem to arithmetic might have been noticed at any time. The actual history of mathematical tables is, however, entirely different from what we might expect it to have been, owing to the wonderfully early invention of logarithms: and it was, in fact, only just about that time that the importance of tables as an aid to general calculation was beginning to be felt. The date of Herwart ab Hohenburg's great double-entry multiplication table, extending to $1,000 \times 1,000$ (the same limit as Crelle's table, and which has never been exceeded), is only four years earlier (1610) than that of Napier's "Canon Mirificus" (1614).

It is interesting to notice that the method of quarter squares is more closely connected mathematically with the method of prosthaphæresis than with that of logarithms; in fact, from the formula

$$\sin a \sin b = \frac{1}{2} \{ \cos(a-b) - \cos(a+b) \}$$

we readily deduce

$$ab = \frac{1}{4} \{ (a+b)^2 - (a-b)^2 \}$$

by expanding the sines and cosines in ascending powers of their arguments and equating the terms of two dimensions.

The method of quarter squares enables us to multiply together two numbers of n figures each if we have a table extending to 2×10^n . If the latter only extends to 10^n three entries are required, and the final result has to be doubled whenever the sum of the numbers exceeds 10^n (as in Laundy's table). If we consider the question of the multiplication of two numbers of n figures each by means of a table extending only to 10^n , the same process being employed in all cases, it appears

that three entries are necessary, and that it would be better to tabulate half squares, using the formula

$$ab = \frac{1}{2}a^2 + \frac{1}{2}b^2 - \frac{1}{2}(a-b)^2.$$

In tabulating the half squares the fraction $\frac{1}{2}$ would be thrown off, so that if a and b were both uneven, unity would have to be added to the result.

It would, however, we think, if the table is not to go beyond 10^n , be more convenient to employ a table of triangular numbers. The n th triangular number is $\frac{1}{2}n(n+1)$, and if we are provided with a table extending to 10^n , we may multiply any two numbers not exceeding 10^n by means of the formula

$$ab = \frac{1}{2}(a-1)a + \frac{1}{2}b(b+1) - \frac{1}{2}(a-b-1)(a-b);$$

or as we may write it

$$ab = T(a-1) + T(b) - T(a-b-1),$$

$T(n)$ denoting the n th triangular number.*

Thus, to multiply two numbers we subtract unity from the larger number, and enter the table with the larger number so diminished, with the smaller number, and with the difference of these two numbers. For example, to multiply 5289 and 2156, we add the tabular results corresponding to 5288 and 2156, and subtract from this sum the tabular result corresponding to 3132.

The mode of construction of a table of triangular numbers is almost the simplest possible, the numbers being formed by adding to zero the natural numbers 1, 2, 3, . . . *e. g.*

$$0+1=1, 1+2=3, 3+3=6, 6+4=10, 10+5=15,$$

and so on. It may be noticed also that any two consecutive triangular numbers are the most nearly equal parts into which a square of points can be divided by a line parallel to the diagonal. For example, in the square of 16 points, the two most nearly equal triangular parts are $1+2+3=6$, and $1+2+3+4=10$. This is shown in the following diagram:

Whether the square contains an even or an uneven number of points, the diagonal, which is in the middle, has to be given to one of the two parts, which therefore necessarily differ by the number of points it contains. In the square n^2 , the two consecutive triangular numbers which form it are $\frac{1}{2}n(n-1)$ and $\frac{1}{2}n(n+1)$, differing, as they should, by n , the number of points in the diagonal. Viewing the same matter from a slightly different point of view, we see that any two consecutive triangular numbers always make a square, *e. g.*

$$1+3=4, 3+6=9, 6+10=16, \text{ etc.}$$

It is interesting to exhibit by means of a diagram the manner in which the rectangle representing the product ab is derived from the three triangular numbers corresponding to $a-1$, b , $a-1-b$. As an example, the mode of formation of the product 8×4 is shown below, the triangular number corresponding to 7 being represented by dots and the triangular number corresponding to 4 by stars.

The dots above the line form the triangular number corresponding to $7-4=3$.

It is not suggested that the method just described by means of triangular numbers is comparable to that of quarter squares. It is certainly better to double the extent of the table and have but two entries. Still, it is interesting to note how readily a table of triangular numbers extending only to 10^n is available for the performance of multiplications of two n -figure numbers. So far as we know, only one extended table of triangular numbers has ever been published. This table, which gives the value of $\frac{1}{2}n(n+1)$ from $n=1$ to $n=20,000$, was published at the Hague, by E. De Joncourt, in 1762, under the title "De Natura et Præclaro Usu Simplicissimæ Speciei Numerorum Trigonalium." The book is a small and handsomely printed volume of 267 pages, 224 of which are occupied by the table.

In tabulating quarter squares, the fraction $\frac{1}{4}$ which occurs when the square is uneven is omitted. If we

* It is interesting to compare the two formulæ which involve half squares and triangular numbers respectively. In the former case we tabulate a discontinuous function, and in the use of the formula a unit has sometimes to be arbitrarily added. In the latter case we tabulate a continuous function, and the formula always holds good (the larger of the arguments being always reduced by unity). One formula depends on squares, n^2 ; the other on factorials of the second order, $n(n-1)$.

† We might of course also perform the multiplication thus:

corresponding to the formula

$$ab = T(a) + T(b-1) - T(a-b).$$

But if unity is subtracted from the smaller, instead of from the larger, number, slightly higher numbers are involved in the process.

denote by $qsq\ n$ the tabulated quarter square of n , we have therefore

$$qsq\ (2n) = n^2, \\ qsq\ (2n+1) = n^2 + n.$$

A table of quarter squares may be formed by adding to zero the numbers 1, 1, 2, 2, 3, 3, . . . *e. g.* $0+1=1, 1+1=2, 2+2=4, 4+2=6, 6+3=9, 9+3=12$, and so on. Its construction, therefore, is very similar to that of a table of triangular numbers, the only difference being that the added numbers 1, 2, 3, . . . are each twice repeated. We may also regard the tabulated quarter squares as defined by this rule: The quarter square of n is equal, if n be even, to the sum of all the uneven numbers less than n , and if n be uneven, to the sum of all the even numbers less than n . For evidently the series $1+3+5+\dots+(2n-1)=n^2$, and the series $2+4+6+\dots+2n=n^2+n$.

By means of this definition of a quarter square we may exhibit the method of quarter squares diagrammatically as follows.

Taking as examples the products 8×3 and 7×4 , we have—

$$qsq\ 11 - qsq\ 5 = 8 \times 3,$$

which may be represented by

and

$$qsq\ 11 - qsq\ 3 = 7 \times 4,$$

which may be represented by

The number of points in the extreme left hand column of the difference of the quarter squares is always equal to the smaller of the numbers to be multiplied. If this number is uneven, there will be one middle line containing a number of points equal to the greater of the two numbers; the points in excess of this number are to be transferred from the line below the middle one to the line next above it, the excess from two lines below is to be transferred to make up the deficiency two lines above, and so on. If the smaller number is even, as in the second diagram, there are two middle lines differing from each other by two points; one point from the lower of these lines is to be transferred to the upper, three points from the line below the lower middle line to the line above the upper middle line, and so on.

It will be noticed that the tabulated quarter squares are, as it were, a species of triangular number in which the succeeding lines of points differ by two, instead of by one, as in ordinary triangular numbers (*i. e.* viewing the matter arithmetically, the quarter squares are derived alternately from the series $1+3+5+\dots$ and $2+4+6+\dots$, and the triangular numbers from the series $1+2+3+\dots$). It is the fact of the lines differing by two which enables us in all cases to adjust the points in the difference of two quarter squares so as to form a rectangle in the manner indicated above.

J. W. L. GLAISHER.

MAGNESIUM AS A REAGENT.

By H. N. WARREN, Research Analyst.

MAGNESIUM, both on account of its purity and at the same time the speed with which it facilitates reduction, either in the moist or dry way, causes it to rank high, if not the first, among reducing agents. Thus, for instance, on account of its purity from arsenic, no agent is, perhaps, better suited for the elimination of pure hydrogen when using Marsh's test for that substance. Secondly, its absence from iron renders it one of the safest reagents for the reduction of ferric salts previous to titration. The speed of reduction, at the same time, being about three times that of zinc, and giving practically no reaction with potassium ferri-yanide, the desired absence of both phosphorus and sulphur may also be relied upon. The ease with which it precipitates, from solutions, for instance, zinc, distinguishes it practically from all other metals, separating the same so perfectly that, if to a solution of zinc acetate containing also metals of the fourth group some pieces of magnesium be introduced, and the solution brought to a state of ebullition, the whole of the zinc is thus separated, so as to cause, after the introduction of the necessary reagents to retain in solution the magnesium salts, no precipitate on the addition of ammonium sulphide.

Several samples of zinc ashes, containing, as impurities, calcium, magnesium, and other carbonates, were quickly assayed by this method, and the zinc estimated in the metallic form.

Its general behavior with iron salts is somewhat more complex, but, strange to say, from a solution containing a chromic salt the iron may, with a slight experience, be entirely separated. Take, for example, a sample of chromic iron, to the acid solution of which has been added a sufficiency of sodium carbonate to replace the combined acid. To the precipitated carbonate is added acetic acid in slight excess; if, now, to the acetic acid solution thus formed about 1 grm. of magnesium is introduced, a somewhat violent reaction takes place, the ferric salt becoming almost instantly reduced, and the reaction thus terminating. On the application of heat to the solution the liquid now changes from the deep green characteristic of chromium chloride to a slight pink, thence to a red, violet, gradually to a pure green, thus assuming its original tint, during which time the iron contained in solution has been precipitated upon the magnesium in a somewhat coherent form, not unfrequently contaminated with oxide of the same. This is to be withdrawn from the solution containing the chromium, boiled with a few drops of HNO_3 , and tested for the presence of iron by

means of potassium sulphocyanide. As a reagent in the dry way magnesium reduces all metals with the exception of the alkalis and alkaline earths, and may possibly aid in the decomposition of the latter, provided a suitable combination of the same could be obtained.

As an exception to reduction in this line may be mentioned the powerful manner in which it attacks molybdic anhydride, when fused with the same, the combination being so intense as to be accompanied with loud detonations.

Even the refractory substances silica and boric

higher than the average low water, and five feet above the average high water of the Elbe; and are allowed to escape from time to time so as to wash out the "Geeststammsiel."

Three large boats lighted by torches lay at the underground landing, and these the excursionists entered. The Stxy-like weirdness of the trip was increased by the white cloaks and caps which the passengers wore over their ordinary clothing to protect it from dripping water. The boats glided noiselessly through the mysterious darkness, being propelled by short oars.

last arms of the system are not yet completed, but the work is being vigorously pushed forward.—*Illustrirte Zeitung*.

INTERNATIONAL PLACARDS AT THE EXPOSITION.

THE international placards which we have here brought together were one of the curiosities of the exposition, and, with their many-colored tints, enlivened the interminable palisades that took the place of

THE THIRTY-THREE NOTICES OF THE EXPOSITION RAILWAY.

anhydride may be also reduced when intensely heated in closed vessels with the same.—*Chem. News*.

A TRIP THROUGH THE SEWER IN HAMBURG.

WHILE Hamburg above ground was busy with its exhibition, trying to entertain the crowd of visitors and to win their praise, underground Hamburg wished to do its part, feeling that it also had something worth showing; so the exhibition committee and the department of buildings arranged a number of excursions which would afford an interesting view of part of the immense sewerage system of the city. The trips planned were to be on that branch known as the "Geeststammsiel."

The excursionists gathered at the Lombard Bridge, under which is the spacious "Spulkammer" (Rinsing Chamber). This they examined first. Here the waters of the Alster are dammed up at a level eleven feet

At regular intervals there were glimpses of the sunny upper world through the shafts, and then torch-lights could be discerned again at the sides of the aqueduct, held by mysterious figures that proved to be watchmen standing in niches formed in the wall of the conduit. Suddenly distant, reverberating and strange noises were heard; the gates in the "Spulkammer" had been opened, and the mass of water came thundering and rushing on; but it was some time before it reached the boats. First the foam could be seen at a distance in the half light, and then the large waves raised the boats and carried them on with great velocity. So the strange journey continued until the goal was reached, and gloomy darkness was left for the beautiful daylight. The length of the main conduit—through which the excursionists passed—from the Lombard Bridge to its opening into the Elbe, is nearly two miles; it is circular in cross section, and has a diameter of about ten feet.

The entire system, as completed in 1886, extended over 164 miles and cost more than \$4,500,000. The

horizon to the travelers on the Decauville railway. They were the piquant note of the trip and exercised the idiomatic faculties of the polyglot visitors. In the first place, at the beginning of the exposition, there was but one warning designed to prevent accidents, the track being necessarily closely hemmed in by the trees of the counter alleys of the Quai d'Orsay. This warning, which was at first placarded in French, was afterward translated into English, Italian and Spanish; and then, as the number of foreigners was constantly increasing, it was found necessary to add new placards reproducing the original notice in Russian, Danish, Swedish, Norwegian, Flemish, etc.

The public was amused at this text reproduced in fifteen different languages, and followed the extension of it with so much pleasure that it was deemed well to add new placards in Arabic, Roumanian, Greek, Turkish, and Chinese, and these were printed in characters so strange that it became necessary to indicate at the bottom what the idiom was that the odd vocabularies belonged to.—*Le Monde Illustré*.

A TRIP THROUGH THE SEWER AT HAMBURG.