This article was downloaded by: [New York University]

On: 28 April 2015, At: 00:29

Publisher: Taylor & Francis

Informa Ltd Registered in England and Wales Registered Number:
1072954 Registered office: Mortimer House, 37-41 Mortimer
Street, London W1T 3JH, UK

Annals and Magazine of Natural History: Series 5

Publication details, including instructions for authors and subscription information:
<http://www.tandfonline.com/loi/tnah11>

VII.—Note respecting butterflies confounded under the name of *Delias belladonna* of Fabricius

Arthur G. Butler

Published online: 07 Oct 2009.

To cite this article: Arthur G. Butler (1885) VII.—Note respecting butterflies confounded under the name of *Delias belladonna* of Fabricius , Annals and Magazine of Natural History: Series 5, 15:85, 57-58, DOI: [10.1080/00222938509459295](http://dx.doi.org/10.1080/00222938509459295)

To link to this article: <http://dx.doi.org/10.1080/00222938509459295>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of all the information (the "Content") contained in the publications on our platform. However, Taylor & Francis, our agents, and our licensors make no representations or warranties whatsoever as to the accuracy, completeness, or suitability for any purpose of the Content. Any opinions and views expressed in this publication are the opinions and views of the authors, and are not the views of or endorsed by Taylor & Francis. The accuracy of the Content should not be relied upon and should be independently verified with primary sources of information. Taylor and Francis shall not be liable for any losses, actions, claims, proceedings, demands, costs,

expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden.

Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

as it "mimics" another insect belonging to a different family and to another zoological region—the *Olesterus australis* of Gorham (Cleridæ), from Australia. The type of the genus, *C. methocoides*, Prof. Westwood compared to a genus of ants (*Methoca*).

Distenia Pryeri.

D. elongata, fusca, sat dense subtiliter albido-pubescent; capite inter oculos lineatim longitudinaliter impresso; antennis leviter pilosis; prothorace subtilissime confertim punctato, disco quadrituberculato, apice tubulato; elytris longe cuneatis, seriatim punctatis, punctis postice gradatim obsoletis, apice singulorum bispinoso, spina exteriori longiore; pedibus sparse pilosis, pilis longis adspersis; femoribus muticis. Long. 11 lin.

Hab. Ellopura.

In its uniform coloration this species is allied to *D. japonica**, but is at once differentiated by its two-spined elytra. The fine whitish pubescence on its brown derm makes a clear dark greyish colour. I have named this graceful Longicorn after Mr. H. Pryer, who is an observer as well as a collector.

VII.—*Note respecting Butterflies confounded under the name of Delias belladonna of Fabricius.* By ARTHUR G. BUTLER.

FOR some years past it has been maintained by most lepidopterists that *Papilio belladonna* of Fabricius, figured by Donovan in the 'Naturalist's Repository,' is the female of *Delias Horsfieldii* of Gray's 'Insects of Nepal;' on the other hand, I have always insisted that, however bad Donovan's figure might be, it represented a brown and not a black species, a male and not a female, and certainly a species in which the whole abdominal border of the secondaries was yellowish white, not partly white and partly yellow.

Amongst the Lepidoptera of the late Mr. Charles Horne, collected in the North-west Provinces of India, I found a single specimen of a *Delias* which, after comparison with Donovan's figure, I am satisfied represents the true *D. belladonna*; it is a brown, not a black insect, it is a male, not a female, and the abdominal border of the secondaries is wholly creamy

* *D. japonica*, Bates (1873), is said to be synonymous with *Apheles gracilis*, Blesig (1872), from Amur-land; the species should therefore stand as *D. gracilis* (Blesig).

white; the yellow spot at anal angle is also perfectly separate from the yellow commencement of the discal macular band, as in the figure by Donovan; the shape and markings of the primaries are quite as in that figure, and therefore quite dissimilar from *D. Horsfieldii* ♂.

I think therefore that we may conclude that *Delias belladonna* is a species confined to the North-west Provinces, that *D. Horsfieldii* is confined to Nepal, and that other forms allied to these may yet be expected to turn up. The two following have been confounded with *D. Horsfieldii* :—

Delias Hearseyi, sp. n.

♂. Nearly allied to the Nepalese *D. Horsfieldii*, slightly smaller; the yellow patch at base of secondaries duller and of a more orange colour; the yellow patch on abdominal area only represented by a dull yellowish nebula at the extremity of the white area, which is restricted owing to the anal angle being broadly black-bordered; the yellow spot near anal angle represented by a few white scales, whereas in *D. Horsfieldii* it is always present as a squamose yellow spot subconfluent with the abdominal patch: on the underside all the yellow spots are of a dull chrome-yellow, not bright gamboge, as in *D. Horsfieldii*; they are also rounded and narrower than in that species, so that they are in consequence smaller and further apart. Expanse of wings 79 millim.

Barrackpore (*Sir John Hearsey*). B.M.

Delias Boyleæ, sp. n.

♂. Form and size of *D. Horsfieldii*, but in the coloration of the wings much nearer to *D. ithiela*, the ordinary markings being represented by internervular grey streaks, upon which the submarginal spots of the primaries and three or four small spots on the disk of all the wings alone show white; base of secondaries almost brick-red (or dull orange); basal half of abdominal area grey, anal half bright chrome-yellow: below all the spots smaller and much more restricted than in *D. Horsfieldii*; all the yellow spots duller, chrome-yellow. Expanse of wings 84 millim.

Darjiling (*Mrs. R. V. Boyle*). B.M.

Females in this group seem to be very rare; of the four species here mentioned we only have male examples in the British-Museum series.