TRANSACTIONS
OF THE
NEW YORK SURGICAL SOCIETY.

Stated Meeting, November 27, 1907.

DR. WILLY MEYER in the Chair.

FUNCTIONAL RESULT AFTER AMPUTATION OF THE BREAST.

DR. ELLSWORTH ELIOT, JR., presented a woman 35 years old who was operated on June 26, 1906, at the Presbyterian Hospital for a carcinoma of the right breast. The history she gave on admission was that she had one child four years ago which she had nursed for 18 months. About six months ago she noticed a small lump in the right breast to the inner side of the nipple. This gradually increased in size, and as it was regarded as a fibroma, it was removed in the dispensary. Upon microscopic examination the growth proved to be a carcinoma, and the patient was sent to the hospital for a radical operation. This was done by the Meyer-Halsted method, and the patient left the hospital July 6, 1906. The functional result of the operation was practically perfect, all the usual movements of the arm being unimpaired.

SPLENECTOMY FOR RUPTURE OF SPLEEN WITH
SUBSEQUENT CHOLECYSTECTOMY.

DR. ELLSWORTH ELIOT, JR., presented a woman thirty-four years of age, who was one of two cases of subcutaneous rupture of the spleen that were reported by Dr. Eliot at the last meeting of the American Surgical Association, and published in detail in the *New York Medical Journal* for July 13, 1907. The history of the case was as follows:

Eight years ago patient was curetted for retained secundines; four years ago she was operated upon for ruptured ectopic gestation sac, in a condition of almost complete exsanguination. Since then she has been in excellent health.

On May 14, 1906, at ten o'clock, patient fell a distance of nine