lens, held in front of the lenses already chosen, which are placed in the trial frame. By this reversal of the axis we alternately increase and diminish the cylindrical effect of the combination before the eye. The lens with the round handle is made by Messrs. Bonschur & Holmes, and the lens with the flat handle is made by Messrs. Wall & Ochs, all of Philadelphia. The one with the round handle (see illustration) is the more easily turned; while the one with the flat handle can be more

readily kept exactly perpendicular to the visual axis. Either will be found of great practical value in refraction work.

DISCUSSION.

DR. W. F. SOUTHARD, San Francisco—Five years ago I devised a method of transillumination for the ophthalmometer, which I believe was the first effort in this direction. I first used ground glass, and gave that up because there was not a good distribution of light, and then I began to use porcelain [exhibiting the instrument]; a 16-candle power, 110-volt lamp is used. I did not think that it was obligatory to have it patented, but that any instrument-maker could be privileged to make it without danger of prosecution and such difficulties as I have experienced. I have testimonials from several manufacturers to show that this instrument was the first of the kind that they saw. One of the gentlemen spoke yesterday of the blurred edges of the mires as they approach each other. That is true and always will be, because it is due to an optical defect. I pay no attention to that myself, knowing that it is an optical illusion, and when at the point of contact there will be a little whitening more than we see here. I have demonstrated this instrument before a number of medical societies, and I desire the privilege of having my claim of priority published now in the transactions.

DR. A. EDWARD DAVIS, New York City—I may say that there was a transilluminated mire in use several years ago. Javal himself had one, but it was not so perfect an illumination as this.

DR. MILES STANDISH, Boston—I want to illustrate a little scheme I have adopted on my ophthalmometer, which aids me greatly in judging the point of contact. I have painted in the center of the solid white portion of the first step a little spot 1 mm. square, so that when the mires approach you have a point still beyond the light diffusion, and by watching this point you can judge the point of contact very accurately. It can be very easily accomplished by pasting a little black square on the mire, and once having placed it there you will never take it off again.

1. Ophthalmic Record, April, 1900.

SYMPTOMS AND DIAGNOSIS OF HYPERTROPHY OF THE PHARYNGEAL TONSIL.*

GEORGE MORGENTHAU, M. D.

CHICAGO.

Symptoms.—Hypertrophy of the pharyngeal tonsil is, in many cases, combined with enlargement of one or more other parts of Waldeyer's so-called lymphatic ring of the pharynx, and of the adenoid or lymphatic tissue found in all the mucous membranes. The symptoms produced by these abnormal conditions, while mainly traceable to the pharyngeal tonsil, may not be ascribed to it entirely, but must be considered in reference to a possible constitutional anomaly of which the adenoid vegetations are but a local manifestation. The most common of these is the excessive discharge which flows either into the nasopharynx or into the nose. It is of thick consistency, sometimes with a slight odor, and comes from Luschka's tonsil itself or from the nose. The tonsillar secretion in conjunction with the venous stasis, especially of the posterior ends of the lower turbinals, and the mechanical interference with drainage of the nasal chambers, by the enlarged tonsil, result in chronic rhinitis. This, in turn, is favored by various malformations found so often in patients with adenoid growths, be the relation causal or accidental. The growths being situated behind the choanæ, inspiration may be performed, while in expiration the growths fall against the nose, acting as valves. "Blowing the nose," is, therefore, quite inefficient. They are but seldom large enough to obstruct more than the upper part of the choanæ, so that the children can, although with some effort, breathe through the remaining lower part of the canal. But when this, too—as is most apt to be the case when the children are in a horizontal position—is either obstructed by the accumulating mucus or by the swelling of the posterior ends of the turbinals or even by the action of the erectile tissue in the tonsil, then they are forced to breathe through the mouth. Infants are sometimes unable to nurse at the breast. Bosworth states that acute rhinitis is comparatively rarely met with in a child. In most cases, when it has an apparent cold in the head, it is really suffering from a subacute inflammation of the pharyngeal tonsil. The nose is rendered still more inadequate for breathing by a condition to which Ingals drew attention, abnormal narrowness of the choanæ—a factor which accounts for some disappointing results after the growths have been removed as thoroughly as possible. Nosebleed sets in often, from small capillaries in the anterior lower angle of the septum or in the tonsil. If the blood can not escape, small petechiæ are to be seen in the submucous tissue, especially of the soft palate and the uvula, or in the tonsil itself. The discharge from the nose is apt to excite eczema at the introitus, thickening and roughness of the upper lip, etc., symptoms often attributed to scrofulosis. The chronic inflammatory condition of the nose is, in many cases, followed by impairment of smell and taste; frequently by conjunctivitis, occasionally by blepharitis, and even by keratitis. Or, as discharge gathers in the nasopharynx during the night, it may trickle into the stomach or be unconsciously swallowed, causing vomiting and gastric mischief, thus adding to the malnutrition of our little patients. The laryngeal and bronchial

*Read in a Symposium on Hypertrophy of the Pharyngeal Tonsil, before the Chicago Climatological and Laryngological Association, July 5, 1900.

symptoms, again, are either brought on by the descending mucus or by mouth-breathing. The most common symptoms are cough, hoarseness and false croup, and bronchitis and bronchopneumonia. While Bosworth states that, in laryngismus stridulus, there is acute swelling of the lymphoid tissue in the larynx proper, it does not seem unlikely that the mechanical irritation of the mucus might excite such attacks. Adenoid growths are often accompanied by nervous disorders. Caution is therefore necessary in drawing deductions from a comparison of the state of the patient before and after operation. The psychical effects of the operation itself might prove corrective of some of the nervous disturbances, or the greatly improved general condition of the patient after operation might be responsible for the beneficial change. In a word, the association of nervous predisposition and of lymphatic growths might be accidental, as the latter are of such common occurrence, especially in children. However, cures of asthma, chorea, even of epilepsy have been reported. Stammering and stuttering, not uncommonly met with in such cases, have been greatly benefited by the operation. Major ascribes enuresis nocturna to carbonic-acid gas poisoning from insufficient ventilation of the blood, the breathing being too superficial because of the stenosis. Of this, interrupted sleep and snoring are a further result. The best explanation is advanced by Fraenkel, that the base of the tongue falls backward on the epiglottis, thereby obstructing the larynx. The deformities of the chest are, according to some authors, caused by the necessity of breathing through the mouth, the respiration being superficial, with limited expansion of the chest. Fraenkel finds that in the chest, as in all stenoses of the upper air tract, the upper parts are emphysematous, the lower atelectatic. Trautmann and others speak of a paralytic form, predisposing the patient to tuberculosis. Talbot, however, considers the deformities of the chest—and of the face—not the result of adenoid growths, but, with them, signs of a general degeneration of the body combined with degeneracy of the mind. The children appear stunted in their physical growth, but it is common enough to see them increase in size and weight very soon after the operation. The peculiar facial appearance is produced by the broadening and flattening of the bridge and root of the nose, the narrowness of that organ, while the nasolabial folds are not well defined, the mouth is open, the eyes seem veiled. The dull expression is enhanced by the impairment of hearing so often present.

In marked cases, the jaw seems compressed from side to side; the hard palate is not flat but forms a high arch; the alveolar border is not rounded but comes to a point in front, is V-shaped; the teeth override each other; the upper incisors protrude from the mouth and beyond the lower incisors, which are usually normal; the teeth are more liable to diseases. Koerner distinguishes two forms according to the presence of adenoid growths before or after second dentition. In the first case, the palate is higher; in the second, the V-shape is characteristic, occurring only when there is nasal obstruction by adenoid growths. While the patient may appear to be stupid, he is not necessarily so. It is certain that such patients are often enough the brightest of children. They do, however, often complain of headaches and inability to concentrate the mind. Guye has termed the latter aprosexia. Whatever theory may be based on the anatomical connection between the nasopharynx and the brain, the symptoms are greatly alleviated by the removal of the pharyngeal tonsil.

Scanes describes a distinct transverse vein across the root of the nose. Schmidt relates that, in two children with continued elevation of temperature, the removal of adenoid growths did away with the fever. The importance of such an observation must be acknowledged in its bearing on a doubtful case of tuberculosis, etc. Most characteristic is the "adenoid" speech. The whole nasal cavity being excluded as a resonator, the voice sounds dead, different from that state when the nose is closed in front, or when enunciation becomes indistinct from enlargement of the faucial tonsil. Certain consonants are mispronounced; m becomes b, and n becomes d, etc. Lately, it has been shown by actual count that the blood of children with adenoid growths is impoverished and that, soon after their removal, it quickly approaches the standard averaged from examinations in normal children.

Thost asserts that symmetrical swelling of the numerous small glands of the superficial cervical plexus, in the lower triangle of the neck, behind the sternocleidomastoid, is characteristic of Luschka's tonsil.

Last, but not least, in this brief summary, are to be mentioned the effects of hypertrophy of the pharyngeal tonsil on the organ of hearing. While not as obvious and manifest as, for instance, the evils of mouth-breathing, there is hardly any condition which calls more imperatively for removal of the offending growths than the various affections of the ear whose cure is otherwise often impossible. Adenoid growths are the most prolific cause of certain diseases of the ear in childhood, excepting, perhaps, the infectious diseases. And these themselves seem to be less of a menace to that organ when the children are relieved of the growths. Simple, so-called, catarrhal otitis media, tubal stenosis, purulent otitis media, with its many sequelæ and complications, eczema of the external canal, etc., often owe their origin to inflammations which either start from the hypertrophied tonsil or are due to conditions dependent on its presence. Otology dates one of its greatest advances from the recognition of the supreme influence of nasopharyngeal and nasal diseases and abnormalities on the anatomically and physiologically intimately related hearing apparatus.

Diagnosis.—The combination of the characteristic facial appearance, of the manifold signs of nasal stenosis, of the voice—which, to be sure, is very similar in paralysis of the soft palate—combined with the aural symptoms, make the diagnosis possible almost at a glance. Sometimes the growths are abundant enough to hang below the soft palate and to become visible as soon as the tongue is depressed. The posterior wall of the pharynx may be studded with lymphoid follicles enlarged from the irritation of the secretion flowing from above. If the children be not too nervous or unwilling, they can be examined by posterior rhinoscopy. The growths can be divided into, practically, two varieties: Either there appears in the mirror a whitish, firm, semispherical cushion, resembling an enlarged faucial tonsil, lying between the openings of the Eustachian tubes, encroaching on the upper part of the septum; or real vegetations are seen, resembling stalactites, berries, etc., and filling the whole or parts of the nasopharynx, spreading from the choanæ into the tubes and far over the vault, the posterior wall of the pharynx, and sometimes the pharyngeal wall of the soft palate.

With anterior rhinoscopy, the growths may be distinguished as they are lifted by the soft palate in phonation, the light reflexes moving with them. They can also be felt with a probe introduced through the nose.

Oil sprayed with an atomizer into one nostril returns with almost undiminished volume through the other if it does not meet with an obstruction. Bosworth advises this method for confirming the diagnosis of adenoid growths. Palpation with the index finger introduced gently behind the soft palate is a quick and valuable way to examine the nasopharynx, even if it is not pleasant nor elegant.

The question when lymphatic glands are pathologically enlarged is usually answered without hesitation. As even the most superficial ones can not be felt through the skin when they are in normal condition, it seems reasonable to pronounce tonsillar tissue pathological when it is plainly visible on macroscopic examination.

The differential diagnosis should be quite clear in the vast majority of cases; as retropharyngeal neoplasms and nasal growths extending into the nasopharynx are exceedingly rare in patients between the ages of 5 and 15, when adenoid growths are most prevalent and troublesome. Palpation would, generally, when the patient is well anesthetized, enable one to decide between uncomplicated adenoid vegetations, on the one hand, and nasopharyngeal sarcoma, or fibrosarcoma, or unusual prominence of one of the retropharyngeal lymphatic glands, on the other. An appeal to the microscope might, however, be necessary if the tonsil should be invaded by malignant or tuberculous disease.

IODIN USED HYPODERMICALLY IN THE TREATMENT OF PULMONARY TUBERCULOSIS.

ALFRED CARENO CROFTAN, A.M., M.D.

Formerly Assistant Professor of General Diagnosis and Clinical Instructor of Diseases of the Chest, College of Physicians and Surgeons, Chicago.

PASADENA, CAL.

The action of iodine on the healthy body has been exhaustively studied; ideal conditions, we may say, were naturally created for such investigation. It seems that the water, soil and air, and consequently the flora of Geneva, Switzerland, are singularly free from iodine; the inhabitants of that region, therefore, constitute a virgin soil for the drug and display a remarkable susceptibility to its action.

Coindet, of Geneva, first described the physiologic effects of the drug; he was led to his studies on iodine while investigating the curative effects of sponge-preparations on goiter—bronchocele—which occurs with such frequency in Geneva; he suspected that iodine was the active remedial principle in the sponge and the other marine products that had for generations been successfully employed for the cure of this affliction. Rilliet, also of Geneva, continued these investigations and in a remarkable "Memoir" presented to the French Academy of Medicine, in 1860, first described iodism and introduced the drug into practice. Trousseau reports on this "Memoir" in the 25th Bulletin of the French Academy. From the reports of these two investigators, that have since been repeatedly corroborated and amplified by reliable authorities—Binz, Niemeyer, Kaemmerer, Sée, Stillé—we glean the following information in regard to the pathogenesis of iodine.

Iodine is peculiarly a drug against which different subjects show marked idiosyncrasies; some people can take large doses with impunity, others are speedily and violently affected by almost incredibly small doses. Thus Trousseau reports that the Geneva goiters are cured with doses of iodide of potash of less than a

milligram, the sponge preparations necessarily contain most minute quantities; in certain sensitive subjects, Rilliet reports "grave accidents" from using quantities of table-salt that contained one part of iodine to 10,000 parts of the salt. He even saw iodism develop in these people from a sojourn at the seashore where, it is estimated, they inhaled from a fiftieth to a tenth of a milligram a day. Cod-liver oil contains about one part of iodine to 40,000 parts of oil, and it is not impossible that the curative effects that this marine product exercises on certain "strumous" diseases and the benefits that are claimed for it in tuberculosis may be attributed to the presence of even so minute a quantity of iodine.

The chief symptoms observed are emaciation, usually accompanied by profuse sweats, some pyrexia and an accelerated pulse; a peculiar psychical depression develops a form of hypochondriasis, "anxietas."

A large field of action is the mucous lining of the respiratory tract; when the system becomes saturated with iodine, almost any of the manifestations of inflammation of these tissues may supervene, coryza, angina, laryngitis "with hoarseness, aphonia and chronic inflammation, even simulating laryngeal phthisis;" the lungs show "congestive oppression, and hemoptysis may occur;" Zeiss¹ reports an "abundant serosity" in the lungs of dogs that had been subjected to subcutaneous injections of iodine and an iodide in solution. The serous membranes, too, become affected and pleuritis has been observed. These effects on the respiratory passages, it must clearly be understood, are remote effects of the drug and follow the administration of iodine wherever in the body it may have been introduced; they must not be confounded with its local irritative action following inhalation or topical application.

The lacteals and the lymphatic system are probably the main sphere of action; iodine acts as an irritant, and possibly this explains the emaciation, inasmuch as the chyle is absorbed more slowly, is insufficiently elaborated by the mesenteric glands, and the absorption of fats is, as a consequence, reduced. Hence, probably, too, the profound remedial action of iodine on bronchocele, lymphatitis—scrofulosis, strumous diathesis—and tabes mesenterica, in all of which afflictions the lymphatic and lacteal systems are inclined to hyperplasia, with resulting sluggish function, etc.

If we study the symptomatology of incipient tuberculosis we find that essentially the same tissues are stimulated to a reactive display of energy by the tubercle virus as by iodine. Again, we must carefully exclude the local, mechanical, and destructive action of the living parasite, and try to draw into the circle of our considerations only the remote effects that are exercised by the products of microbial life near the focus of infection and in distant parts of the body.

Locally, we have that "liquefaction of tissue" that is so commonly attributed to the action of iodine; remotely, the specific selective affinity for the respiratory mucosa, the serous membranes, the lympho-lacteal system; we have the emaciation accompanied by pyrexia, tachycardia and sweats; probably the initial psychosis so often observed is the same.

In the light of our theoretic beliefs the administration of iodine should act curatively in pulmonary tuberculosis; the virus of tuberculosis, entering the blood at first in minute quantities, produces certain slight reactive phenomena from the

1. La Clinique, Feb., 1900.