always likely to cause abortion. No matter how likely the appearances are that a woman is likely to have obstruction during confinement, it rarely occurs; and unless there are positive indications we should not interfere in any way with the pregnancy and not consider the induction of abortion.

DR. A. H. TUTTLE, Cambridge, Mass., referred to two cases of pregnancy complicated with uterine myomata. In the first case the patient was very much upset with symptoms of inflammatory and strangulatory complications, which progressed until, when in about the fourth month of her pregnancy, it was deemed necessary to make an exploratory incision and determine the exact condition of things. A diffuse myomatous condition of the posterior wall of the uterus was found which interfered with the upward extension of the organ. With gentle manipulation the uterus was loosened and raised from the pelvis and the abdomen closed without further treatment. In spite of the fact that the manipulations of the uterus were of the most gentle nature, severe contractions set in and seriously threatened to terminate the pregnancy. Under full doses of opium this complication was controlled, and the case progressed to term without further trouble. A difficult labor was rewarded with a living child, and since the uterus has been removed.

The second case presented four uterine myomata on the anterior surface. The largest, of about hen's egg size, was undergoing gangrenous changes when it first came under his notice, and the indications for their removal were complete. The mother was then between four and five months pregnant. The tumors were completely removed, every care being exerted not to disturb the uterus. The parts healed by primary union without any complication, and a perfectly normal child was delivered at term.

From the total absence of contraction pains and complications in the second case where there was surgical interference without manipulation of the uterus, and the presence of severe contractions in the first case where there was no surgical treatment but considerable manipulation, it would seem that much manipulation of the pregnant uterus will not be tolerated and that in the surgical treatment of the gravid uterus, handling or disturbing the position of the uterus should be avoided as much as possible.

DR. A. E. BENJAMIN, Minneapolis, said that there are three indications for operating on these cases. First, the pain experienced; second, the location of the tumor; and third, degeneration. A great many of these women complain of pain which gradually increases, with symptoms of abortion. Two cases of this sort he operated on without any trouble following. He had one degenerative case. The patient made a good recovery. Dr. Boldt referred to the position of the tumor. In one case with a large tumor filling the pelvis, death of the child resulted, because the attending physician was unable to deliver the child without crushing the head. The woman passed through the ordeal and later came to operation. This tumor was so large and fixed low in the pelvis that it seemed almost impossible that a child's head could pass through the opening.

DR. J. H. CARSTENS, Detroit, said that if the tumor is at the fundus he lets it alone; if it is down low it ought to be enucleated, or if it is in the posterior cul-de-sac and adherent, it must be operated on. If it can be lifted over the brim of the pelvis, operation is unnecessary. Sometimes a tumor is found between the uterus and the bladder. A tumor at this point grows rapidly and interferes with the process of delivery. These tumors may be removed by vaginal section.

DR. S. P. WARREN, Portland, Me., said that the first Cesarean section made in Maine was performed for a fibroid complicating delivery. He has seen but two cases in an obstetric practice of thirty years in which fibroids obstructed labor; one case was associated with eclampsia and acute nephritis at the sixth month, the mother dying of sepsis a week postpartum, and the other case required craniotomy before delivery. At the second labor of the same woman there was no fibroid found in the uterus. She had not been operated on for it, but the fibroid, which was as large as a child's head at the first, had entirely disappeared at the second labor.

DR. C. C. FREDERICK, Buffalo, N. Y., has seen five cases of pregnancy complicated with fibroids in the last ten years. He agreed with Dr. Carstens that fibroids below the ileo-pectineal line, if seen early enough, should be enucleated or Cesarean section must be resorted to. He has had the same experience with fibroids at the junction of the cervix and body. These should also be enucleated. In answer to the question last asked, he has twice seen labors with fibroid tumors in which the tumor was discovered at the time of the labor, and in both these cases the tumors were absorbed or disappeared, leaving only a little nodule. What occurs is a simple process of absorption during involution.

DR. W. B. DORSETT, St. Louis, Mo., referred to a case in which he wanted to do an enucleation, but found the tumor holding the uterus down under the promontory of the sacrum. He was unable and afraid to remove it because it seemed to dip too deeply down. He simply delivered the uterus from the pelvis and the woman went on to labor. He said that a great many ovarian tumors are removed during pregnancy, and they are probably attended by more danger to the life of the child than when a myomectomy is done, because the nerve supply is more apt to be interfered with and as a result miscarriage occurs. In the case of a soft myomatous tumor the question of operation depends on the individual case.

DR. MILES F. PORTER, Ft. Wayne, Ind., differed with Dr. Carstens. He thinks that any tumor, no matter where it is located, that is likely to interfere with pregnancy or labor, should be removed by operation. He has seen a pregnant woman who also carried an eight pound fibroid, but the abdomen would not carry both, hence the tumor was removed, the child was carried to term and born without accident.

DR. E. E. MONTGOMERY, Philadelphia, was in hearty concord with those who advocated non-interference in these cases unless there are absolute indications for interference. He said that it is true that in the subsequent development of pregnancy, the distension of the uterus would lift the tumor out of the pelvis and not interfere with delivery. It is only in the early stages of pregnancy that enucleation should be performed. With regard to the tumor being an interstitial one, in all three of his cases the tumors were interstitial, and at least two-thirds of the uterine wall was entered, and in one the opening was nearly five inches across the denuded surface. It seemed as though the fetus, covered by little more than a membrane, was projecting through. The patient recovered without marked inconvenience and later gave birth to a 13-pound child. These tumors are very vascular and great pressure may lead to rupture of the vessels and hemorrhage and even death of the patient. If the patient is not suffering inconvenience from the too great size of the tumor, he should prefer her to go on to the completion of the pregnancy.

EXCISION OF CANCER OF THE HEAD AND NECK.

WITH SPECIAL REFERENCE TO THE PLAN OF DISSECTION
BASED ON ONE HUNDRED AND THIRTY-TWO
OPERATIONS.*

GEORGE CRILE, M.D.
CLEVELAND, O.

Though signal advances have been made recently in many surgical problems, the treatment of cancer of the head and neck has, it would seem, neither received the attention nor kept the pace of progress in other fields. These unhappy cases are too often regarded as specters at the clinic. The operative treatment is hampered by tradition and conventionality, and the tragic ending of so large a proportion of these cases has held back lay and even professional confidence.

In this paper it is intended to present an outline sketch of the conclusions regarding the surgical treat-

* Read in the Section on Surgery and Anatomy of the American Medical Association, at the Fifty-seventh Annual Session, June, 1906.

ment of cancer of the head and neck in the curable stage. The etiology, the diagnosis and the pathology will not be considered. It is generally admitted that cancer is primarily a local disease. Each case, then, is presumably at some period curable by complete excision.

The immediate extension from the primary focus is principally by lymphatic permeation and metastasis in the regional lymphatic glands. Secondary foci in distant organs and tissues are probably due to cancer emboli. A careful study of 4,500 cases, exclusive of the thyroid gland, traced to their original report in literature, made for me by Dr. Hitchings, showed that in less than 1 per cent. have secondary cancer foci been found in distant organs and tissues. That is to say, in cancer of the head and neck, death almost always occurs by local and regional development of the disease. The collar of lymphatics of the neck forms an extraordinary barrier through which cancer rarely penetrates (Figs. 1 and

Fig. 1.—(From Gray, Poirier, Cuneo and Toldt.) The distribution of the superficial lymphatics. A, Posterior auricular glands. B, occipital gland. C, superficial cervical gland. D, submaxillary gland.

2). Every portion of this barrier is surgically accessible. Paired organs or distinctly one-sided foci usually metastasize regularly, while unpaired organs, as the tongue, or the mesial tissue, such as the nose and the middle of the lip, metastasize irregularly (Fig. 3).

After the lymphatic stream has been blocked, as by carcinomatous invasion, it may flow in any direction and every sort of irregularity in the further metastases may follow, but always somewhere within the accessible lymphatic collar. After all, how much more favorable such distribution is than that of certain other organs, as, for instance, the breast with its thoracic and abdominal metastases, the stomach and intestines with their inaccessible retroperitoneal metastases?

What, then, is the best method of surgical attack? An incomplete operation disseminates and stimulates

the growth, shortens life and diminishes comfort. Local excision of the primary focus only is as unsurgical as excision of a breast, leaving the regional glands. Excision of individual lymphatic glands, as one would excise a tuberculous gland, not only does not afford permanent cure, but is usually followed by greater dissemination and more rapid growth. Judged by analogy and experience, the logical technic is that of a "block" dissection of the regional lymphatic system as well as the primary focus on exactly the same lines as the Halstead operation for cancer of the breast. Such a dissection is indicated whether the glands are or are not palpable. Palpable glands may be inflammatory and impalpable glands may be carcinomatous. A strict rule of excision should, therefore, be followed (Figs. 4 and 5).

In the last 63 cases we have divided them into two classes: (a) those in which the lymphatics were obviously enlarged, (b) those in which they were not en-

Fig. 3.—(Gray.) Emphasizing the venous tree, easy to excise in toto but difficult to dissect individually from lymphatics.

larged. The enlarged lymphatics were presumed to be carcinomatous (this was, of course, not always correct). In this group the lymphatic-bearing tissue was widely excised, because when once the lymph stream is choked by metastasis further dissemination may travel in any direction. When there are no palpable glands the lymphatic drainage area only is excised.

Next in importance to the block regional and local excision is that of a strict avoidance of all handling of carcinomatous tissue so long as the lymphatic channel remains intact, thereby avoiding further dissemination of the growth. Retraction should be entirely on sound tissue and blunt dissection should be the exception.

If the foregoing be mainly true, what, then, are the salient features of the plan and technic, and what are the principal dangers to be avoided? We shall not here con-

sider the preliminary preparation of the patient nor the after-care, but shall address ourselves to the immediate operative problem as to how to secure the highest immediate recovery rate and the most probable permanent cure. What, then, are the immediate dangers attending such operations? Infection, local and broncho-pulmonary, stands first, then hemorrhage, shock and exhaustion.

Infection.—In the present status of surgery, infection may be dismissed without discussion in all clean dissections of the neck in which the wound does not communicate with the mouth, the air passages or the esophagus. The greatest infection risk exists when an extensive dissection wound of the neck communicates with the mouth or air passages, as in cancer of the tongue or of the larynx. In the latter instance experience has emphatically taught us that the primary wound in the neck should not be made at the same time that the larynx is excised. A preliminary tracheotomy is doubly indi-

of passing closely fitting rubber tubes through the nares into the pharynx, opposite the epiglottis, then close gauze packing of the pharynx, after drawing the tongue well forward. With the exception of operations requiring the removal of the entire base of the tongue, complete control of blood inhalation is thereby accomplished. Incidentally, this separates the surgeon and the anesthetizer, permitting each to do his work unhampered by the other. Mucus that otherwise might collect in the throat is absorbed at once by the gauze packing. The prevention of vomiting is almost wholly under the control of an even anesthesia. The many other features of the prevention of infection will not here be considered.

Hemorrhage.—The control of hemorrhage is important on its own account and almost equally on account of the maintenance of a dry field, affording opportunity for a clean and precise dissection. The arterial hemorrhage is best controlled by temporary closure of the common or external carotid artery. Permanent closure of the common carotid, on account of the high percentage of cerebral softening in the cancer period of life, should be avoided. Permanent closure of the external carotid, while it is not attended by this risk, carries with it approximately a 2 per cent. mortality rate from the washing away of the thrombus of the ligatured stump, causing cerebral embolism. At all events, it is unnecessary. In 61 cases I have temporarily closed the common or the external carotid without immediate or remote complications. Proper closure of this vessel should be attended with little more risk of thrombosis or embolism than closure by tourniquet or by pressure applied on the skin. This part of the technic must be done with absolute gentleness, the lumen merely closed, the walls not compressed. The troublesome venous hemorrhage may be minimized by placing the patient in the partial upright posture with a sufficient even pressure on the lower extremities and the trunk up to the costal borders to prevent gravitation of blood. Probably the most convenient means for accomplishing this is by my rubber pneumatic suit. By this device almost any posture may be assumed without serious risk of cerebral anemia by gravity. It is interesting to note the collapse of the veins as the patient is taken from the horizontal to the head-up inclined posture. In atheromatous subjects, presenting a high blood pressure, I have seen marked hemorrhage, even pulsating hemorrhage, from the branches of the external carotid when the common was closed. This is easily demonstrated to be due to the passing of the blood of the opposite artery through the brain, causing a reverse stream through the internal carotid of the clamped side to the bifurcation of the common, then up the external carotid, causing pulsating hemorrhage from the branches of the latter. When this occurs, as soon as the external has been exposed, its

Fig. 4.—(From Poirier and Cuneo.) In cancer of the tongue the metastases extend well down as this cut illustrates. Sometimes the first point of metastasis is in the gland lying between the internal jugular and the omohyoid. A, posterior belly of digastric. B, submaxillary gland. C, thyrohyoid. D, omohyoid. E, sternohyoid.

cated, as, aside from the short circuiting of respiration and fixing the trachea, it produces a wall of protective granulations across the top of the dangerous mediastinal area. After the establishment of this barrier, together with the superficial treatment of the stump of the trachea, but little risk of broncho-pulmonary infection remains.

In the case of the tongue the operation is best made in two stages. It is now well known that the immediate and perhaps the most important factor in the causation of broncho-pulmonary infection in operations within the mouth is the inspiration of blood at the time of the operation. This may be absolutely prevented by the adoption of a simple method which I have heretofore described in the *Annals of Surgery*. The method consists

Fig. 5.—An incision which exposes the regional lymphatics which are excised in cases of lateral cancer of the lip without palpable metastases. After metastases have occurred, excision of the entire lymphatic bearing tissue of the side of the neck should be done.

of passing closely fitting rubber tubes through the nares into the pharynx, opposite the epiglottis, then close gauze packing of the pharynx, after drawing the tongue well forward. With the exception of operations requiring the removal of the entire base of the tongue, complete control of blood inhalation is thereby accomplished. Incidentally, this separates the surgeon and the anesthetizer, permitting each to do his work unhampered by the other. Mucus that otherwise might collect in the throat is absorbed at once by the gauze packing. The prevention of vomiting is almost wholly under the control of an even anesthesia. The many other features of the prevention of infection will not here be considered.

When this occurs, as soon as the external has been exposed, its

lumen may be closed by a clamp, thereby ending the reverse as well as the direct stream of blood.

Shock and Collapse.—In these extensive operations prevention of shock and collapse may be effectively accomplished. The important shock-producing factors are: the number and intensity of surgical contacts, as forcible retraction, vigorous and repeated sponging, blunt dissection, tearing, etc.; the loss of blood; mismanaged anesthesia, and the duration of the operation. The principal factors causing collapse are: interference with the trunk and certain branches and terminals of the vagus, excessive hemorrhage, air emboli and anesthetic accidents.

To minimize the foregoing, the operative field should be subjected to the very minimum trauma by the employment of ample operative space, sharp dissection, minimum retraction, by preserving the field free from blood and minimizing the sponging. The loss of blood

This is analogous to the ether-morphin anesthesia in certain prolonged abdominal technics, in which the less painful part of the operation is done principally under the morphin factor of the anesthetic alone. The factor of hemorrhage may now be practically eliminated, and with it air embolism.

In operations on the tongue it is well to remember that the lingual arteries may be closed by the firm pressure of the gauze packing at the base of the tongue in the technic of tubage. In neck dissections the desired standard of the technic is attained when the field is so clear that the minute tissues, such as individual lymphatic vessels, are clearly and distinctly seen, and when one would not expect to damage even the sympathetic nerve unintentionally.

In estimating the possible shock, collapse or exhaustion, it is well to bear in mind that in the cancer period of life there are apt to be cardio-vascular changes of im-

Fig. 6.—(Continuation of Fig. 5.) The regional lymphatic bearing tissue is removed by a block dissection, leaving the salivary gland in the cases in which the lymphatics are free from metastases, and excising it when any of the regional glands are involved. A, submental. B, facial veins and artery. C, gustatory. D, digastric. E, submaxillary. F, platysma. G, parotid gland. H, lingual. I, digastric. J, sternocleidomastoid.

may be reduced to a minimum by methods already suggested. A hypodermic injection of atropin is given half an hour before beginning the operation for the purpose of paralyzing the nerve endings of the vagus in the heart, thereby wholly preventing an inhibitory collapse from direct or reflex inhibition through the vagus or its branches and controlling bronchial secretion. Morphin favors quiet anesthesia and partially supplements the general anesthetic. By taking advantage of the distribution of the sensory nerves, the supply of which is rich in the skin and superficial fascia while scanty in the deeper planes of the neck, but little ether, in addition to the morphin, is required during the latter dissection.

Fig. 7.—Drawing from a case of epithelioma of the face near the ear. There were no palpable glands. The local block dissection here included a portion of the sternomastoid so as to afford a better exposure of the parotid group of glands, thereby protecting against injury of the facial nerve or the jugular. In this instance the lower part of the parotid gland was excised. The submaxillary was left. A, sternohyoid. B, digastric. C, internal carotid. D, carotid. E, sternocleidomastoid. F, submaxillary gland. G, hypoglossal. H, thyrohyoid. I, external carotid. J, omohyoid. K, platysma. L, internal jugular.

portance. A high degree of atheroma, a senile heart, or a chronic myocarditis are all important factors which should be met before the day of operation by such therapeutic measures as digitalis, nitroglycerin, etc., as may be indicated. Unless these factors are anticipated but little can be done in the midst of an operation toward their mitigation. When the wound is large, one must not forget the importance of moist saline compresses to prevent the effect of exposure to the air.

The practical application of the foregoing data may be illustrated in the technic for the excision of carcinoma

of the floor of the mouth invading the lower jaw with extensive metastases in the submaxillary lymphatic glands. In this instance one would be compelled to take the double risk at one seance. Assuming all preliminary preparations to have been made, the patient is given an injection of $1/4$ morphin and $1/100$ atropin half an hour before beginning of anesthesia. After completion of anesthesia the pharynx is cocaineized to prevent reflex inhibition from manipulation; two rubber tubes closely filling the nares and having perforations at the distal end are pushed down to the level of the epiglottis, the tongue is then drawn well forward, a large piece of gauze packed firmly into the pharynx, completely filling it, the brunt of the packing being made at the sides of the tubes, preventing their compression; the patency of the tubes and easy respiration is readily verified; a T-tube is then connected up with an inhaler and the anesthetizer takes his place a foot or more from the field of operation, giving him the opportunity of continuing an

temporary closure made. The complete skin incisions are then made, the skin reflected back over the entire area of the field. The sternomastoid is divided, the internal and the external jugulars are secured, tied double and divided at the base of the neck (Fig. 7). The dissection is then carried from below upward into the deep plane of the neck behind the lymphatic glands, working first at the sides, then posteriorly, carrying upward all the fascia, muscles, veins, fat and connective tissue until the floor of the mouth is reached (Fig. 8). The lower jaw is then divided at a safe distance on each side of the growth. The floor of the mouth and the border of the tongue are then similarly divided, completing the block.

In very critical cases, an assistant experienced in the use and purpose of the sphygmomanometer is placed in charge of the management of the circulation. If the pressure falls below a certain point the pneumatic pressure is increased, and after the operation the suit is gradually decompressed, but in operations in the mouth in which there may be some oozing, and in which the posture of the patient may measurably prevent inhalation of blood, by continuing the rubber suit the patient may be allowed to come out of the anesthetic in the head-up posture. The pneumatic suit may be gradually decompressed after the circulation has reached a safe physiologic status.

Clinical.—My experience consists of 132 operations for cancer of various parts of the head and neck. A study of the literature convinces me that one could obtain the safest conclusions from his own experience, as it is impossible to get a precise knowledge of the plan, purpose and descriptions of the work of other surgeons. I have, therefore, depended largely on my own tables for my conclusions.

In the first part of this series we now appreciate as we could not then, that incomplete operations do more harm than good. Our principal errors on this point consisted in the removal of individual glands, in allowing insufficient space for the dissections, in regarding hemorrhage too lightly, in handling carcinomatous tissue, in limiting the dissection only to tissues known to be infected; in short, in not doing a complete, comprehensive operation. The early and vigorous return growth clearly proved the fallacy of our technic. Experience, too, more clearly taught us what cases should be considered inoperable. We have never seen benefit from excising a recurrent malignant tumor which had transgressed lymphatic glands and freely invaded the deeper planes of the neck. If, on the other hand, the deep planes were free and the skin involved, there still remained hope of cure.

Superficial epitheliomata in aged subjects, appearing on various parts of the skin of the face, usually about the nose, rarely metastasize. The block dissection when done in cases in which the lymphatic glands of the neck are already involved, is most radical. In all these the key to the situation is the complete ex-

Fig. 8.—(Continuation of Fig. 7.) A method of closing the deeper plane of the neck by suturing the longitudinally split border of the sternomastoid to the platysma and the digastric. A, submaxillary gland. B, posterior belly of digastric. C, platysma. D, sternomastoid.

even, uninterrupted anesthesia, allowing the surgeon full control of the operative field, absolutely preventing the entrance of any blood into the pulmonary tract, and permitting the operator to place the patient in any position he wishes.

The rubber suit has been placed and laced during the administration of the anesthetic, and is now sufficiently inflated to merely support the circulation of the extremities and the abdomen up to the costal arches. The patient is then placed in the inclined posture, head up, and the skin incision over the common carotid artery just above the clavicle is made (Fig. 6). The artery is exposed by an intermuscular separation of the sternomastoid, its or

Fig. 9.—Skin incision for block excision of the gland bearing tissue of the entire side of the neck. A procedure always done when glands are involved.

cision of the internal jugular vein (Fig. 2). In one case both the internal and external jugulars of one side were excised their whole length from the bottom to the top of the neck; when, several months later, metastasis appeared on the opposite side of the neck, a similar excision of both veins was made on this side also, but only after it was discovered what the compensating route for the return circulation was. It was found that some of the superficial branches of the jugular had assumed the size of the internal jugular and were evidently the principal veins of compensation. Following the excision of the internal and external jugulars of the opposite side of the neck there was not the slightest circulatory disturbance, congestion, or any hint of insufficient return circulation (Fig. 9).

Since the adoption of the plan and practice as here described, in 1898-1899, we have seen most encouraging results not only in the immediate recovery rate, but in the permanent cures. We have now apparent cures in patients in whom at the time of operation, various stages of cervical metastases were demonstrated. Among them was a case that was so large and so extensively involved the skin that the patient's physician lanced it, supposing it to be an abscess. This has passed the three-year period. Better selection of cases and improved technic have materially reduced the immediate mortality rate.

It is not intended here to assume that all cancers in the various tissues of the head and neck are alike curable, but when once the lymphatics of the neck are involved the surgical problem and the risk are independent of the location of the primary lesion.

SUMMARY OF AUTHOR'S OPERATIVE CASES.

NUMBER OF CASES, 132.

Age.

Age.	
Mean age 50, youngest 28, oldest 77; 90 observations.	
Cases between ages of 20 and 30 years.....	1
Cases between ages of 30 and 40 years.....	14
Cases between ages of 40 and 50 years.....	21
Cases between ages of 50 and 60 years.....	18
Cases between ages of 60 and 70 years.....	29
Cases between ages of 70 and 80 years.....	6

SEX AND SOCIAL STATE.

Females, 15; males, 96. Married, 40; single, 10.

LOCATION OF PRIMARY FOCUS AND IMMEDIATE RESULTS.

Location.	Operations.	Recoveries.	Deaths.
Epithelioma lateral margin eye.....	6	6	0
Carcinoma of face (including maxillæ).....	15	14	1
Carcinoma of lips.....	31	31	0
Carcinoma of floor of mouth.....	4	1	3
Carcinoma of soft palate.....	1	1	0
Carcinoma of alveolar process.....	2	2	0
Carcinoma of pharynx.....	1	1	0
Carcinoma of tongue.....	12	10	2
Carcinoma of tonsil.....	2	2	0
Carcinoma of nose.....	5	5	0
Carcinoma of scalp.....	2	2	0
Carcinoma of lymphatic glands.....	23	20	3
Carcinoma of parotid.....	5	5	0
Carcinoma-branchiogenic.....	4	4	0
Carcinoma of larynx.....	15	13	2
Carcinoma of thyroid.....	4	4	0
	132	121	11

8 per cent.

OTHER DATA.

Mean duration of disease prior to operation, 1 year. In the last 21 operations there has been no fatality.	
Total number of cases operated.....	132
Total number of cases traced.....	106
Number of these known to be living.....	47
Number of these that did not have radical block dissection.....	96
Number of these operated over 3 years ago.....	67
Number of these traced.....	48
Number who have passed 3 years.....	9
Number of radical block dissections.....	36
Number of block dissections over 3 years.....	19
Number of these traced.....	12
Number of block dissections of patients living who have passed the 3 year period.....	19
0 to 1 year.....	5
1 to 2 years.....	5
2 to 3 years.....	0
3 to 4 years.....	2
4 to 5 years.....	2
5 to 6 years.....	3
6 to 7 years.....	1
7 to 8 years.....	0
8 to 9 years.....	0
9 to 10 years.....	0
10 to 11 years.....	1
Number that did not have block dissections now living.....	28
0 to 1 year.....	6
1 to 2 years.....	10
2 to 3 years.....	3
3 to 4 years.....	0
4 to 5 years.....	3
5 to 6 years.....	2
6 to 7 years.....	3
7 to 8 years.....	1
8 to 9 years.....	1

There were twenty-three operations for cancer metastases in the cervical lymphatics following operations for the excision of the primary focus alone. Some of these were in my own early cases in which I was led to excise only the primary focus, either by the pleading of the patient or because the trifling primary lesion scarcely seemed to warrant excision of the regionary lymphatics (Figs. 10 and 11). The majority have been drawn from other sources. It would seem that the end results in this class of cases are the real test of the effectiveness of any method. The end results here are certainly equal to that of cases of axillary involvement in breast cancer. Increasing experience confirms us more and more in the belief that there has been too much conventionality in the technic. The evolution of the technic has so much minimized the operative risks that the chief consideration at present may be given to the complete excision of the last vestige of the disease along the lines of the principles underlying operations for cancer in general.

I would direct especial attention to the fact that among the 48 traced cases operated on more than three years ago, in which the radical block dissection was not made, nine are living; while among 12 traced cases among the block dissection cases operated on more than 3 years ago nine are living. The radical block dissection has, therefore, shown itself to be four times more effective than the less radical (Fig. 12).

Our general conclusions are that, since the head and neck present an exposed field, cancer here, unlike that of the stomach, the intestines, or even the breast, may be recognized at its very beginning; that every case is at some time curable by complete excision; that the field of regional metastases is exceptionally accessible; that cancer rarely penetrates the extraordinary lymphatic collar of the neck; that the growth tends to remain here localized; that by applying the same comprehensive block dissection as in the radical cure of breast cancer and by freely utilizing the modern researches of surgery the final outcome in cases of cancer of the neck and head should yield better results than that of almost any other portion of the body.

DISCUSSION.

DR. H. M. S. DAWBARN, New York City, took a somewhat different ground, though with the same object in view, the prevention of hemorrhage. Although he has tied the external carotid artery over 100 times, he does not expect to tie this artery hereafter with any such frequency. In the June number of the *International Journal of Surgery*, he said, he gave his technic and reported four cases of brain operation with so little hemorrhage that the artery forceps were hardly needed at all, and in which for half an hour at a time the anesthesia could be laid aside because the patient was lying there with just such an anemia of the brain as is present in normal sleep. By constricting the limbs at the junction of the body tightly enough to accumulate blood in them, there results partial anemia of the head and trunk, and then there is no need of the rubber suit or the posture described by Dr. Crile. There is also very much less pressure at the mouth of the cut vessels. By this simple procedure one can operate in the most vascular region almost without any severe bleeding at all. Such usually bloody operations as excision of the upper jaw and removal of adenoids can be performed comparatively bloodlessly. It is carrying out the same principles in head, neck and trunk, as is done in operations on the extremities by bleeding into the trunk. When operating on the trunk he advised bleeding into the extremities. As soon as the tourniquet is released, as an essential corollary, Dr. Dawbarn applies boiling water by means of gauze wrung out of it with thick rubber gloves. The surface is blanched at once and then there is not that bleeding that ordinarily follows the release of the tourniquet.

If skin and mucous membranes are avoided, the heat will never be found injurious even to delicate tissues.

DR. D. N. EISENDRATH, Chicago, said that in every operation it is essential to remove the lymphatics which drain the field of operation. There are, however, early cases, he said, where it is hardly justifiable to adopt the thorough blocking method Dr. Crile suggested, where a less radical procedure is preferred. Three years ago he began in every case of carcinoma of the lips and tongue to remove the lymphatics in a thorough manner. In these cases nine-tenths practically of all the lymphatic drainage is toward the glands which lie in the submental region between the two anterior pillars of the digastric muscles and in the submaxillary lymph glands lying on the capsule of the salivary gland and within the gland itself. In every case of that kind it is necessary to remove all of the contents of the submaxillary triangles on both sides of the neck at the primary operation, and this should be done as thoroughly as possible, removing not only the fat and fascia, but also the submaxillary and salivary glands. In carcinoma of the tongue surgeons ought to adopt a procedure almost as radical as that mentioned by Dr. Crile. The lymph drainage from the tongue not only goes into the glands into which the lip drains, but also into the deep glands under the sternocleidomastoid muscle, and in some instances even into the glands of the parotid region itself. In these cases one ought to make an incision something like Dr. Crile described. The incision Dr. Eisendrath makes in these cases is slightly different from Dr. Crile's incision. He begins midway between the lower border of the lower jaw and the upper border of the thyroid, from one sternocleidomastoid muscle to the other. By retracting the flap he completely exposes the submaxillary triangles. When the incision is extended this is done from the upper angle of the transverse incision, i. e., from the middle of the sternocleidomastoid muscle down to the clavicle. This enables him to remove every gland from around the muscle without resecting it. In case the internal jugular is involved, it is well to respect that also. Dr. Eisendrath said that no surgeon should content himself simply by removing the primary growth and then dismissing the case. He should remember that in 99 per cent. of the cases metastases occur not in the body, but locally in the regional glands. By removing these and their surrounding fat and fascia, as well as the veins and muscles, if they are involved, the permanent results will be better.

DR. GEORGE CRILE, Cleveland, excises the vein and muscle only in cases in which metastases have already occurred in the lymphatic glands. When metastases can not be demonstrated, only the regional glands that are known to drain the field of the original focus are excised. He said that the functional value of the internal jugular vein and the sternomastoid muscle is not such that one need hesitate to remove them, provided it will add, even to a slight degree, to the certainty of complete excision of the disease. The provision for return circulation of the veins is many times greater than is needed. Dr. Crile in one case removed both the internal and the external jugular veins. Later metastasis occurred in the opposite side, and after the neck was opened he found that compensation had taken place in some subcutaneous veins near the median line, these having enlarged to the size of the internal jugular. He then removed both the internal and the external jugulars of the remaining side, and there was not the slightest circulatory disturbance at any time.

Give Drugs for Exact Purposes.—Shattuck, in the *Boston Med. and Surg. Jour.*, lays down general rules for the giving of drugs: to do no harm; to have a clear idea whether we give the drug—as a specific, a curative, a palliative or a placebo; to give the drug uncombined, if possible, though to this rule there are many exceptions. When using an efficient drug one should be sure that the preparation is efficient; and should continue its use until either the desired effect or toxic manifestations are produced. The *Medical Times* adds the comment that one should be sure the druggist is an accurate compounder and uses fresh material.

DISPENSING VERSUS PRESCRIBING.*

M. H. FUSSELL, M.D.
PHILADELPHIA.

Much is written in various medical journals, much is spoken by men, leaders in the medical profession, of the nostrum evil, of the impropriety of physicians using any preparation of drugs the full formula of which is not known.

Many writers, with great justice it seems to me, insist, as Billings has done in a recent article, that mixtures put out by manufacturing firms, especially when such compounds contain on their labels the various diseases for which it is a specific, are just as objectionable as various secret remedies in which we are not certain of the exact proportions of the various ingredients.

Sal vitæ has on its label a list of fourteen or fifteen ingredients said to be in the compound. It also states just exactly the therapeutic use of each supposed ingredient. It is a blotch on the escutcheon of a manufacturing pharmacist.

A certain preparation containing eight ingredients, the exact proportions of each being given, is said to be a formula used by a famous French specialist. It is prepared by a reputable firm; it is handsome in appearance and pleasant to the taste. It is an excellent example of the kind of mixture against which Billings speaks. The fact that it is said to have originated with an eminent physician is used by the smooth-tongued agent as a specious argument for its use. Doubtless if it does not cost too much it graces the shelves of hundreds of dispensing physicians.

The great advance made in the practice of medicine and surgery in the past century has come because of the heroic labors of men working in pathologic and bacteriologic laboratories, delving for the cause of disease and its remedies; because of men working in hospitals and at the bedsides of private patients studying the clinical aspect of diseases and how best they are prevented and controlled; because of men working in pharmaceutical laboratories searching for the best forms of drugs and their most useful and active ingredients. Of these three classes of men, the second division claims or should claim the great majority of practicing physicians.

The opportunity is given to each individual practicing medicine to add something, be it ever so little, to the sum total of medical knowledge, and thus to the good of mankind. It can be done, is daily done, by thousands of men and women by a careful physical examination of each patient, by recording of observations, by a careful selection of the proper treatment and the means of applying that treatment, and is made efficient by submitting these observations to the medical public.

This paper will have to do with one of the last statements: The means of supplying to patients the treatment necessary by drugs. What shall a young physician do? Shall he dispense from his own office or from his medicine case? Or shall he write a prescription to be filled by a pharmacist? What I shall have to say must relate only to a physician practicing in proximity to a reputable pharmacist. I can see no way at present for the country doctor to work except by dispensing. It appears to me that the choice of these two methods will depend entirely on the education and the character of the young physician.

If he has been instructed that the giving of drugs is a small, but important, item in the duties of an honest

* Read in the Section on Pharmacology of the American Medical Association, at the Fifty-seventh Annual Session, June, 1906.