had two months of tropical service. Although the skin had been noticeably bronzed since childhood nothing indicating gall-bladder trouble could be elicited. A careful inquiry as to gastro-intestinal disturbances and family history was also negative. Heart, lungs and kidneys were apparently normal.

Examination.—On admission to this hospital there was marked rigidity of right abdomen and extreme tenderness over the appendix; temperature 100.4 F.; pulse 78, regular and full; respiration normal; tongue coated and bowels costive. The attack began with epigastric pain and nausea on the day before admission, the pain in a few hours being centered over the appendix; patient vomited before reaching the hospital; white count 13,000. An immediate operation was decided on.

First Operation.—The abdomen was opened by the Battle incision and the appendix found greatly distended, with a small gangrenous perforation at its base, through which fecal matter had escaped. There was also a gangrenous spot on the cecum surrounding the base of the appendix. The appendix was removed, and in addition the gangrenous spot on the cecum was surrounded and buried by a purse-string suture, over which a continuous Lembert suture was introduced. The parts were dried and cleansed by mopping with gauze and a cigarette drain of iodoform gauze brought out through the wound. A rubber drainage tube was passed down into the pelvis and the wound closed with the exception of a small opening for the drains. The patient was placed in Fowler's position and Murphy's proctoclysis started.

Subsequent Course.—On March 10 there was a free discharge of pus from the cecal drain which had a fecal odor. This gradually lessened and the patient passed through an uneventful convalescence with the exception of the development of a small abscess in the rectus muscle, the result of a stitch becoming infected. On March 24 the sinus had completely closed, one of Bier's hyperemia cups being used for five days.

April 1, 1910, the patient had a slight chill in the afternoon followed by profuse sweating. The temperature ranged from 99 F. to 103 F., and a slight pain developed over the gall-bladder region, but the liver dulness was not increased. For seven days after this there was an afternoon chill, fever and sweat, the temperature reaching usually 104 F. Daily white counts never ran above 8,000 and numerous differential counts were normal. The test for iodophilia was negative at this time as were also the urine, sputum and feces. The tongue was thickly coated and anorexia was marked. Repeated blood examinations were negative for the malarial parasite and several Widal reactions were also negative. The Cambridge reaction was negative, as was also a culture of the blood. Iodophilia now began to be positive and patient developed a slight cough, though no lung involvement could be made out. The pulse at times reached 100 but was usually between 70 and 80. Although there was no pain or tenderness in the loin or operation scar the latter was incised and probed but nothing unusual found. The pain over the gall-bladder region increased somewhat and at this time there was quite a perceptible deepening of the bronzed condition of the skin. The lungs and entire abdomen were repeatedly examined but, with the exception of the tenderness over the gall-bladder region, nothing abnormal could be found. At no time could the spleen be palpated nor was there ever the slightest pain elicited by the deepest pressure in this region.

Second Operation.—April 8, a Bevan incision was made over the gall-bladder and the latter found to be imbedded in a rather dense mass of adhesions and these when liberated disclosed a very much thickened and dwarfed gall-bladder. This was incised and explored, as were also the ducts, but no stones were found. The head of the pancreas seemed to be enlarged and harder than normal. The gall-bladder was drained with a Mixter tube and for two days there were no chills or fever, but at the end of that time the pyemic condition began again. The gall-bladder drained well but the symptoms were practically the same as before operation with the exception of the development of a systolic murmur heard loudest at the base. April 31 an aspirating needle was introduced in the eighth intercostal space in the axillary line and the liver was needled in five different directions without locat-

ing pus. The symptoms of marked sepsis continued and the patient finally died of exhaustion on June 9.

Necropsy.—Cutaneous surface was deeply bronzed; emaciation marked; adhesions in right iliac fossa broken up and revealed small, firm, smooth scar in cecum at the site of the excised appendix; ascending colon dissected up to right kidney pouch with no indication of any previous inflammatory condition; gall-bladder firmly adherent to abdominal wall at site of operation wound. On palpating in left abdomen it was found that a mass of small intestine and omentum was closely adherent over the spleen, which could hardly be felt through the distended intestine. The intestinal adhesions were carefully broken up, showing the spleen to be about two and a half times as large as normal and closely adherent to the stomach and diaphragm. In breaking up the diaphragmatic adhesions a large pus sac was opened in the discolored, friable spleen. This sac was located in the center of the spleen and was about the size of a goose-egg. Two smaller abscesses about the size of a pigeon's egg, were located at the periphery. The splenic vein was thrombosed up to the vena cava. On sectioning the liver, which was slightly enlarged, there was to be seen widespread fatty degeneration, with here and there small necrotic spots about the size of a pea. Both ventricles of the heart were dilated and the left ventricle very much thinned. There were to be seen greenish-yellow nodules and patches on the flaps of the mitral valve and left ventricular endothelium, showing the presence of endocarditis. Lungs and pleurae were apparently normal. The head of the pancreas was enlarged and nodular. Both kidneys were slightly enlarged and showed minute areas of fatty necrosis. The small intestine was blanched and contained a small quantity of tenacious greenish yellow material.

The post-mortem findings clearly indicated a widespread pyemic condition with a primary focus in the spleen, and it would seem that the lesson to be learned from the history and findings in this case is that the splenic condition could only be diagnosed by exclusion. Systematic writers on this subject invariably refer to pain in the left side high up. At no time was there the slightest pain in this region, nor could the spleen be palpated. It is believed that the splenic abscess resulted from the gangrenous pyogenic condition of the appendix and cecum. Had there been the slightest suspicion that the spleen was diseased a splenic puncture would have been done. It is further believed that the splenic puncture should have been done shortly after the gall-bladder operation, in view of the fact that everything but the spleen had been excluded. Had this been done the abscess in the spleen could have been drained posteriorly with a possible chance of saving the patient. Of course this latter conjecture is not free from the criticism that the patient could not have lived any way on account of the pyemic condition which was in all probability present at that time, but there is no question as to the fact that at least the patient's chances of recovery would have been greatly improved had the abscess been located and drained, and of course the sooner this was done the more favorable the prognosis.

IDIOSYNCRASY TO ASPIRIN

CYRUS GRAHAM, M.D.

HENDERSON, KY.

In THE JOURNAL I recently read a communication¹ in regard to idiosyncrasy to aspirin, of which I had an almost parallel case on November 19.

On the advice of a friend a young man took a five-grain tablet of aspirin for headache. In about one hour, still suffering, he took another, and then lay down on a cot in his office. In a few moments he arose and remarked to some one present,

1. Berl. klin. Wchnschr., 1893, No. 19.

"I feel mighty queer, and can hardly see." He then looked in the mirror and exclaimed, "What is the matter with my face? I am going to a doctor!" He walked nearly two blocks to my office, and in attempting to walk up to the door (just three steps up from the pavement) fell, and then had to be assisted into my office. He hurried in without waiting to knock, and in a voice which indicated great distress, exclaimed, "Doctor, some one has poisoned me." This was probably about half-past one, and about an hour and a half after taking the first tablet. The mucous membranes of his eyes, nose and mouth were very edematous. The eyelids and lips were everted and swollen to twice the normal size, the conjunctivæ very much congested; the patient could hardly breathe through his nose at all. Tonsils and uvula were in a state of congestion, the articulation was very indistinct, and when the patient attempted to raise his voice to a higher pitch, it had a peculiar metallic sound, which would end with a muffled tone. The veins in his temples and throat were knotted, and in places were standing out like hard cords. I found his circulation 136, respiration 24, afterward going down to 12, temperature normal. His face was greatly flushed, and this, with the eversion and swelling of eyelids and lips, made him a ghastly-looking object.

I made him lie down on a cot, and as he seemed to have considerable trouble in breathing, removed his collar and opened his shirt. I had the nurse give him a hypodermic injection of digitalin, bathe his face, neck and head with hot water, and apply continuous hot towels, and keep him quiet. Later, as he complained of great pain in the shoulders and neck, I also applied continuous hot pack to those parts, and, as the circulation remained weak, ordered hypodermic of strychnin. At half-past two he suddenly fell into a profound sleep; his circulation slowed down and became stronger, but respiration continued very slow, and seemed almost to stop at times. His slumber was so profound that the removal and the replacing of the hot towels did not seem to disturb him at all. I allowed him to sleep until nearly four o'clock, when I awakened him and had him conveyed to his home, which was about three miles in the country. He stood the trip all right, and after getting into bed he almost immediately fell asleep again. In about three days the edema had disappeared from face, nose, eyelids and mouth, but the throat remained irritated for at least ten days longer, and the patient was slow in regaining his strength.

The report of H. E. W. and this case show that some have a decided idiosyncrasy to this drug, and that it should be listed as one of the dangerous drugs, which should not be retailed indiscriminately to the laity.

THE USE OF CYLINDRICAL GAUZE AND COTTON DRAINS IN DISCHARGING EARS

MARK D. STEVENSON, M.D.
AKRON, O.

Aural surgeons must have witnessed the efficiency of gauze drainage in the ear following mastoid operations. The pus is constantly carried by the moist gauze packing in the external auditory meatus to the preferably wet gauze protective dressing. The solution with which this dressing is moistened should not be strongly antiseptic or irritating. Normal saline is excellent, although to encourage osmosis I sometimes employ a heavy sterile solution of magnesium sulphate. The packing in the auditory canal should be carried down to the tympanic membrane. There should not be any space left to serve as a reservoir for pus. This obliteration of space may be readily accomplished by placing over the outer end of the auditory canal a few folds of the end of a narrow 1 inch or 2 inch strip of gauze packing which has no

loose threads on its margin, and then with a rough or screw-ended probe pressing the gauze directly in, until it touches the drum head. If a smooth probe is used or no fold made in the gauze the probe is likely to slip through the single layer of gauze, irritating the canal or injuring the ear-drum instead of carrying the gauze before it.

In acute or chronic otorrhea, a gauze drain is also very useful, but requires changing once or twice a day, which must be done by the surgeon, since such a drain cannot be inserted safely by untrained attendants. A few aurists, in certain cases, advocate gauze drainage but the patient must be in a convenient hospital or make frequent visits to the aurist's office which in practice is likely to be either too inconvenient or too expensive, or both. To overcome these objections I employed for some time a drain which my office assistants made by rolling a narrow strip of gauze doubled on itself on the end of a sterile toothpick, giving it stability by wrapping its central part with thin gutta-percha gummed together by any unirritating sterile ointment. It presented the appearance of a diminutive cigarette, $\frac{3}{4}$ to $1\frac{1}{4}$ inches long, about $\frac{1}{8}$ inch in diameter, with the gauze projecting about $\frac{1}{8}$ inch beyond the surrounding gutta-percha and with a toothpick sticking in the unravelled end of the gauze. The drain was made loose on the toothpick before being introduced. After its partial introduction the toothpick was withdrawn and its opposite dull end was used to press the gauze drain well into the ear. The outer end of the canal was then packed with gauze to help retain the drain in place and to absorb any excess of discharge when there was more than the drain could contain. Gauze is much more serviceable than cotton for this purpose.

Similar drains are now being manufactured. At first drains made altogether of gauze tightly wrapped by machinery were tried, but they were too stiff, hurting the patient on introduction. Drains made wholly of cotton were not stiff enough to be introduced. The cylindrical gauze and cotton drains which I recommend consist of cylinders of cotton wrapped in gauze, which is both soft and pliable, so as not to injure the delicate tissues of the ear, although sufficiently firm so that they can be readily introduced. They are supplied sterile, ten in an envelope, each six inches long. They may be cut by sterile scissors to any desired length, depending on the depth of the external auditory canal, and are supplied, of proper length, to the patient, in dry sterile two-dram vials or wrapped in sterile gauze. These drains are also very useful in draining small abscesses or cavities anywhere, and are easily introduced. They may be used any desired length. Several, side by side, may also be employed for drainage, as in some mastoid wounds.

They are easily introduced into and removed from the external auditory meatus by any simple forceps. They may also be introduced by first impaling them on the end of a sterile toothpick, using the latter as a handle. When there is much discharge I have the patient keep one in the ear day and night, putting a fresh one in every morning and evening; but after the discharge is much lessened it is employed only at night. After its removal, if there is any excess of discharge, it should be wiped away with cotton on the end of a sterile toothpick. Any drops which the aurist may wish to employ, *e.g.* 5 or 10 per cent. phenol in glycerin, compound tincture of benzoin, boroglycerid solution, or boric acid and alcohol solutions in different strengths, may be warmed and poured into the upturned