which means, of course, to operate promptly. Do not wait for all the positive and confirmatory evidences of the trouble or the golden opportunity will have been lost. Nor should we be too greatly perturbed about shock when present, as it is in these cases no indication for delay. Some of the best results have been in patients operated on in shock.

As to the technic of the operation, I would only say, do not try to do too much. Do not lose time in useless tinkering and fussing. Do the work quickly—best through an incision in the right linea semilunaris—close the opening in the bowel, and be sure to amply drain the cavity. If the opening is not readily found, let it be and drain extensively. To prolong the operation needlessly is bad surgery. If the patient is a bad subject for a general anesthesia, use a local anesthetic.

Escher has recommended stitching the edges of the intestinal opening to the parietal incision, and Le Compte advises damming off the infected area, particularly when there are multiple openings near each other, with a gauze wall. An artificial anus presupposes only one perforation. The mattress Lembert suture is looked on as the best. Extensive flushing and leaving some of the normal saline solution in the belly, with of course drainage, is, I believe, the preferable way to overcome the peritonitis, although Finney and others continue to advise dry mopping of all infected areas.

In conclusion, I wish to read the following table of Elsberg and some additional cases I have found in the literature bringing the reported cases to date:

REPORTED CASES OF OPERATION TO APRIL, 1903.

	Adults.	Children.	Total.
Cases operated on.....	264	25	289
Recoveries	59 (22.4%)	16 (64%)	75 (25.9%)
Deaths	205	9	214 (74.1%)

ADDITIONAL CASES.

	Cases.	Recoveries.
Armstrong	8	5
Crile	4*	2
Chevallier and other Parisian surgeons.....	13	5
Frank	1	0
Harte	13	3
Méry	1	0
Total	40	15 37.5%

*One unoperated.

Of these last, 14 are reported as children with 5 recoveries, or 35.57 per cent.; 26 adults, 10 recoveries, or 38.46 per cent.

My own case was as follows:

A man, 32 years of age, in the third week of typhoid fever, living in the country, was seized with a sharp pain in the right iliac fossa, which disappeared after copious movement of the bowels. Two hours later his temperature became very slightly subnormal, and continued so or normal for twenty-four hours. When seen by me 60 hours later his temperature was slightly above 99 F., pulse between 86 and 90, though it rapidly rose to 112. There had been no vomiting, and distension of the abdomen had been present for only a short time after the sudden pain. The patient presented a good appearance, without pain or tenderness, and was not depressed. Tympany was present over the entire abdomen as far as the fifth intercostal space, with marked rigidity and absence of liver dullness. The pulse was hard and characteristic of peritonitis. Diagnosis of perforation was made, with prognosis of death, with or without operation. The patient asked that he be given the slight chance by operation. On opening the abdomen it was found filled with pus and fecal matter and the intestines distended above a point about eighteen inches above the perforation, below this point the intestine was flat. The perforation was closed with difficulty, the cavity washed out and five drains inserted. The patient was put to bed in fairly good condition, and next morning had a pulse of 130, no vomiting or pain. He died on the second day.

CONCLUSIONS.

From the foregoing we can make the following summary:

1. Perforations are to be expected in about 2.5 per cent. of all cases of typhoid fever.
2. Prompt surgical intervention is the best and only logical treatment.
3. Early diagnosis is most desirable, and will be the means of greatly reducing the mortality, as 55 to 60 per cent. would recover.
4. Diagnosis, sufficiently early to achieve these results, can only be made by careful watching, treating all cases as serious ones and a proper interpretation of the early, even the preperforative, symptoms, the suggestions of Cushing as to this stage being of value. At the first indication of this stage have the surgeon in consultation and be prepared to operate.
5. The sphygmomanometer should come into general use as an important aid to diagnosis.
6. More cases die from delay than from errors in surgical technic. Therefore, in doubtful cases, though a mistake may be made and no perforation found, operate.
7. No case, unless dying, is so desperate as to be beyond some hope of saving. So in operating, lose no time and be sure to drain.
8. Get into the abdomen quickly and get out more quickly.

(NOTE.)—Since this paper was written, Dr. W. D. Haggard of Nashville has reported, in a paper read before the American Association of Obstetricians and Gynecologists, three cases, with three recoveries.—L. F. (See THE JOURNAL A. M. A., Oct. 17, 1903, p. 979.)

A SIMPLE, ACCURATE AND RAPID METHOD OF LOCALIZING FOREIGN BODIES IN THE ORBIT.

VARD H. HULEN, A.M., M.D.
SAN FRANCISCO.

When the eye has been injured and there is a possibility that a foreign body remains, the very first procedure is to determine, if possible, the exact location of the object in reference to the eye. If the foreign body can not be seen with the ophthalmoscope we must resort to the x-ray, and should the result of such examination be positive, we then proceed to localization before any means whatsoever are tried for its removal.

As the vast majority of the foreign bodies in eye cases (iron, steel, copper, lead, glass, etc.) are revealed by the Roentgen ray, we can usually be certain of a positive result if the foreign body be present. The circumstances attending the injury often determine the probable substance of the foreign matter present. Fortunately most of such injuries to the eye are produced by pieces of iron or steel, and as these objects are attracted by a magnet we have thus an invaluable method for their removal.

Since the advent of the incomparable giant magnet of Haab, great advancement has been made in the successful treatment of these cases, but I do not at all agree with those who advocate using the Haab magnet in an attempt to determine whether or not an eye contains such a foreign body. I do not consider it good surgery or good sense to attempt to remove a piece of steel from the eye before its exact location is known. This being so, it is most desirable to have a method of localization

that is simple, that is quickly used and that is absolutely accurate.

A number of methods of localization have been advocated, but the only one to be recommended, with which I am familiar, is the method of Makenzie Davidson. In London over two years ago I had the pleasure of seeing Mr. Davidson repeatedly apply his method

Figure 1.

with excellent results. But his method requires considerable special apparatus, a good deal of time, and I find the exact details are not easy to carry in the mind. For these reasons I sought a simpler and more rapid method, and yet one equally as accurate. Since August, 1902, I have employed a method of localization which has fulfilled all the requirements. With me it is original,

Figure 2.

but I do not suggest it as a new method, for not a few things in medicine presented as new later are proven otherwise.

My "localizer" (Fig. 1) can be made by any carpenter, and the photograph presented is explanatory of its mechanism. As the head must be absolutely immovable, I have the patient hold her position firmly with her

teeth, in preference to Makenzie Davidson's head fixation (Fig. 2).

The method of procedure is as follows: A piece of lead of suitable size and shape is securely fastened with adhesive plaster to the margin of the lower lid of the injured eye directly under the center of the pupil. (This marker might be fastened in the conjunctiva.) Then the distance from the upper extremity of this "marker" to the center of the cornea and also the distance the marker is anteriorly or posteriorly to the apex of the cornea is taken and recorded. A Crookes tube is next adjusted vertically in the holder of the localizer so that the luminous point of the anode and the point where the wires cross and the V-shaped sight are in a horizontal line. Then the distance from the anode to the cross wires is measured and recorded. The patient is placed in position, a photographic plate is put in the frame and Photograph 1 is taken. The patient remains in the same position, but the tube is raised a definite distance,

Fig. 3.—Case 1. Photograph 1. X. Foreign body. O. Point of marker. In this photograph, as also in Figures 4, 9 and 10, the foreign body shadow has been marked over to make it more clear.

usually three inches, and Photograph 2 is taken, and both pictures are sent to the dark room to be developed. The marker on the patient's lid is again inspected to be sure it has not shifted its position since first measured.

With these two plates and the accurate measurements one can quickly determine the exact position the foreign body occupies in the orbit.

On a sheet of drawing paper make two vertical lines, the distance between them being that recorded for the distance between the anode and cross wires. Cross each of these lines near the bottom by a horizontal line, and mark the point of crossing of left line "1." Then mark a point "2" on this left vertical line, the same distance above as the tube was displaced vertically when taking the second picture. Then in Photograph 1 measure the vertical distance from the shadow of the horizontal wire to the upper extremity of the shadow of the marker, and

on the right vertical line in the diagram beginning at the point crossed by horizontal line measure vertically this distance, designating that point "0, 1." In the same way measure and mark the vertical distance from shadow of horizontal wire to shadow of foreign body "X, 1." Now take Photograph 2, measure and mark shadows of the foreign body and marker on same vertical line as with Photograph 1, designating the point of the foreign body "X, 2," and that of the marker "0, 2." Then connect by a continuous straight line point "1" on left vertical line with point "X, 1," on right vertical line, also by a broken line with point "0, 1"; connect point "2" in a similar way to "X, 2" and "0, 2."

It is evident that the point where the *continuous* lines cross represents the location of the "marker," and that the point where the *broken* lines cross is the exact location of the foreign body in the vertical plane. As the relation between the eye and the marker is known, draw schematically an eye according to its anatomic

Fig. 4.—Case 1. Position 2.

measurements in its correct position, using the vertical measurement previously made from the center of the cornea to the upper extremity of the marker. The location of the foreign body to the eye in the vertical and lateral meridians is thus made plain by our diagram.

Now for the upper part of the figure. Mark a point on the left vertical line "1" and also a short line at the corresponding horizontal point on the right vertical line. Then in Photograph 1 measure the distance from the shadow of the vertical wire to the shadow of the "marker" and mark the point of that measurement above the crossing of the short line "0', 1'"; also measure the distance from the shadow of the vertical wire to the shadow of the foreign body, and as above described, mark it "X' 1'." Connect point "1" by a continuous line to "0', 1'" and by a broken line "X', 1'." Then erect across the paper a perpendicular continuous line through the point corresponding to the marker and a

broken perpendicular line through the point corresponding to the foreign body in the lower part of the diagram. Again it is evident that the points where these lines cross the above lines will mark the exact location of the marker and the foreign body in the horizontal plane. Again draw the eye in its correct relationship to the "marker," using the other measurement from the marker to the apex of the cornea. Thus the location of the for-

Fig. 5.—Case 1.

foreign body in the relation to the eye in its horizontal meridian is revealed and the localization is complete and perfect.

I need hardly say that after the idea has been grasped, one can draw this complete diagram unhesitatingly in far less time than it takes to read the description of how it is done.

Three foreign body cases are chosen from my records

Fig. 6.—Case 2. Photograph 1.

and reported for their interesting points, as well as to illustrate this method of localization.

CASE 1.—Male, aged 43 years, expert machinist. Fourteen years ago he was struck over the right eye causing loss of vision of this eye from traumatic cataract. The sight of the left eye was perfect until Aug. 14, 1903, when a flying chip of cast iron from a neighboring workman's chisel injured it. He

placed himself in the care of an oculist within an hour after receiving the injury. Not much pain followed, and he was able to perform his toilet, but vision was rapidly lost. His oculist reported that at the first examination there was a small wound at the sclerocorneal margin below and to the temporal side, and a clot of blood or something in the vitreous to the nasal side. The patient did not believe the iron had entered the eye

Fig. 7.—Case 2. Photograph 2.

at all. A small magnet was applied and later the Haab magnet was tried repeatedly without any result or any sensation of pain or drawing feeling about the eye. Very slight reaction followed the injury, but he was kept under daily observation.

Four months later the patient consulted me. The left eye was free from pain and inflammation, a traumatic cataract obscured the pupil, a very insignificant scar marked the site of

Fig. 8.—Case 2.

the injury. Vision $= 1/\infty$. Projection faulty. $T = -\frac{1}{2}$. An x-ray examination showed the presence of a foreign body in the orbit.

Localization by the method described above revealed the fact that the piece of iron was in the inner and lower part of the wall of the eye. As this condition seemed improbable I felt that some slight error had been made in my measurements, and that probably the iron was inside the eye, resting on the

wall at the site shown. Then I tried my giant magnet again and again with a negative result. The patient was sent to the hospital, and next day, under chloroform, an incision, guided by the localization, was made with a cataract knife, cutting into the iron, but it was even then found impracticable to remove it with the magnet or forceps, and as the eye was hopelessly blind in any case, enucleation was done. Thus I became possessed of a most interesting and beautiful specimen, for the small elongated piece of iron was found to be in the wall of the globe, one point without and the other point within the eye. My localization had been absolutely accurate.

Subsequently the cataract was successfully removed from the right eye.

This case proves that the giant magnet is not at all a reliable test in the presence of an injury to the eye from iron or steel, for its negative result misled the patient's former surgical adviser, who counseled a let-alone plan of treatment.

CASE 2.—Male, aged 22 years, logger in lumber camp; referred to me by an oculist confrère with the history that on

Fig. 9.—Case 3. Photograph 1.

March 17, 1903, while trimming a recently spliced wire rope with a hatchet, his left eye was injured by a flying particle. He was treated by a local surgeon until the eye became quiet. A month after the injury he had a severe pain in the other (right) eye, which led him to seek advice in San Francisco.

My examination, April 30, 1903, showed the right eye to be perfectly normal. In the left eye there was a small corneal opacity at upper edge of pupil, traumatic cataract, posterior synechia at point corresponding to corneal macula. Vision $= 3/200$. Tension normal, eye perfectly quiet.

The x-ray revealed a small foreign body in the orbit. The repeated applications of the Haab magnet resulted negatively, as in Case 1.

Next day localization showed the particle to be in the eye below and slightly to the nasal side, probably resting in the posterior surface of the swollen lens. Not knowing the conditions I again applied the giant magnet, having the patient turn the eye as strongly upward as possible. After a moment he complained of pain; reducing the strength of the current I directed the patient to look straight ahead and then down, and

the piece of steel entered the anterior chamber. A small section was made in the cornea and the foreign body was easily extracted. No reaction or change of vision followed the operation.

This case again shows the unreliability of the Haab magnet to reveal the presence of a piece of steel in the eye; it shows the accuracy of this method of localization, but—what is here most important—the exact loca-

Fig. 10.—Case 3. Photograph 2.

tion of the foreign body enabled me to remove it successfully with the Haab magnet, while before localization I failed.

CASE 3.—Male, aged 36 years, blacksmith. Aug. 1, 1903, while striking a flying piece from the sledge hammer struck him in the left eye, destroying the sight instantly. Aug. 3, 1903, the patient was brought to my office by his oculist to try the effect of the giant magnet. Examination showed a penetrating wound of the cornea extending from the center down-

Fig. 11.—Case 3.

ward and slightly outward 8 mm., a corresponding injury to the iris and the lens, no reflex from fundus. Vision = 0, globe soft.

The result of application of the Haab magnet to the eye in various positions was entirely negative.

An x-ray photograph revealed the presence of a large foreign body in the orbit. Localization by the method detailed above showed the foreign body to be entirely outside the eye (see

diagram). Although it was not necessary, enucleation was done by the oculist in charge of the case as the eye was hopelessly blind, and the healing would require the patient's detention in the hospital, etc. Thus we obtained the proof that my localization was correct; the wound of exit of the foreign body was found in the posterior and inner wall of the globe. Both the wounds of entrance and exit were as clean cut as if made with a sharp knife. The axis of one wound being almost at a right angle to that of the other, showed that the foreign body twisted on itself in its passage through the eye. The wall of the eye was unusually thin.

This case again shows the necessity of exact localization, for it seemed quite improbable under the circumstances that so large an object would go entirely through the eye.

Because of the known position of the foreign body in the orbit the eye could have remained with perfect safety had a cosmetic result been of more importance than time and expense. The piece of steel in this case was 8 mm. in one diameter by 5 mm. in the other. We knew the foreign body to be quite thin from the faintness of the shadow in the x-ray picture, as well as from the clean-cut lines of the wounds.

751 Sutter Street.

STRICTURES OF THE FEMALE URETHRA.

WITH REPORT OF CASES.*

H. L. E. JOHNSON, M.D.

WASHINGTON, D. C.

Strictures of the female urethra are narrowings of the intima of the canal of a greater or less degree, which impede or prevent the passage of the urine by the natural course from the bladder. They are congenital or acquired, single or multiple, and may be classified as stenosis or obstructive, atresia or retentive, atresia, non-retentive, when the urethra becomes occluded from disuse associated with vesicovaginal fistula and atresia with urine discharged through the umbilicus via a patulous urachus. Various modifications of these varieties are found associated with malformation of the bladder.

Congenital atresia are usually located at the neck of the bladder, or in the upper third of the urethra, but obliteration of the entire canal, or its absence, has been noted. Acquired cases, if traumatic, are situated at or near the site of the injury, while, if non-traumatic, they are found usually in the middle third of the urethra.

Statistics are lacking as to frequency. The few authors who mention the condition agree that it is uncommon. About twelve years ago I read a paper before a Washington society, reporting a case of acquired stenosis, with retention, calling attention to the absence of statistics, and the large number of standard authors who made no mention of the disease. Some recent text-books mention the subject only casually, while others do not mention it at all.

The causes are congenital malformation, traumatic, internal or external, associated with abnormal, prolonged or instrumental labors. Injuries of the genitals from burns, scalds, falls, cicatrices following operations and contractions from free use of caustics. Inflammation and infection from gonorrhea, urethral chancre and chancroid, septic instruments, urethral masturbation, tumors and adhesion bands in vagina. Tumors within the urethra are said never to cause stenosis or atresia.

The pathologic alterations resulting from congenital malformation, inflammation, infection, pressure and trauma, may present any degree of structural change

* Read before the Tri-State Medical Association of the Carolinas and Virginia, Danville, Va.