the clamps, care should be taken not to exert too much pressure as it would interfere with the nutrition of the seat of operation and probably give rise to sloughing. Frequent cleansing should be carried out by means of a spray of Dobell's solution or other antiseptic wash. The clamp should remain in position for three or four days, after which I use plugs of carbolyzed oakum, as they are more cleanly than hard plugs of ivory or wood and exert sufficient pressure to hold the part in the required position. They should be changed at least once daily. In some cases it will be sufficient to plug only the formerly obstructed side with the oakum, after removal of the clamp. The evening temperature may rise above 101 degrees on the second and third days after operation, but soon subsides. The three following severe cases of deflected septum, all in male patients, and of traumatic origin, which I have recently operated upon by Delstanche's method with the most favorable results, both as regards the nasal stenosis and the facial deformity, have satisfied me as to the efficiency and value of this operation:

Case 1.—A. W., age 21, consulted me, Oct. 26, 1892, for complete obstruction of the right nostril and catarrh, and expressed himself as desirous of undergoing any operation that would give him relief from the distress he experienced at not being able to breathe through the right side of his nose. At about 5 or 6 years of age a severe fall upon the face had broken the nose, nothing having been done at the time to set it, so that from that time he could not breathe through the right nostril properly, and this difficulty increased in after years, until almost complete obstruction existed. Upon examination I found the right nasal chamber almost entirely occluded by deflection of the septum in both its cartilaginous and bony portions, and of undoubted traumatic origin. The left inferior turbinated body was greatly enlarged by puffy hypertrophy, yielding easily to pressure of the probe, and it almost filled the large concavity caused by the deformity of the deflected septum. In this case as there was so much inflammatory thickening of the projecting portion into the right nostril, I removed a piece about an eighth of an inch in thickness with the saw, in order to obtain space for the introduction of the flat blade of the forceps, as well as with the view of obtaining a better result after crushing. The blades of the forceps being placed in position in their respective nostrils, the anterior portion of the deflection was crushed and cut; this was repeated in two or three places back to the posterior part of the deviation, after which all resiliency was overcome, and the finger could be passed into the occluded nostril. The clamp was then placed in position and firm but not tight pressure exerted. The temperature rose to 101.2 degrees on the evening of the second day after operation, but soon fell to normal. On the morning of the fourth day I removed the clamp, and plugged both nares with carbolyzed oakum which was renewed daily. After two weeks' treatment the plugs were abandoned, firm union having taken place, and the patient could breathe freely through both nostrils. The facial deformity was also markedly reduced, the nose being now fairly straight. This patient returned by appointment in about a month from date of discharge when the relief of the stenosis was found to be permanent, and the catarrhal condition had nearly subsided. At this time the left inferior turbinated body was cauterized two or three times by galvano-cautery and the patient has since remained free from any obstruction or inconvenience from catarrh.

Case 2.—F. M., age 30, consulted me Feb. 3, 1893, for almost complete obstruction of the left nostril, very marked facial deformity and catarrh. About seven years previously he had received a severe blow on the right side of the nose. Upon examination I found the left nasal chamber almost entirely occluded, and post-nasal catarrh. The deviation was both cartilaginous and bony, and of unmistakable traumatic origin. I operated by Delstanche's method with the result that in two weeks the patient was breathing as freely through the left nostril as through the right, and upon inspection there was almost as much space. In this case also the deformity of the nose externally was decidedly improved and now, in appearance, is comparatively straight.

Case 3.—G. H., age 17, consulted me, April 6, 1893, for ob-

struction of the left nostril, which was causing him great discomfort. About five months previously, he had received a heavy blow on the right side of the nose. Upon examination I found marked deflection of the cartilaginous portion of the septum to the left, causing partial stenosis, and there was also post-nasal catarrh. The facial disfigurement consisted of rather an abrupt turning of the tip of the nose to the right. I did Delstanche's operation in this case with the most satisfactory results in regard to both the nasal stenosis and the external deformity. During the past year I have treated several additional cases by means of this operation and with equally good results.

A CASE OF RETENTION CYST OF THE FRONTAL SINUS.

Read in the Section on Laryngology and Otology, at the Forty-fifth Annual Meeting of the American Medical Association, held at San Francisco, June 5-8, 1894.

BY F. B. EATON, M.D.

PORTLAND, OREGON.

In January, 1893, an Italian aged about 26 years, consulted me concerning what appeared to be an osteoma at the upper and inner angle of the right orbit. Three years before, the skin over the right brow and upper lid became red and swollen. The redness disappeared in three weeks; the swelling remained. Six months later a soft swelling appeared where the tumor is. Sometimes there was pain over the right brow. Nose used to run before the swelling but has "dried up" since. The hard swelling has increased slightly in the past year. No specific history.

Objectively the right eye was pushed downward and outward and there was some proptosis. The hard tumor, about the size of a hickory nut, was not tender to the touch. Nasal respiration was impeded by a very large polypoid in middle turbinated bodies, the right being fully one-half inch wide. Believing this condition might be connected with that of right frontal sinus, I snared off with a powerful instrument the anterior portion of the right middle turbinated bone, the piece being the size of a hickory nut. That night a large amount of fluid escaped from the right nostril. Later I used a small electric light and found I could illuminate the left, but not the right frontal sinus when it was placed under the brow.

In consultation I made a small incision over the projection, ten millimeters above the tendo-oculi, and with an electric drill perforated the bone which proved to be thin. The drill entered a cavity which could be probed to great depth.

Two weeks later at the Good Samaritan Hospital I made an incision parallel with the eye-brow joining it by a vertical one at the inner end, exposed the bone and with a beak-shaped bone knife removed a piece of the bony shell one inch long and one-half inch wide. The sinus was found completely and tightly packed with a fatty granular substance, which was removed with a curette as rapidly as possible, there being profuse hemorrhage. There was no capsule. The sinus was as large as a good sized egg, extending far back, and its outer and lower wall was very thin. This cavity was packed with iodoform gauze, which could not be changed for several days, owing to the free hemorrhage which returned on each attempt. I decided later on to endeavor to obliterate the cavity by keeping it packed with gauze, since there was a continued tendency to mucopurulent discharge, owing to infection from the nose. I made an attempt to secure free drainage into the nose by passing a long slender silver probe by way of the infundibulum into the sinus. To the end I attached a bit of silk suture tying it to a strip of gauze which I drew into the nose. On the following day I found the gauze caused so much irritation that I withdrew it, and continued the daily packing with gauze, which, in about four months, resulted in obliteration of the cavity. The projection of the outer wall remained, however, as well as the displacement of the eye.

The substance removed, examined under the microscope, proved to be composed mainly of fusiform cells of an epithelial character, with some amorphous fatty substance.

DISCUSSION.

DR. MANSFIELD—I have had a case of a gentleman under my observation for the past five or six years with about the same amount of aphonia as the case reported by Dr. Eaton. He has not had catarrh, and being a vigorous male can not

be suspected of hysteria; he is also able to make a much louder and better voice when he sings in a high key than when speaking. He has paresis of the left vocal band probably of cerebral origin.

REPORT OF CASE OF TOTAL ADHERENT SOFT PALATE; OPERATION AND RECOVERY.

Read in the Section on Laryngology and Otology, at the Forty-fifth Annual Meeting of the American Medical Association, held at San Francisco, June 5-8, 1894.

BY J. E. SCHADLE, M.D.
ST. PAUL, MINN.

Eighteen months since, James S. consulted me in reference to complete stenosis of the nasal passages and deafness which had existed for a period of seventeen years. On examination of the nasal passages they were found to contain polypoid growths varying in size, which by their presence seemed to annoy considerably the patient by creating a continuous flow of serous fluid from the nostrils and bringing on attacks of violent sneezing, which condition ceased soon after the tumors were removed.

Inspection of the throat revealed partial obliteration of the pillars of the palate, entire loss of the uvula, and total adhesion of the velum palati to the posterior wall of the pharynx. What still remained of the soft palate consisted chiefly of cicatricial tissue as was evidenced by the firmness and density of the structure. After explaining to the patient the character of his trouble, informing him of the difficulty surrounding such an operation and advising him against one being performed, he left me, determined to seek advice elsewhere. Two months ago he returned to me and insisted on my giving him relief, if it be possible.

The patient is a book-keeper by occupation, was born in Manchester, England, and came to the United States in 1885. As to family history; mother died early in life from the effects of puerperal septicemia and the father three years ago from an attack of gout. The subject of this sketch is the eldest of six children all of whom except himself are in the enjoyment of good health. The history is negative, so far as tubercular or malignant diseases are concerned. Mr. S. is now 32 years of age, married, weighs 140 pounds, is in a general way robust and healthy and complains of the local malady only, on account of which he seeks advice. He denies ever having had acquired syphilis. When 10 years old he developed a sore throat which continued to disturb him for some months when the family physician was consulted, who pronounced the disease "an ulcerated sore throat" and made repeated applications of lunar caustic. The morbid process of the throat seemed to have remained unarrested as is shown by the fact that at the age of 14, interference of nasal respiration was markedly experienced, and at 16, the sense of smell was lost, the function of the ears was impaired and the power to breathe or blow through the nose was totally abolished.

Cocainizing the parts to be operated upon as thoroughly as possible, I divided the adhesions by the use of an angular probe-pointed knife and curved scissors such as are employed in gynecologic work.

After puncturing on either side of the adherent parts with a sharp-pointed bistoury, I insinuated through the opening thus made the probe-pointed

blade of the angular knife and severed the attachments by first cutting downward to the lowest point of adhesion, thence upward to the median or uvular line where the adhesions were freed with the scissors.

These different procedures effected a communication between the naso- and oro-pharynx of sufficient spaciousness for me to introduce my index finger into the post-nasal cavity for purposes of exploration. By means of this manipulation, numerous bands of adhesions extending from the superior surface of the palate to the vault of the naso-pharynx were discovered, which bands I divided with the knife guarded by the finger. The loss of blood was slight. The patient was at once enabled to blow his nose with entire freedom and nasal respiration was restored. In a few days after the performance of the operation, a sense of smell in a measure returned and hearing without any other aid became perceptibly improved. Regarding after treatment which I deem most essential, the plan adopted to keep the denuded surfaces from reuniting, was as follows:

A piece of iodoform gauze, two inches wide, fifteen inches long, was in half its length divided, which parts were respectively drawn through the nostrils from behind the palate forward, thus by means of slight traction bringing the crotch of the tape to rest firmly against the posterior extremity of the septum narium. The broad or oral end of the tape was brought forward through the mouth, and united with the ones passing through the nose over the upper lip. This appliance was well borne by the patient, its presence interfering somewhat with the acts of deglutition and phonation. Apart from this inconvenience no demurrer was made. The dressing was changed every twenty-four hours for two weeks, after which it was omitted during the day and worn through the night only. Mono-chloroacetic acid (Smith and Delavan) was applied occasionally to the raw surfaces to insure if possible a favorable issue. Whenever a point of adhesion would show itself, it was immediately broken up. The nostrils and post-nasal cavity were kept clean by the employment in spray form of antiseptic washes.

REMARKS.—1. The cause of the ulcerative processes of the soft palate followed by adhesions, was inherited syphilis. This opinion is confirmed by the fact that the patient had typical Hutchinson teeth.

2. The operation is justifiable and made successful by effecting a thorough division of the points of adhesion, and devoting most zealous attention to all details of after treatment until recovery takes place.

TREATMENT OF HAY FEVER.

Read in the Section on Laryngology and Otology, at the Forty-fifth Annual Meeting of the American Medical Association, held at San Francisco, June 5-8, 1894.

BY A. C. WOLFE, M.D.
COLUMBUS, OHIO.

Bostock, himself a sufferer from hay fever, gave the first detailed account of this disease in 1819. In 1869 Helmholtz held that the symptoms were produced by vibrios existing in the nasal passage. In 1873, Blackley, by carefully conducted experiments, was led to believe that the pollen of grasses and flowers was the exciting cause of hay fever. In 1883, Roe advocated that hyperesthesia is associated with, or occasioned by a diseased condition, either latent or active, of the naso-pharyngeal mucous membrane. Harrison Allen attributed the disease to permanent