traction of teeth these methods are applicable. In uterine hemorrhages, due to different causes, large irrigations with a 5 per cent. solution should be terminated by the application to the neck of the uterus of a tampon wet with the same solution. In hemorrhage due to endometritis fungosa, equal parts of salol and antipyrin are melted at the fusion-point of salol, which gives a brown, syrupy liquid, and this liquid is used to moisten a tampon which can be placed into the uterine neck, a speculum being used, and over it a few other tampons. After amygdalotomy moistened tampons are efficacious. For hæmoptysis, inhalations or sprays of a 1 per cent. solution have been effectual when other means, including subcutaneous injections of ergotin, have failed. For open wounds a solution or a gauze impregnated with the powder coagulates the blood, which resists for a long time decomposition, because of the antiseptic and cicatrizing properties of the drug.—*Journal des Praticiens*, 1895, 2e semestre, No. 19, p. 289.

THE ABSORPTION OF THE ALBUMINOIDS AND FATS OF A NUTRITIVE ENEMA.

DR. R. LÉPINE, reviewing the physiological literature, acknowledges that the question, whether absorption takes place in the large or small intestine, remains open, because the ingenious hypothesis of Grützner, that there is a retrograde transportation of fine alimentary particles without visible antiperistalsis, has not yet been established. If this should fail of proof, it is strongly probable that some of a nutritive injection could pass through the ileocecal valve, and thus would be partly explained the fact that a patient can live and keep his weight entirely upon rectal enemata, for all that the large intestine does not show digestive phenomena which are hardly appreciable. Physiological observation shows that the ileocecal valve is permeable, and it is only necessary for nutritive substances to penetrate a certain distance into the ileum to meet with digestive juices which can change them and render them absorbable.—*La Semaine Médicale*, 1895, No. 45, p. 389.

IZAL.

MR. SHERIDAN DELÉPINE makes a laboratory-report upon this compound, which is comparatively insoluble and non-volatile at the ordinary temperature. It can be freely administered internally, used over extensive wounds, or injected under the skin without bad effects, and, finally, does not damage surgical instruments. The organisms upon which he has used this drug are: (1) bacillus tuberculosis (hominis); (2) bacillus coli communis; (3) staphylococcus pyogenes aureus; (4) bacillus anthracis (sporing-stage). Summing up the results of his observations he concludes that when diluted with 100 or even 200 parts of water it is a powerful and reliable antiseptic when contact of a sufficient length of time is secured. As an antiseptic it is more powerful than carbolic acid, and there is little doubt that it possesses immense advantages over it in many directions.—*The Medical Chronicle*, 1895, No. 6, p. 434.

[Izal is said to be a by-product obtained in the process of coke-formation. It is also known as Thornccliffe disinfectant.—R. W. W.]