CASES 6 AND 7.—J. S., coxswain, aged 28 years, presented reducible oblique inguinal hernia on both sides. The right was said to have resulted from exertion on a boat davit, eighteen months before operation, and the left appeared one month before. Both have been continuously retained with trusses. His general condition was excellent, and double operation was performed on July 27, 1899. The patient reacted well from the ether. No sac was demonstrated on either side. The post-operative condition was normal, and the wounds united by first intention. He is still under observation, but the result is perfect.

The herniotomies that were recorded in 1898 were performed admirably, by Assistant-Surgeon James R. Whiting, U. S. N., a satisfactory conclusion resulting in each case; all the men having been returned to duty, and no relapse has as yet been reported in any case.

The herniotomies of this year were all performed by Passed Assistant-Surgeon Charles F. Stokes, U. S. N. —my executive surgeon—and I can not too earnestly commend his care and skill, and do honor to his admirable technic and successful result in every case.

VASOMOTOR SYSTEM, AND THE IMPORTANCE OF THE DRUGS WHICH ACT ON IT, IN OPHTHALMOLOGY*.

BY DAVID W. STEVENSON, M.D.

RICHMOND, IND.

A study of the pulse and vascular tension is of the utmost importance to the ophthalmologist. Although clinical reports and practice show that while very slight local eye symptoms are noticed, yet how seldom is the pulse recorded, even previous to a cataract operation or during the crises of a glaucoma. It would be well if all oculists would always report the tension of the pulse with the same care, and use the same signs which are used in regard to the eye. The scale would thus run from $+3$ to -3 .

Rolling the artery under the fingers with varying pressures, or carrying the skin along longitudinally over the vessel, will decide the tension as well as the inequalities of thickness and density.

Ewart, in a classical work on pulse sensations, has proved that the tension may be best noted by the amount of pressure that will cause the pulse impression to pass from the distal to the proximal side. Although tension is the one attribute of supreme importance in the pulse, there are at least four others that must be noted, frequency, size, the condition of the arterial wall, and the characteristic shape of the pulse curve; whether its access is gradual or sudden, its crest's duration long or short, its subsidence abrupt or slow; note being taken whenever it is dirotic.

The great vasomotor system is controlled by both augmentor and inhibitory nerves to the heart; as well as the vasoconstrictor and vasodilator nerves to the vessels. The augmentor and vasoconstrictor nerves are largely associated in their course with the sympathetic system.

If it is essential to the oculist to be able to estimate the pulse tension, it is of more importance to be acquainted with the drugs that will control this tension. Inflammation causes a paralysis of the vasoconstrictors, and the area involved is swollen with sluggish and poison-laden blood which, pressing on the nerves, causes pain and further obstruction. The sensible thing is to bleed the patient into his own vessels so that the hyper-

emic area will be drained of the sluggish fluid. Vasodilators, such as aconite, are used in tonsillitis, and the result is better drainage and less pain in the diseased parts. There is not as much fever, ordinarily, in the inflammations of the eye as in the tonsil, yet the vasodilators, properly handled, will do a surprising amount of good.

There is need of such remedies as bromid of sodium, acting as a vasoconstrictor of the vessels of the eye and brain, and of aconite to dilate the general vascular system; also strychnin, the tonic and regulator for the nervous system, and which prevents any depression that is apt to follow the use of any vasodilator. Such remedies in addition to saline laxatives and the local use of eserine may obviate the necessity of an iridectomy in glaucoma.

Opium and potassium iodid are efficient vasodilators. Dr. George F. Butler says he uses five minims of the deodorized tincture of opium every four hours for months. The dose is so small, he says, that there is no danger of forming a habit. Potassium iodid, so often used in eye diseases with such good result, may be efficient, not that the patient is always syphilitic, but that the contracted vessels are dilated.

Hare gives a list of vascular depressants in the order of their efficiency; amyl nitrite, nitroglycerin, veratrum viride, antimony, aconite, alcohol in large doses and jaborandi.

Haig has shown how the xanthins and uric acid in the blood act as marked vasoconstrictors. This shows how sodium salicylate and mercury favor the dilatation of the vessels under such circumstances and prevent high tension.

Eyes operated on for cataract have the tension suddenly removed from the choroidal and retinal vessels. The wonder is that there are no more cases of disastrous hemorrhages after the operation.

Wide dilation of the trunk vessels, with contraction of the vessels of the brain by the bromids, is the best preventive, of which all such patients should have the benefit. It would be safer if all operations opening the eyeball had this preliminary treatment.

Bartholow states that nitroglycerin stands in the front rank of remedies for anemia, and that a much larger quantity of blood is obtained by the tissues, while the mere giving of iron is one of the most common of therapeutic fallacies of the day, because of the contracted vessels. Nitroglycerin is transient in its effects and ought to be given in broken doses of one-thousandth of a grain, often repeated. Sluggish conditions of the retina, like retinitis pigmentosa and optic nerve atrophy ought to have this treatment, and it promises the best results in such delayed conditions as slow corneal repair after cataract extraction. Strychnin will be added to this, because it prevents side-tracking of the waste material and venous blood. Its tendency is to balance-wheel the body mechanism in all its parts, following an inflammation or a degeneration. It is like replacing the telegraph lines after a storm. It increases connections. The rule again is broken doses often repeated and effects carefully watched.

Atropin dilates the vast cutaneous system, and is thus one of our most efficient remedies for internal bleeding from nasal or other internal organs. As it dries up glandular secretion, it is suitable in bleeding from mucous membrane or internal glands. As to pilocarpin, experiment proves that it acts by paralyzing the vasomotor system. The increased action of the heart and flushing of the skin is due to the dilation of the arteri-

*Presented to the Section on Ophthalmology, at the Fiftieth Annual Meeting of the American Medical Association, held at Columbus, Ohio, June 8-9, 1899.

oles. DeWecker has shown how it brings about absorption and resolution in inflammatory eye diseases with effusion and exudation. DeSchweinitz has also shown its wide field of usefulness if used in broken doses.

In albuminuric retinitis and allied conditions, the salvation of the retina, as well as the life of the patient, may depend on the reduction of tension by the general dilation of the vascular system.

INTRANASAL ANGIOMA: BLEEDING POLYPUS OF THE SEPTUM.*

BY W. E. CASSELBERRY, M.D.

Professor of Laryngology and Rhinology in Northwestern University Medical School; Laryngologist to St. Luke's and Wesley Hospitals, etc.
CHICAGO.

A typical *angioma simplex* is composed of new-formed, very tortuous blood-vessels, held together by a small amount of connective tissue and situated usually in the skin, e. g., nevus. The normal vessels of the part where it is situated may enter more or less into its formation, becoming dilated and tortuous, and it has therefore become customary to include in the class of angiomata tumors composed entirely of dilated pre-existent vessels, although the term telangiectasis or telangiectoma is by some preferred to this sort of angioma where dilatation is more prominent than actual new formation.

FIGURE 1.

A typical *angioma cavernosum* is composed chiefly of blood spaces lined with endothelium, separated by a stroma of connective tissue. The spaces contain venous blood, but seem to represent capillaries, being interposed between arteries and veins. The structure is similar to that of the corpus cavernosum. Such are often found in the liver.

The vascular tumors which grow on the septum narium do not conform exactly to either of these types. They contain both blood-vessels and blood spaces, thus far representing rather a combination of the two types, and they contain also a larger proportion of reticular connective tissue, with either round or spindle cells. In some this reticulum has a fibrillary character justifying the term angiofibroma. Among themselves,

however, as occurring in this situation on the septum narium, they do present a tolerable uniformity of structure, the differences being variations in degree rather than of kind and, as their salient characteristic is extreme vascularity, they are best grouped under the term "intranasal angioma." The condition is exemplified in the following case:

Mrs. T., aged about 30, married, pregnant, has been annoyed for the past few months by frequent bleeding from the right nostril, and gradually increasing obstruction. Examination disclosed a soft, reddish, irregularly nodulated tumor the size of a small bean, attached by a rather broad pedicle to a small excrescence of the septum at the point of junction of the cartilaginous segment with the septal process of the superior maxilla, (Fig. 1) just within the nostril and close to the floor. It bled easily on contact with a probe. I removed it with a cautery snare and cauterized the base with chromic acid. Three weeks later one small vascular point, still unhealed and disposed to bleed, was again cauterized with chromic acid. There has been no recurrence for two years. The microscopic examination, made by Dr. R. H. Harvey, shows the tumor to be composed largely of blood-vessels and blood spaces. The blood-vessels vary in size and most of them contain more or less blood. Some are collapsed. The blood spaces are partly lined with epithelium. Between these vessels and spaces is connective tissue in which are numerous round

FIGURE 2.

and young connective tissue cells. On one edge of the tumor is a remnant of mucous membrane. The diagnosis is angioma. Microphotographs of sections under high and low power are appended (Figs. 2 and 3).

I can not think that these tumors are as rare as has been represented, although they have attracted little notice until recently, when, following Victor Lange¹ in 1892, papers have appeared by Schadowaldt, Alexander, Sheier and Heyman, all in the *Archiv für Laryngologie und Rhinologie* (Band I, 1894) under the title of bleeding polypus of the septum. Their detailed pathologic descriptions tally so closely with each other and also with that of my own case as above given, that they must all be the same sort of tumor, notwithstanding one is spoken of as soft fibroma, and another showing some glandular tissue as lymphangioma telangiectaticum. Since then Pierce² has reported a case under the name

*Presented to the Section on Laryngology and Otology, at the Fifth Annual Meeting of the American Medical Association, held at Columbus, Ohio, June 6-9, 1899.