that vital organ upon which a high temperature exerts such a destructive influence. We must therefore look elsewhere for the immediate cause of death; and every item of evidence, whether it be pathological or clinical, points to the same inevitable conclusion—namely, that the fatal termination of acute pneumonia is the direct result of cardiac failure.

There are two general factors which are capable of producing the condition of cardiac insufficiency—an increased resistance to the propulsive action of the heart, and a progressive deterioration of its muscular substance. And in acute pneumonia both these factors are conspicuously present; the former in the increase of tension in the pulmonary circuit consequent on the consolidation of a portion of the lung, and the latter as the direct result of a high temperature. The action of these two forces is to compel the heart to beat more forcibly and more quickly, while at the same time it is steadily deprived of the power to do either. In the asthenic variety of the disease the heart is intrinsically feeble from the outset, and hence the malign influence of fever is an unnecessary element in the production of its insufficiency.

That these *a priori* considerations are in accordance with actual facts has already been shown, for it has been proved that death is most liable to supervene at two periods in the disease: about the fifth day, when the fever is at its height, or at the crisis, when cardiac collapse is apt to result from its sudden withdrawal. One step further. Two great methods of treatment have been examined in detail as to their influence upon the mortality from the disease. In the one we find that the chief remedies were aimed at relieving the condition of the lung; and while stimulants were freely administered with the object of whipping up the flagging heart, the cause of its physical lameness was allowed to proceed unmolested. The result was that among 552 cases so treated the mortality exceeded 23 per cent., although alcohol was exhibited in no less than 70 per cent. In the second case a method of treatment was adopted with one special object: to economise cardiac force by minimising the injurious influences of fever. In 108 cases of similar severity to the foregoing the treatment consisted in the systematic reduction of temperature by means of sponging or ice cradling. Of this number only 45 (41 per cent.) received alcohol, and only 10 per cent. died.

Again, among the former class 46 deaths resulted from collapse at the crisis of the fever. Among the latter, where special attention was paid to this source of danger, not a single death is recorded from this cause. Nothing could be more conclusive, for these facts show to what a considerable extent the mortality from acute pneumonia may be reduced when prophylactic treatment is applied with the view of strengthening that vital position against which death almost invariably concentrates the force of his attack. And the success which has attended this rational treatment of acute pneumonia appears as a still further argument in favour of the specific febrile nature of the disease; for it would indicate that the consolidation of the lung is after all but an analogue of the typhoid ulcer and of the scarlatinal sore-throat—a characteristic result, but never the cause, of the disease. It has also been shown that, firstly, the quantity of albumen in the urine is of considerable prognostic value; secondly, that the crisis is often a period of great danger to life; and thirdly, that those cases which commence with a severe gastro-intestinal attack are twice as liable to end fatally as those which exhibit the more usual initial rigor.

In conclusion, I beg to thank the various members of the medical staff of the London Hospital for their kind permission to publish in this manner the results of their cases.

WEST KENT GENERAL HOSPITAL.—The fifty-eighth annual report records the munificent gift during the year of the Dowager Lady Howard de Walden. The entire top floor had been very completely fitted up by her ladyship as a women's ward for eight beds, together with a ward for special cases and bedrooms for the staff. Ample funds (£10,000) had been placed in the hands of trustees as an endowment. During the year there had been a considerable increase in the number of in-patients. A generous gift of £200 had enabled the committee to avert any addition to the adverse balance.

Clinical Notes:

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

A CASE OF GASTROSTOMY.

BY HENRY J. ROPE, F.R.C.S.

W. R.—, aged sixty-four, a waiter, was admitted to the Salop Infirmary on Sept. 15th, 1890, complaining of difficulty in swallowing for the last three weeks. Cannot take anything solid, and fluids come back through the nose. Rapid emaciation; weight, 7st. 10lb. Has been a free drinker. Has a soft blowing systolic murmur. (Esophageal bougie passed thirteen inches, and found a hard unyielding obstruction. The patient was put on fifteen grains of iodide of potassium three times a day. Sept 17th: He retains about half of the fluids that he takes. He cannot retain puddings or arrowroot.—Oct. 4th: He feels much better, and takes exercise daily. Bougie has been passed at frequent intervals. and to-day was passed fifteen inches.—19th: He does not complain of any pain, and is always anxious to have the bougie passed, as he says it gives him great relief. He seems weaker. Weight, 7st. 6lb.—28th: He complained of pain when the bougie was passed, referring it to the right mammary region. He does not retain his food so well, and is weaker. Weight, 7st. 3lb. Is unable to leave his bed.—31st: Gastrostomy (first stage) was performed under ether. The stomach was easily found, and was attached to the abdominal wall by Mr. Howse's method. He rallied well, but complained of great pain.—Nov. 1st: No pain; temperature normal; mouth very dry. Dressing soaked; fresh gauze packed over it.—2nd: More pain; mouth still very dry. Pulse 90, weak, respiration 30.—Nov. 3rd: Cough troublesome, with profuse expectoration of very tenacious, non-aerated sputa.—4th: Dressings removed; wound perfectly sweet, edges almost in contact. The edges were separated and held apart, and the stomach was pinched up and carefully opened with sharp-pointed scissors; two ounces of strong beef-tea and milk were at once introduced through a catheter, to the intense satisfaction of the patient, giving him a very pleasant sensation. The catheter was tied in, and he was fed every two hours. At each feeding a slight flow of saliva was noticed in the mouth.—5th: Slept well and is cheerful. On coughing some fluid escapes through wound, bile-stained, very acid.—6th: Slept well, but is weaker; more cough, more escape of gastric juice. Skin round wound red and irritated. Rubber tracheotomy tube used instead of catheter.—7th: The introduction of food causes a cough, which expels some of the food again; nutrient suppositories at frequent intervals.—8th: Pulse quicker and weaker; temperature 101°, delirium, jactitations; free return of food through wound.—9th: Temperature 102°, pulse almost imperceptible; Cheyne-Stokes breathing. He died the following morning.

Necropsy (Nov. 11th).—Great emaciation; well-marked rigor mortis. Stomach firmly adherent by strong adhesions all round the wound, of normal size and thickness, containing a small quantity of brownish fluid. Intestines much contracted. Kidneys small; capsules adherent. Aorta atheromatous; aortic and mitral valves puckered. The upper part of the œsophagus was dilated and much thickened. Three inches and a half above its cardiac end was a tight stricture, barely admitting a No. 8 English catheter; and here the œsophagus was greatly thickened and as hard as cartilage. There was ulceration of the mucous membrane for half an inch above the stricture, with perforation into the root of the right lung, where there was a sac, the size of a filbert, containing grumous fluid similar to that found in the stomach. The bases of the lungs were œdematous. The mediastinal glands were enlarged.

The patient was persistent in his opposition to any operation, and when at last his consent was obtained he was too weak to derive more than a few days' advantage from it. The notes were kindly supplied by Mr. Harold McLeod.

Shrewsbury.