valve formation at the opening of the ureter into the pelvis. An incision was carried through the spur or valve formation downward onto the pelvis and ureter an equal distance, about one and a half centimeters (see Fig. 8), the edges of the incision were then pulled apart at their mid-points, and the edge of the incision in the pelvis sutured to the ureter in a transverse line. (See Fig. 9.) Richardson made this an extrapelvic operation rather than transpelvic, as carried out by Fenger.

FIG. 10. Uretero-pyeloplasty. Longitudinal incision a-b in Fig. 9 made into a transverse incision and sutures placed.

Fenger's first operation simply made a nick in the spur and made no attempt to lower the opening; the second operation, as carried out by Richardson, did lower the opening somewhat, but this was not a object of the operation. Fenger's first operation could hardly be dignified by the name of uretero-pyeloplasty, but the second, as carried out by him and Richardson, should be so called.

REFERENCES.

- ¹ Simon: Chir. der Nieren, 1871.
- ² Landau: Berl. klin. Wochenschr., 1888, pp. 47 and 48.
- ³ Terrier and Baudoin: Rev. de Chir., 1891, pp. 719, 833 and 1055.
- ⁴ Duval et Grégoire: L'Asso. Française d'Urologie, Paris, 1906, p. 1.
- ⁵ Küster: Arch. für klin. Chir., 1892, vol. xlv, p. 850.
- ⁶ Moynihan: Brit. Med. Jour., 1904, April 30.
- ⁷ Bazy: Rev. de Chir., 1903, pp. 1-27.
- ⁸ Tuffier: Ann. des Mal. des Org. Gén.-urin. 1894, vol. xii, p. 14.
- ⁹ Fenger: Jour. Am. Med. Assn., 1894, vol. xxii, p. 335.
- ¹⁰ Gardner: Opérations dans le Traitement des Retentions du Rein, Paris, 1904.
- ¹¹ Ekehorn: Arch. für klin. Chir., 1907, vol. lxxxii, p. 955.
- ¹² Merkel: Virchow Arch. für Path. Anat. und Physiol. und für klin. Med., 1908, Bd. 191, Folge 19, Seite 534.
- ¹³ Brewer: Med. News, 1897, July 31, p. 128.
- ¹⁴ Carlier: L'Asso. Française d'Urologie, Paris, 1906.
- ¹⁵ Albarran: Ann. des Mal. des Org. Gén.-urin., 1896, vol. xiv, p. 987.
- ¹⁶ Trendelenburg: Volkmann's Sammlung Klin. Vorträge, 1895, no. 355.
- ¹⁷ Küster: Arch. für klin. Chir., 1892, p. 850.
- ¹⁸ Fenger: Jour. Am. Med. Assn., March 10, 1894.
- ¹⁹ Richardson: Trans. Am. Surg. Assn., 1897, p. 555.
- ²⁰ Israël: Chir. klin. der Nierenkrankheiten, 1902.

(To be continued.)

TREPONEMATA (SPIROCHÆTÆ) IN SYPHILITIC AORTITIS. FIVE CASES, ONE WITH ANEURISM.

BY JAMES HOMER WRIGHT, M.D., S.D.,
AND

OSCAR RICHARDSON, M.D.,
(From the Pathological Laboratory of the Massachusetts General Hospital.)

THE essential points in the histories and gross anatomical appearances of the lesions in the cases recorded here are as follows:

CASE I. Service of Dr. F. C. Shattuck and Dr. R. C. Cabot. A man, thirty-six years of age. Eight years before entrance to the hospital he had chancre and buboes. There was a slight scar on the glans penis.

The heart weighed 802 gm., and showed marked hypertrophy and dilatation, the latter condition being more marked in the case of the left ventricle. The coronary arteries were free and quite smooth. All the valves showed considerable increase in circumference, and excepting the aortic they were quite smooth. The aortic cusps showed much fibrous change which greatly deformed the valve and increased its circumference. (See Fig. 1.)

The aorta showed marked areas of aortitis. These areas extended down as far as the aortic cusps and the process in the wall of the aorta was apparently common with that in the cusps.

CASE II. This case was kindly furnished us by Dr. George B. Magrath, Medical Examiner for Suffolk County. A man, about thirty years of age, with no definite history of syphilis.

The heart, in this case, showed no particular enlargement, and the aortic cusps presented only a slight amount of fibrous change.

The aorta was the seat of numerous areas of aortitis. These areas extended down as far as the aortic cusps, and the process in the wall of the aorta almost occluded the opening of the right coronary artery.

CASE III. Service of Dr. H. F. Vickery. A man, thirty-nine years of age, who had had gonorrhea three or four times, the last, five or six years ago, when he also had soft chancres. He denied ever having had a hard chancre, but sixteen years ago had a non-itching rash, extending over the body, with sore throat and alopecia. Mercury was administered for two or three months with benefit. It was also stated that he had buboes eighteen years ago.

The heart weighed 710 gm., and showed marked hypertrophy and dilatation, the latter condition being more marked in the case of the left ventricle. The coronary arteries were free and quite smooth. The valves were not remarkable, except in the case of the aortic, which showed marked fibrous changes with a few very small areas of calcification, all of which greatly deformed the valve. There was no particular increase in the circumference of this valve.

The aorta presented many areas of aortitis. These areas extended down as far as the aortic cusps and the process in the wall of the aorta was apparently common with that in the cusps.

CASE IV. The heart in this case was a museum specimen, and we are without any history in regard to it.

The organ weighed 563 gm. and showed considerable hypertrophy and dilatation, the latter condition being more marked in the case of the left ventricle. The coronary arteries were free and showed only a few small fibrous plaques in the intima. The valves, other than the aortic, require no mention. The cusps of the aortic valve showed a moderate amount

of fibrous changes and the circumference of the valve was moderately increased.

The aorta showed many areas of aortitis. These areas extended as far as the aortic cusps, and the process in the wall of the aorta was apparently common with that in the cusps.

CASE V. Service of Dr. R. H. Fitz. A man, forty-three years of age. Venereal disease denied.

The heart in this case was not especially enlarged, although it was perhaps slightly larger than normal. The organ showed nothing for record except in regard to the aortic valve, which showed some fibrous change in its cusps, with a moderate amount of increase of its circumference.

The ascending portion of the aorta and the aortic arch showed considerable increase in their circumferences. The internal aspect of the aorta in these regions presented numerous furrows, roughenings and yellowish discolorations. Just above the upper margin of the sinus of Valsalva, from which the right coronary artery originates, there was a more or less irregular opening in the ascending portion of the aorta, which measured 4 cm. in diameter. The edges of this opening were rounded off and were continuous with the lining of the aorta and with the lining of an ovoid sac measuring 6 cm. in greatest diameter. This sac was lined by a firm, grayish-white tissue, resembling the furrowed lining of the aorta. The sac projected forward and somewhat downward on to the external surface of the right ventricle and right auricle, which latter rested behind it and above it. It thus projected into the cavity of the pericardium. At a point at or near the posterior aspect of the sac the wall was considerably thinned and generally less than 2 mm. thick. In this thinned area in the wall of the sac there was a small opening about 2 or 3 mm. in greatest dimension with ragged red margins. Through this opening in the wall of the aneurism a hemorrhage into the pericardium had taken place. What appeared to represent the right coronary artery had its origin just at the margin and at the inferior left border of the opening of the sac in the ascending aorta. This vessel was somewhat smaller than usual for the right coronary artery. The thoracic and abdominal portions of the aorta showed some but not an excessive amount of furrowing and yellowish patching of the intima.

It is apparent, from the above data, that the condition of aortitis is common in all of the cases, and that certain other conditions are associated with most of them. Thus, the ages range between thirty and forty-three years, and hypertrophy and dilatation of the heart is marked in three of the cases. The dilatation is most marked in the cavity of the left ventricle and in all of the cases the aortic valve shows more or less well-marked fibrous changes. There are only a few small areas of calcareous change, and this in one of the cases. The other valves in all of the cases are not remarkable. The coronary arteries are not remarkable except as their openings are more or less occluded by the fibrosis in the walls of the sinuses of Valsalva.

The lesion of aortitis, which is present in all of the cases, presents certain quite definite characters. The regions involved are mainly those of the first portion and the arch. Beyond this the areas of change in the wall of the aorta are much less marked, fewer in number, and in the abdominal portion they have practically disappeared or give place to small sclerotic

areas, indistinguishable from those of ordinary arteriosclerosis. The areas of aortitis are mainly fibrous in character, and it is only in one case that any calcareous element enters into the process. The wall of the first portion of the aorta, extending up from the region of the cusps, shows the typical condition. In this situation the intima presents smaller and larger, irregularly margined, discrete and confluent areas, the surfaces of which are thrown into irregular, smaller and larger areas of elevation with furrows, and smaller and larger areas of depression between. This change in the intima gives it a grayish, granular aspect. Between these areas of change in the intima, areas of quite normal appearing intima may intervene. On cross section through one of these typical areas of aortitis, the wall of the aorta is always found to be markedly thickened. In instances the wall may reach a thickness of 7 mm. The elevated areas in the intima are seen to consist of rather dense, firm, grayish, fibrous-like tissue, which rests over a layer of rather soft, yellowish, stringy, necrotic-like tissue, extending into the media for varying depths. The media is thickened and between and about the necrotic-like areas mentioned appears as fairly firm, yellowish, homogeneous tissue, resting over and along a thickened grayish, slightly translucent, somewhat fibrous-looking layer of adventitia. On the whole the media appears macroscopically to be the coat which is involved to the greatest extent by the process. We find no definite changes to record in the appearance of the great branches of the aorta.

Microscopic examination of sections from the aortic wall shows that the lesions in all of the cases are essentially the same and like those described as characteristic of syphilitic aortitis.

The most striking feature is the extensive transformation of the media and intima into fibrous tissue and the degeneration and necrosis of this tissue. In places there are areas of infiltration with plasma cells and lymphocytes. The degenerated and necrotic tissue in Cases I and II is infiltrated in places with polynuclear leucocytes. (See Fig. 2.)

The adventitia is more or less extensively thickened and is infiltrated in places with plasma cells and lymphocytes. The vasa vasorum show proliferative endarteritis.

The intima in places shows much fibrous thickening, and in Case I contains many fibroblasts, together with some giant cells.

Thus the process as a whole, in these cases, seems to be of long standing, but in places it is apparently recent, as is shown by the occurrence of necrosis in the media without fibrous changes.

In sections from all of these cases prepared by Levaditi's first published method, treponemata identical in morphology with *treponema pallidum*, shown by the same method in syphilitic lesions, were found. They were found practically only in or about the degenerated and necrotic areas of the tissue in the media and intima. In only one instance was one found in a granulomatous area in the media.

FIG. 1. CASE I. Showing the gross appearances of the lesions of the wall of the aorta and of the aortic valves. The very marked hypertrophy and dilatation of the left ventricle is shown by comparison with the centimeter scale.

FIG. 2. CASE I. Low-power photograph of a section of the wall of the aorta through a portion corresponding to the elevated areas in the intima just above the aortic valve. The necrosis of the media and of the new-formed connective tissue as well as the marked infiltration with polynuclear leucocytes is shown. Some of the larger dots in the hyperplastic intima are giant cells.

FIG. 3. CASE I. Treponemata in degenerate new-formed connective tissue of the intima. The field lies immediately beneath the surface of an elevated area in the intima similar to those shown in the photograph of the heart near the aorta just above the aortic valve. Many treponema are not in focus and some are only indicated by obscure shadows or blurs.

Photographs by L. S. Brown, Pathological Laboratory, Massachusetts General Hospital.

FIG. 4. CASE IV. Treponemata in the midst of necrotic tissue in the wall of the aorta.

FIG. 5. CASE V. Treponemata in a necrotic portion of the aortic wall near the site of an aneurysm.

Their numbers varied greatly in the different cases and in different places in the lesions. In and about many areas of necrosis none at all could be found. When present in a given area, they were usually few in number, but in Case I, in an area of degenerated hyperplastic intima, they were present in enormous numbers throughout a portion of the area (see Fig. 3), and in Case II they were fairly numerous. In Cases III and IV they were found only in small numbers and atypical forms of the micro-organism predominated. In Case V only atypical forms were found in small numbers. Some of these atypical forms were of the same length and width as the typical treponema, but had less numerous and less abrupt curves or windings and sometimes presented a granular appearance; others were like the typical treponema except that they were shorter. Inasmuch as we found these atypical forms, not only along with the typical treponema in the other cases, but also in the liver of congenital syphilis among abundant typical forms, we consider them to be degenerated forms of the treponema. That they are not fragments of the tissue fibrils we have convinced ourselves after comparing them with such products in the same sections.

From the study of the microscopic appearances in these lesions, we consider the process to be one of primary necrosis of the media, and probably also of the intima, with the development of reparative connective tissue, followed by secondary degeneration and necrosis of this connective tissue. The finding of the treponemata in an area of primary necrosis in the media, as well as in association with the necrosis in the fibrous tissue, seems to justify the belief that the micro-organisms are the cause of the necrosis by their local action upon the tissue, and that they are to be regarded as the cause of the whole process, although the possibility that they are merely secondary invaders cannot be denied. Their numbers and distribution in the lesions would suggest that they rapidly multiply at a given point, produce necrosis and then degenerate and disappear.

Although the micro-organisms have the same morphology as the treponema pallidum, we are not thoroughly convinced that they are identical with that parasite.

A search through the literature revealed only two cases of syphilitic aortitis in which treponemata were found in the aortic wall.¹

TREATMENT OF OBSTRUCTION OF THE LACHRYMAL DUCT.*

BY JOHN C. BERRY, M.D.,

Ophthalmic and Aural Surgeon, Worcester City Hospital; former Surgeon Doshisha University Hospital, Kyoto, etc.

I. IN INFANTS.

DURING an animated discussion on dacryocystitis in the Ophthalmological Section of the British Medical Association at Toronto, in 1906,

¹ Reuter: *Zeitschr. für Hyg.*, 54, 1907, p. 49; Schmorl: *Münch. med. Wochenschr.*, 1907, p. 183.

* Read before the New England Ophthalmological Society, Boston, Jan. 12, 1909.

reference was made to "congenital dacryocystitis," when Dr. Knapp, Sr., made the statement that these cases (dacryocystitis in infants) never require treatment, — nature taking care of the difficulty in due time. I wonder if this is the experience of members of this society. In my own it has been largely so, and yet only recently I was obliged to interfere in the case of a child ten months old, suffering from obstruction of the duct and blennorrhea of the sac. I would remark in passing that this term, "congenital dacryocystitis," is unfortunate, for the sac is, I think, never inflamed at birth, this inflammation being always secondary to the morbid process set up within the canal itself. In this particular case (and I select a clinical case as the most convenient and concise way of presenting treatment for other similar cases), the blennorrhea began within a week after birth, and though mild disinfecting collyria were used, pressure made over the sac to force its contents on into the nose, and alkaline-antiseptic nasal sprays employed daily, yet the morbid process continued, and, indeed, grew worse, producing congestion and swelling of surrounding tissue. Fearing that a phlegmon would result if the case was allowed to pursue its course further, the child was put under an anesthetic, the canaliculus partially slit and a No. 3 olive-tipped probe passed on through the canal, a single passage being sufficient, as is usual, to completely relieve the difficulty. In these infantile cases the growth of micro-organisms is favored within the canal by the retained debris of fetal life therein, resulting in inflammation and softening of the lining membrane, thus making the passage of a small probe, especially one with a stiff shank, somewhat hazardous. I therefore hesitate to use a probe smaller than the No. 3, and an olive-tipped probe with a small flexible neck that will easily bend and allow the tip to follow the canal is favored. It should be remembered that the course of the canal in babyhood is quite irregular and radically different from that in the adult, the direction being backward, at times, to an angle of nearly forty-five degrees.

These cases of dacryocystitis in infants are almost always from obstruction, — the abnormal proliferation of the epithelium of the canal obstructing its lumen, — especially at the lower end of the duct, which is very small at birth and further obstructed by pressure from the lower turbinate. Such pressure and occlusion produces stagnation in the canal itself, with resulting decomposition of its contents. Under such a condition it requires but a few days' exposure of the eyes to atmospheric conditions for the lachrymal secretion with micro-organisms from the eyelids and surface of the eyeball to be forced on into the sac and the whole gelatinous mass to become the seat of decomposition, inflammation and the appearance of the characteristic discharge of dacryocystitis at the inner canthus of the eye. According to the report of the London Ophthalmic Hospital for last January, there have been found in these discharges