THE BRITISH JOURNAL
OF
DERMATOLOGY AND SYPHILIS
MAY, 1920.

STUDIES ON THE EFFECT OF LIGHT ON
VITILIGO.*

CARL WITH,

First Assistant of the Department of Skin-Diseases of the Finsen Institute.

*From the Dermatological Department of the Finsen Medical Institute.
(Director Dr. Axel Reyn.)*

WHILE working at light-baths in the Finsen Institute I clearly perceived that there was a problem of uncommon interest as to the relation of vitiligo to light. On December 4th, 1918, I demonstrated to the Danish Society of Dermatology† a patient suffering from a decided macular pigmentation produced by light-baths. To this demonstration I added some general views regarding the effect of light on vitiligo. It was my intention at that time to discuss the question on a broader basis, and I wanted especially to examine the pigmented spots by the dopa-reaction. However, Dr. Kissmeyer returned home from Bruno Bloch in Zürich with a supply of dopa,

* In part delivered as a lecture at the Fourth Scandinavian Dermatological Congress in Copenhagen in June, 1919.

† *Hospitalstidende*, Nr. 14, 1919, pp. 122-124.

and he took over this part of the examination, of which he gives an account in the current number of this Journal. This treatise must be interpreted as a provisional report, especially as the treatment of the literature is rather incomplete.

It is generally thought that pigmentation is a considerable protection against injurious effects of light, and that the human organism, so to speak, protects itself normally against these by means of pigmentation.

Before discussing the reaction of the non-pigmented parts of the skin to the light, I want to say a few words about the pigmented spots which are produced in vitiligo patches by the action of light.

I. NOTES ON PIGMENTED SPOTS IN VITILIGO.

The first person who succeeded in producing pigmentation in vitiligo was Montgomery (8), who treated patches of vitiligo in a Mexican about 19 years old; some of the patient's relatives were similarly affected. The patient was treated with Finsen light (London Hospital lamp), and was given five exposures of ten minutes on each patch. As a result "the skin was well reddened as in a sunburn. Some of the more recent patches about the mouth showed immediate improvement on the subsidence of the inflammation. Some of the older patches were much slower, both in reaction and improvement." In fact he stopped treatment long before restitution of pigment in most of the patches.

This case, the description of which is rather incomplete, is almost unique, and later authors—for example, Buschke and Ehrmann—have not succeeded in producing pigment in vitiligo by Finsen light. Buschke (2), in two patients suffering from vitiligo, succeeded in producing pigmented spots, similar to freckles, arranged around the follicles after exposure to the quartz-lamp. A single exposure at a distance of 10 cm. and of a duration of a few minutes produced in a fortnight or three weeks distinct pigmentation; the small spots of pigment extended little by little, so that the pigmentation became diffused. This pigmentation almost completely disappeared in about four months. Together with Mulzer (3) the above-mentioned author arrived at the same result.

Moser (9) mentions the result of treating a patient suffering from

vitiligo with "blue light." With seven exposures in a month the vitiligo was not only arrested in its development, but the hyperchromatic regions around the vitiligo patches became less marked, and the vitiligo became partly pigmented in the form of freckles. Ehrmann (4), at a somewhat later date, succeeded in producing pigmentation in patches of vitiligo, as well as in albinos, by the quartz lamp.

Stein (14) discussed the question later. By means of the Kromayer lamp he succeeded in producing pigmented spots like freckles by exposing the vitiligo patches for ten to twenty minutes at a distance of 0.5 cm.; the spots appeared in four or five weeks. In contrast to Buschke he was not able to produce pigmented spots by means of exposures of a few minutes. With cantharides it was not possible to get pigmentation, but by freezing for two or three minutes with carbonic acid snow pigmentation was obtained.

Bruno Bloch (1) has dealt with this question in one of his papers on the dopa-reaction, and has propounded a new and very interesting theory as to the nature of vitiligo. Dr. Kissmeyer intends to report on this subject.

At the Finsen Institute we have worked almost exclusively with the universal carbon arc bath for about two hours and a-half every day or every second day. This treatment was originally introduced by Finsen, and later, in the year 1913, used by Reyn and Ernst (10, 11), as well as by Strandberg (13), with excellent results when used in cases of lupus and surgical tuberculosis. Reyn, who has worked out this method of treatment more than anyone else, uses two carbon arc lamps, the current used being 20 ampères if a single patient is treated, and 75 if several patients are treated; the E.M.F. is 50-55 volts. For details I refer to Reyn's papers (10, 11). In this way we have treated seven patients (five men and two women), who, in addition to their vitiligo, were all suffering from tuberculosis of the skin, of the lymphatic glands, or of the bones (with the exception of one woman). The vitiligo was rather extensive in all these patients, of whom the eldest was 70 years and the youngest 41. In these patients the vitiligo was, as a first result of treatment, far more visible than before, because of the stronger pigmentation of the other part of the skin. In six cases we obtained similar results to Buschke and Stein, for pigmented spots like freckles little by little became apparent in the vitiligo patches

In a single woman, aged 56 years, who was shown before the Danish Dermatological Society on account of a widely-distributed vitiligo, and who from May 10th to September 29th had been treated with 44 quartz-light baths, the vitiligo became more obvious, and pigmented spots of a similar kind as in the above-mentioned patients were found. It deserves notice that the new pigmentation was less marked, and the small spots appeared to be more scattered in this woman than in the patients who had been treated with the carbon arc baths. In this patient, before the treatment, pigmented spots were observed in the edge of the vitiligo patches as well as outside, probably because the vitiligo was disappearing. When we remember that some of our vitiligo cases did not show the usual picture of a hyperchromatic marginal zone, and that the way in which a vitiligo reacts towards light is dependent, to a certain degree, upon its age and physiological cause, we are obliged to be very cautious in drawing conclusions from a single case; nevertheless we are bound to remember the slight therapeutic effect of the quartz-light baths compared to that of the carbon arc baths as shown by Reyn and Ernst.

In one of our cases of vitiligo, vitiligo upon the hands was present fairly distinctly at the first examination (July 31st, 1918). At the examination January 1st, 1919, the vitiligo was somewhat indistinct upon the hands; in some places, upon the body as well as upon the extremities, it was scarcely possible to make out what was vitiligo and what not (see fig. 1). On February 12th after a few baths the pigmentation was strong, in consequence of which the vitiligo was fairly visible. On February 24th after 8 light-baths the first small pigmented spots in the vitiligo were observed. This is the only patient in whom a latent vitiligo has become manifest and well circumscribed before our eyes.

It often happened, however, that a vitiligo remained unnoticed at the first examination before the light-bath was given, and also while it was concealed by a universal light erythema. Then a fortnight afterwards we are told that there is in the light-bath a man singularly spotted, and we are presented with a well-developed vitiligo, more marked than is usually seen in our country, because of the strong pigmentation of the skin. Unlike the above-mentioned authors, we have had no opportunity of observing vitiligo increase during the light-bath, and we might reasonably have expected that

FIG. 1.—L. February 1st, 1919. Before the beginning of the light-baths.

FIG. 2.—L. April 2nd, 1919. After 13 light-baths (last March 1st).

FIG. 3.—L. June 9th, 1919. Right arm after 3 light-baths further (April 7th, 11th and 21st).

TO ILLUSTRATE DR. CARL WITH'S STUDIES ON THE EFFECT OF LIGHT ON VITILIGO.

FIG. 4.—B. C. May 30th, 1918. After 21 light-baths

FIG. 5.—B. C. November 20th, 1918. After 74 (altogether 95) light-baths more.

FIG. 6.—B. C. May 30th, 1918.
After 21 light-baths.

FIG. 7.—B. C. November 20th, 1918.
After 74 (altogether 95) light-baths
more.

FIG. 8.—B. C. April 1st, 1919. After 53
(altogether 148) light-baths more.

FIG. 9.—B. C. July 17th, 1919. After 38
(altogether 186) light-baths more.

FIG. 10.—G. S. July 2nd, 1918. After altogether 38 light-baths from April 3rd to July 2nd, 1918.

FIG. 11.—G. S. February 24th, 1919. After 23 light-baths more from January 20th to February 24th, 1919.

FIG. 12.—G. S. June 30th, 1919. Without any light-baths more.

the strong stimulation of the pigmentary function from the light-bath might produce an over-stimulation, which might act prejudicially upon those areas of the skin which were inclined to vitiligo.

On the other hand, the observations made by Stein and Buschke were proved most completely; in all of our patients islets of pigmentation developed from the edge inwards, towards the centre. The pigmentation began as brown spots, quite indistinct, but later on they became more and more visible. The more regularly the patient gets his light-baths the more the vitiligo disappears. This fact is clearly seen in the patient whom I demonstrated before the Association for Dermatology (figs. 4-9). While at the first examination on May 30th, 1918, after 21 light-baths (from April 12th), he had small spots upon the abdomen, but had big, confluent vitiligo patches upon the thighs, he may now, after about 180 light-baths given every day or every second day for a whole year, be called white-spotted. Previously he was black-spotted upon the thighs and abdomen. The patient mentioned above, whose vitiligo was almost invisible during the first investigation, now shows a very strong pigmentation in the patches. This pigmentation began after 8 light-baths (the first on February 5th). Although the patient left off the light-baths about March 1st, the spots of pigmentation increased upon both arms. He had light-baths again April 7th, 11th, and 21st, having the left arm covered up. Seeing him again on June 6th the number of spots was only slightly increased on the covered left arm, whereas the islets of pigmentation on the right side had increased to such a degree that the white could only be seen very slightly (figs. 1-3). Because of the long period of rest, the surrounding pigmentation as well as the islets were only little discoloured. My impression from this patient is that the pigmentation begins around the hairs. That the pigmented spots were perifollicular was distinctly seen by an examination of the vitiligo patch around the umbilicus in a man B. C— (fig. 13) (March 15th, 1920).

In a woman (figs. 10-12) a distinct augmentation of the pigmented spots was observed when she returned to the Institute for examination, though she had not had light-baths for four months. In a third patient, a woman, L. J— (fig. 14), it was evident that the pigmented spots in the vitiligo had increased in size and that the islets of pigment had in some places become peninsulas, though she

had not had light-baths for about a month. In these patients we have proof that the pigmentation is still produced for some time after the direct pigment-inciting agent, light, has ceased to be active.

As you will realise from figs. 13-14 as well as from the other figures on a smaller scale, this process of pigmentation begins in the periphery and spreads little by little towards the centre; the islets gradually become peninsulas and lastly we get diffused pigmentation. The new spots are at the beginning rather pale, but become by degrees darker and darker.

The question of how long these pigmented spots last in the vitiligo patches we have not yet solved. Probably they remain there for several years. The spots which Buschke produced by a few irradiations with the quartz-lamp disappeared in the course of four months. In the woman (G. S—) mentioned above, the spots not only did not disappear after four months, but on the contrary increased in size, and in a man (B. C—), who had taken his last light-bath on July 17th, 1919, numerous spots were observed eight months later. In a man, aged 70 years, who had had several pigmented spots in the vitiligo patches after 75 light-baths from July 29th, 1918 to January 29th, 1919, some of the spots had disappeared seven months later, but many were very distinct; and in a gardener, aged 44 years, who had been treated altogether with 94 light-baths (53 from October 10th to December 16th, 1916, 34 from January 31st to March 13th, 1917, and 10 from March 16th to April 20th, 1918), typical pigmented spots were observed in the vitiligo fourteen months later (July 2nd, 1919). In the patient L— (figs. 1-3), who, as mentioned, had only had 16 baths from February 5th to April 21st, the number of freckles on the right arm was unaltered on February 3rd, 1920; on the left arm, which was covered during the three last baths, the spots had disappeared in the middle.

II. REMARKS ON VITILIGO AND LIGHT-PROTECTION.

If pigmentation were the only protection against the injurious influence of the light, we should expect, *à priori*, that wide-spread vitiligo spots would annoy the patients very much during the light-baths, and we should have to fear a constant recurrence of the

FIG. 13.—B. C. Umbilicus of fig. 9 to show the appearance of the pigmented spots (July 17th, 1919, after 186 light-baths).

FIG. 14.—L. J. Spots in the left groin about September 19th, 1919, after about 48 light-baths.

TO ILLUSTRATE DR. CARL WITH'S STUDIES ON THE EFFECT OF
LIGHT ON VITILIGO.

disagreeable erythema. What are the facts of the case? As far as all our seven patients are concerned we see that this effect has only partially taken place.

We obtained the usual reactions after the first light-baths, and in the case of some patients the vitiligo patches were somewhat more prominent; but all our patients maintained that the vitiligo spots did not annoy them very much after some time had passed, unless special circumstances asserted themselves. A patient who, because of a foot-splint, was obliged to remain in the same position, and who had light-baths upon a single vitiligo patch for the whole two and a-half hours, was somewhat annoyed by irritation and pricking in this patch. The acute erythema disappeared from the vitiliginous patches somewhat later than from other regions, and it annoyed the patient a little longer. However, this was not always the case. In some of the patients I have an impression that the scaling chronic light erythema is inclined to be localised in the vitiligo patches more than in other parts. When observing the patients during the light-baths one sees a well-developed erythema in the vitiligo patches; little by little it disappears; even twenty-four hours after one observes a feeble rosy colour of the patches.

I did a control experiment in a woman patient. From the beginning of light-baths (January 29th) the right groin was covered up, and after 14 light-baths the patient was exposed uncovered to the light. For two hours she turned the front to the light. In the course of twenty-four hours a very strong, light erythema developed, being more marked on vitiligo than upon the surrounding, normally pigmented parts. On the left side the patient felt no annoyance.

In a woman who had been treated with 41 quartz-light baths from June 14th, during which the right side of the abdomen was covered up, a similar effect of the light was observed; on September 24th the right side of the abdomen was uncovered during the bath and exposed to the light for an hour and a-half. On the right side, in due time, a very strong erythema appeared, which annoyed her very much, while the left side did not show any erythema.

From the previous exposition as well as from these experiments, which, strictly speaking, are superfluous, we must logically arrive at the following conclusion:

The human organism has the power of working out other means

of protection than that of pigmentation in order to stand, without annoyance, a light effect which, without gradual acclimatisation, would act in an injurious way.

This means of protection must certainly be found principally in the power of filling the superficial vessels as well as in the circulation in them. I do not, however, wish to enter upon this matter here. Haxthausen (6) has hinted at several of these mechanisms of protection in his thesis, *Skin-diseases Caused by the Light*, Copenhagen, 1919.

I shall not, however, omit to draw attention to some interesting investigations made by Aug. Krogh (7), and apparently not quite so well known to dermatology as they deserve. It has been demonstrated in his work, *Exploration of Supply of Oxygen to the Tissues and Regulation of the Capillary Circulation* (1918), that the degree of filling of the capillaries does not, as is usually supposed, depend only upon the contraction of arteries and veins. The capillaries can be contracted, and we find not only arterio-motor fibres but capillary motor as well, and even veno-motor fibres too. If the arteries, veins and capillaries are dilated we get hyperæmia; if the arteries or veins are contracted while the capillaries are dilated we get a local hyperæmia of a bluish colour (this is observed in chilblain circulation, and is very prominent in the flat vascular nævi).

Reyn (11) has propounded this theory: that the effect of the quartz-light as opposed to that of the carbon light is that the former produces stasis and the latter hyperæmia. He is probably right, as the slightly penetrating rays from the quartz-lamp can produce dilation of the capillaries only and not of the arteries. If the carbon arc is used there will be a dilation of the arteries as well as of the capillaries, and thereby strong hyperæmia, deep pigmentation and better therapeutic results. It seems to me very interesting that Reyn has discerned correctly, and has probably found an acceptable theory for the better therapeutic results, although no physiological basis existed for it at that time. I have no idea of the mechanics of vitiligo. I have taken this opportunity of attracting attention to these modern physiological investigations, which seem to be only slightly known, although they present new points of view for clearing up many questions of the greatest importance to our knowledge.

If I am right, that vitiligo patches and other unpigmented parts

possess other mechanisms than pigmentation for protection against light, then the same question must, as a matter of course, be raised as to the normal skin, and thus we raise the question of the importance of pigmentation.

From his experiment of exposing his blackened arm to the sun, Finsen (5) concluded that light produces pigmentation and that the latter is a protection against the chemical rays of the sun. The last part of the conclusion is hardly justifiable. It ought to have run: the light is the condition of pigmentation, and at the same time the skin has acquired a kind of "immunity" from the light. Many things go to show that the protection is given by the pigmentation, but about this we will speak on some future occasion.

As we know, pigmentation is partly resultant on mechanical, thermal and light influences upon human skin; it appears, too, in certain illnesses as well as being caused by certain poisons. The skin is often strongly pigmented on spots where it is exposed to maceration (for instance, at the anus and on the nipples). As far as the genitalia and the nipples are concerned, the internal secretion is certainly of great importance. It is possible that a protection is acquired by this, but till now it has not been proved. Solger (12) maintains this theory, but has not been able to prove it. There is, perhaps—as Solger has suggested and as Dr. K. A. Heiberg has pointed out to me—a certain connection between the pigmentation and the quick renewal of the skin. A decided pigmentation often takes place as a result of Erythema caloricum; it is, however, to be doubted whether this pigmentation gives a greater power of resistance against the effects of heat subsequently. After a strong Röntgen erythema there appears, as we know, a strong pigmentation, which, however, gives no protection against renewed Röntgen treatment.

As far as this question is concerned we can only consider this a strong probability. But we dare not beforehand exclude the possibility of pigmentation (in negroes, for instance) actually being a thing quite secondary, and without any connection with the increased resistance of the skin to the sun.

If we look at this question from a phylogenetic point of view it must be admitted that the organism may perhaps, little by little, have acquired the power of reacting by pigmentation to protect itself

against the injurious effect of the chemical rays of the sun, which is of greater importance than any other injurious effect. As something quite secondary the organism reacts in the same way towards chemical and thermal influences, though the pigmentation is no protection against a renewed action of these agents.

However, there exist so many observations (for instance, light erythema in partially coloured animals and men) that it seems quite natural to consider pigmentation as an important protection, against the sun in particular, because it has, as Unna (15) points out, the colour of yellow-brown, which absorbs the greatest number of violet and ultra-violet rays; and, moreover, the light-rays, by exciting pigmentation, prevent their own entrance and thus provide protection to the skin (Finsen (5)). As mentioned before, pigmentation is surely not the only protective remedy.

I just want to mention that an apparently strong and early pigmentation, congenital as well as acquired, gives a smaller power of resistance against the light than a less strong one which has been acquired later. In a negro, for instance, a very annoying erythema was produced by giving him a continuous light-bath of two and a-half hours at once.

SUMMARY.

(1) During carbon arc light-bath vitiligo becomes visible in regions where formerly it was difficult to perceive; we have, on the other hand, not succeeded in producing new spots of vitiligo.

(2) The carbon arc light produces upon the vitiligo patches a macular pigmentation more or less decided; it spreads from the periphery to the centre, and may last at least fourteen months.

(3) In three cases this pigmentary process continued after the light-bath had ceased.

(4) It is true that the vitiligo patches react with a stronger erythema than the pigmented skin, but little by little the patients are able to stand a quantity of light, to which they formerly reacted with an erythema, not only upon the pigmented but also upon the vitiliginous regions of the skin. Consequently the organism is capable of protecting itself against the light in another way than by pigmentation.

(5) The final proof that the rôle of the pigmentation is to form a

protection against light has not yet been forthcoming. So many observations, however, go to show it that it must be considered the most probable conclusion.

I tender the chief of the clinic, Dr. Reyn, my best thanks for his interest and kind assistance.

LITERATURE.

(1) BLOCH, BRUNO.—“Zur Pathogenese des Vitiligo,” *Archiv f. Derm. u. Syph.*, 1917, cxxiv, pp. 201–32.

(2) BUSCHKE, A.—“Notiz zur Behandlung des Vitiligo mit Licht,” *Med. Klinik*, 1907, Nr. 33, pp. 983–84.

(3) BUSCHKE, A., and MULZER.—“Weitere Beobachtungen über Lichtpigment,” *Berl. klin. Wochenschr.*, 1907, Nr. 49, pp. 1575–76.

(4) EHRMANN, S.—“Pigmentanomalien,” Mracek's *Handbuch der Hautkrankheiten*, ii, p. 750. “Ueber physiologische und therapeutische Lichtwirkung,” *Verh. d. Deutsch. dermat. ges. Arch. Derm. Syph.*, 1908, xci.

(5) FINSSEN, NIELS R.—“Om Lysets Indvirkning paa Huden” (“On the Effect of the Light upon the Skin”), *Hospitalstidende*, 1893, p. 722.

(6) HAXTHAUSEN, HOLGER.—“Sygdomme fremkaldt af Lyset” (“Diseases Produced by the Light”), *Disputats*, 1919.

(7) KROGH, AUG.—“Undersøgelser over Vævenes Forsyning med Ilt og Kapillærkredsløbets Regulering” (“Studies on the Supply of the Tissues with Oxygen and on the Regulation of the Circulation in the Capillaries”), *Det kgl. Vidensk. Selsk. Biol. Medd.*, 1918, i, 6, pp. 1–29.

(8) MONTGOMERY, DOUGLAS.—“Vitiligo Treated with Finsen Light,” *Journ. Cut. and Urin. Dis.*, 1904, xxii, pp. 17–19.

(9) MOSER.—“Zur Behandlung des Vitiligo mit Lichtbestrahlungen,” *Med. Klinik*, 1907, Nr. 45, p. 1363.

(10) REYN and ERNST.—“Om Anvendelse af kunstige Lysbade ved Lupus vulg. og kirurgisk Tuberkulose” (“On the Use of the Artificial Light-baths in Cases of Lupus and Surgical Tuberculosis”), *Hospitalstidende*, 1914, Nr. 43–44, pp. 1297–1311 and 1321–30.

(11) REYN and ERNST.—“Om Anvendelse af kunstige Lysbade ved Lupus vulg. og kirurgisk Tuberkulose” (“Further Studies on the Use of Light-baths in Cases of Lupus and Surgical Tuberculosis”), *ibid.*, 1917, Nr. 19–20, pp. 448–63 and 471–91.

(12) SOLGER, F. B.—“Die Bedeutung des Pigments für die hellfarbigen Menschenrassen,” *Derm. Zeits.*, 1905, x, pp. 516–27.

(13) STRANDBERG, OVE.—“Det universelle Lysbads Anvendelse i Rhinologien” (“Studies on the Value of the Universal Light-baths in Rhinology”), *Hospitalstidende*, 1918, Nr. 7, pp. 193–203.

(14) STEIN, ROBERT.—“Ueber experimentell erzeugtes Pigment in Vitiligo,” *Archiv f. Derm. u. Syph.*, 1909, xcvi, pp. 163–78.

(15) UNNA, P. G.—“Ueber das Pigment der menschlichen Haut,” *Monatshefte f. pr. Derm.*, 1885, p. 285.

This document is a scanned copy of a printed document. No warranty is given about the accuracy of the copy. Users should refer to the original published version of the material.