[Popular Science Review—Abstract.]

HOW MUSHROOMS ARE REPRODUCED.

(*Agaricus lacrymabundus*. Fr.)By WORTHINGTON G. SMITH, F.L.S., M.A.I.,
F.R.HIST.S. IRELAND, ETC.

THE nature of the gills in various members of the mushroom tribe differs very much, and this difference is especially remarkable in the absence or presence of the trama, and in the number, form, and size of the cystidia. In the genus *Coprinus* there is no trama to the gills. The trama in true *Agaricus* is formed by cells, which are of a different nature from the other simple cells which go to form the gills, and this trama forms the intermediate substance between the hymenial surfaces.

Nothing can be more different than the interior structure of the gills in the genus *Coprinus*, and in *Agaricus lacrymabundus* (Fr.). In Fig. 1 is shown, enlarged 150 diameters, a vertical transverse section down a gill of *C. radiatus* (Fr.), which species is (though minute) quite typical of the entire genus. The trama with its large cells would, if present, be at A; but a glance at the figure will show that the individual cells throughout this section are almost identical in size. If reference is now made to the similar section of a gill belonging to *A. lacrymabundus*, Fr. (and shown on Pl. *), the nature of the cells at A, which go to form the trama, will be quite apparent. Wide differences of cell-form like this are common in the mushroom tribe, but the difference of form as found amongst the cystidia is still more striking. In *Coprinus abramentarius* (Fr.) the cystidia are indeed so large that each individual cystidium would hold within itself more than 300 of the ordinary cells which go to build up the pileus. These facts regarding the cap and the hymenial surface (to say nothing of the gills often completely separating from the horny hymenophorum, as in *Paxillus*), and many other known facts in regard to the nature of the stem and its outer surface, go far to disprove the hypothesis that the higher fungi found under the Agaricini may be generalized as mere concreted masses of moulds. No bringing together into a mass of a forest of Penicillium could even make an *Agaricus*, and to me there is nothing in common between the conidio-spores of Penicillium and the basidio-spores of the Agaricini.

The various bodies which, as I believe, belong to the reproductive process in the mushroom tribe are seen in Figs. 1 and 2 and Pl. *. The peculiar cells of the trama are shown at A, the simple cells which form the external and hymenial surface at B, and the privileged cells at C, D. C shows the cystidia and D the basidia. Each basidium in the Agaricini bears four minute spicules, which carry the same number of spores. The difference in size between the basidia and cystidia is often immense, and in *Agaricus lacrymabundus* the perfect forms of the two organs are very different from each other; they, however, so approach each other in this species that every intermediate form between a cystidium and a basidium may be observed. In *Coprinus radiatus*, as I have shown elsewhere, the cystidia have their contents differentiated, and at once produce spermatozooids; or they germinate, and the spermatozooids are produced from a differentiation which then takes place at the end of the thread; these spermatozooids attach themselves to and ultimately pierce the spores, and their piercing causes the discharge of a single cell from the pierced spore, which cell belongs to the pileus of the new plant. As I interpret my observations, this process is carried on upon the gills themselves, as shown in Fig. 1, and at the time of the perfect maturity of the fungus, at least in *Coprinus*, in the sub-genus *Volvari*, in the ordinary mushroom, and in such plants as have fallen under my notice. The basidia with their spicules and spores, the cystidia germinating and bearing spermatozooids, and the first cells of a new plant (E) are all shown in position in Fig. 1.

Agaricus lacrymabundus (Fr.) is a very different plant from *Coprinus radiatus*. It is one of our commonest Agarics of the autumn, and is usually found in damp pastures and about stumps. It bears a considerable resemblance to the common mushroom, and is without doubt often gathered for the table in mistake for that species, but whether with any ill effects or not I am unable to say. The most striking character of this Agaric resides in its gills, which are always furnished with a white edge, which drips with semi-milky tear-like drops.

Part of a section of a specimen of *Agaricus lacrymabundus* is shown at E, Pl. * with the characteristic drops *in situ* on the gills and upper portion of stem, F G. The drops are shown towards the edge of a gill enlarged 100 diameters at H, above which may be seen several drops of a smaller size, which may or may not at length coalesce with the lower drop. These drops invariably dry up on the edges and surface of the gills. As far as I know, milky tears peculiar to *Agaricus lacrymabundus* have never been minutely described, neither has an account of any microscopic examination of them been hitherto published.

These drops are distilled direct from the cystidia. When the fungus reaches maturity the cystidia protrude boldly from the surface of the hymenium, as seen at C; and the apex of the cystidium opens exactly as does the ascus in *Peziza*. At first the cystidium is filled with a thin fluid in which no granules can be detected; but at length a large vacuole is formed at the base of the cystidium, and the contents are forced towards the apex: here the fluid becomes partly differentiated into granules, and at length the cystidium opens at the top and discharges its contents. The protruded drop now swarms with moving atoms of a regular size, which I refer to spermatozooids. They are in every way identical with the spermatozooids of *Coprinus radiatus* (Fr.).

COPRINUS COMATUS. Fr.
Vertical section and surface (F) of Gill enlarged 150 dia.

In my recent paper on the reproduction of *Coprinus radiatus* I adverted to the fact of the cystidia falling bodily away from the hymenium, and exactly the same phenomena holds good in *Agaricus lacrymabundus*; for if a drop of the liquid from the gills is examined under the microscope it will be seen to swarm with free cystidia which have dropped into the fluid from the gills; this liquid not only abounds with spermatozoid and cystidia, but it also swarms with spores which have fallen away from the basidia, as seen in Pl. *

If the spores which have fallen into the drop are examined, they will be found to be very different in aspect from the spores as they are seen upon the basidia; for whilst the spores in the latter position are perfectly plain in outline, as seen at J, enlarged 2000 diameters, the spores within the drop

are studded all over the surface with the spermatozooids discharged from the cystidia. Not only are the spores studded as shown at K, but they are pierced through the external coat by a fine thread protruded from the spermatozoid. The effect of this is that a single free cell is soon after discharged from the spore, and this floats about in the drop, leaving the exospore quite empty. This single transparent cell, which is three or four times the size of the original spore, and is the first cell of the pileus of a new plant, is in its turn commonly attacked and pierced by the spermatozooids. Sometimes a spore is surrounded and pierced by a large number of spermatozooids, whilst at other times it receives the attack of one body only; this is, however, sufficient, for I have frequently seen the first cells discharged from the orifice made by a single spermatozoid, and the latter body again pushed out by the cell. The spermatozooids which have a single turn of a spiral revolve rapidly, and when they come in contact with a spore, not unfrequently creep with an amoeba-like movement over the surface, before they discharge their contents into the protoplasm of the spore. These bodies, as seen in *Coprinus radiatus*, are shown in the accompanying figure, enlarged to 1000 and 3000 diameters, and they are precisely the same in *Agaricus lacrymabundus*. When the spermatozooids of the latter species emerge they are dark brown in color, and here they resemble the spores; the at first colorless contents of basidia and cystidia alike become at first differentiated and then oxidized on exposure to the air. In a minute species like *Coprinus comatus*, where the growth of the plant is very rapid, and hundreds of generations are produced in one season, new plants are rapidly reproduced from the spores, and the growth can be watched; but in *Agaricus lacrymabundus* the case is very different, for it is probable that it would take many months (if not a whole year) to grow a perfect fungus belonging to this species from the spores. It must also be remembered that although this Agaric produces many millions of spores, yet the plant does not increase in actual numbers; for if all the spores produced plants, then in one year the world would be covered with this species only. For one spore which grows, hundreds of millions must be failures.

In a former paragraph reference was made to bodies intermediate in form between cystidia and basidia, and bodies of this nature are common on the upper part of the stem in *Agaricus lacrymabundus*; they are shown on the natural size section at G. The bodies in this position secrete and distil a semi-milky fluid, which they excrete in drops, which drops are invariably dotted all over the upper portion of the stem. It must not be thought that this distillation is peculiar to the species under notice, for it is a character (though till now unpublished) of whole sections of Agarics. For instance, certain groups coming under *Tricholoma* are described by Fries as "lamellis decolorantibus, rufomaculatis l. cinereiscentibus." Now these brown spots on the gills, when examined under the microscope, are seen to be produced by a liquid which changes color, and which is distilled from the cystidia; and it is quite possible that the same phenomenon holds good in many other fungi, as in the tears of *Merulius lacrymans*, and in the drops found upon such species as *Polyporus dryadeus*, *P. hispidus*, *P. cuticularis*, and many others. It seems probable that in some species this liquid, which by a differentiation produces spermatozooids, also sends the spores into a temporary resting condition, and that the spores rest before germination, just in the same way as many seeds rest. This would explain the great difficulty of getting some fungus-spores to germinate. Under any circumstances diverse fungus spores resemble seeds in the fact of some germinating at once, whilst others will not germinate till long periods of time have elapsed.

On examining the semi-milky drops as seen on the stem, I have been unable to observe any production of spermatozooids, but these intermediate cystidium-like bodies are expelled from the stem and fall into the drops of moisture distilled. This is a state of things to be expected, for as these bodies belong to neither basidia nor cystidia, they are as a consequence quite unable to produce spermatozooids as spicules and spores. In Agarics and Boleti the stem may be considered as a mere barren hymenium; when the stem is striate (as it commonly is) these striæ represent the absent gills; in species with decurrent gills, the fruiting surface (as in *Agaricus prunulus*) sometimes quite reaches the ground. When the stem in Boletus is reticulated, the reticulations represent the absent tubes, so that it should be a matter of no surprise to find organs pertaining to the hymenium upon the stem. The external surface of the pileus, for the same reason, often bears basidia and spores, and the stuffed stem and cartilaginous bark answers to the trama and hymenial surface. An Agaric or Boletus with the pileus and gills arrested would answer to a *Clavaria* bearing fruit all over the club, and abnormal Agarics are sometimes found in this condition, whilst the simplest form of *Hymenomyces* is where a merely filmy hymenium is developed, as in *Hymenula*.

When it is remembered how innumerable are the myriads of spores and spermatozooids set free every autumn, and how probable it is that hybrids of every degree are produced from these bodies, the diverse and almost countless forms seen in the mushroom tribe quite cease to be a matter of wonder. Without doubt the spermatozooids of some species commonly pierce the spores of an ally, and so produce plants intermediate between one species and another; such forms are an every-day experience with fungologists and lichenologists, and the more one knows of species as species, the greater are the difficulties to be surmounted in correct naming. Fries himself says, under *Mycena*, that he has only given the best marked species, and that he knows in his mind many other apparently distinct

HOW MUSHROOMS ARE REPRODUCED.

forms and species which he has never published. Our cryptogamic floras require cutting down to at least one fourth of their present dimensions by some competent botanist, who is well acquainted with species. In the majority of instances the species are so ill-defined that it is the easiest possible matter to gather plants which it is simply impossible to refer to any described fungus, simply because the specimens gathered belong to none.

The spore, as found amongst Agarics, I consider female in the sense of its pertaining to the female in the same way as an unimpregnated ovum may be considered a female organ whilst still attached to the mother, and the spermatozoids male, as pertaining to the male in the same way as spermatozoids are always the direct offspring of the male (as ova are of the female). When the spore is pierced by the spermatozoid, the former is capable on germination of producing either or both sexes. If these views are correct, a term is required for the unpierced spore synonymous with the ovule in flowering plants, as distinguished from the seed; and the old term sporule might well be used for indicating the unpierced spore. The spores in *Agarics* must be considered very different in nature from gemmæ, or buds, and they are totally different from the conidia or conidio-spores of *Penicillius* and the acro-spores of *Mucor*. The latter may reasonably be compared with the bulbils of *Dentaria*, and some *Liliaceæ*; but the nature of the basidio-spores (and of the hymenium which carries them) points to a very different conclusion. It is a very common thing for the mycelial threads of all sorts of fungi to break up into bead-like conidia, and as these conidia or secondary spores have a certain reproductive power, there is here certainly a slight analogy with gemmæ. The spores and spermatozoids in the *Agaricini* appear to me to have a strong analogy with the oosphere and spermatozoids of *Fucus vesiculosus*, the cystidia having an analogy with the antheridium in the same *Alga*. I have expressed an opinion elsewhere that Van Tieghem's idea of male and female spores in the *Agaricini* is altogether untenable; such a thing as a male ovum or spore is as unreasonable as a female spermatozoid or female pollen-grain. Seeds of all sorts are capable, on germination, of producing either or both sexes, though it is common enough to see one sex exalted at the expense of another. Even in the highest mammals the males have a trace of the female in the subordinate mammals and other characters, and similar characters which show a trace of the male are found in most female animals.

In the vegetable kingdom nothing is more common than to find so-called male or female plants changing their characters. Males will, under altered conditions, carry female organs, and females will produce anthers and pollen, which conclusively shows that not only are ovules, spores, seeds, and eggs not in themselves male or female, but the produce itself of these eggs is inherently hermaphrodite. It is convenient to name many animals and plants "male" and "female," because they are almost but not entirely so.

Several papers have been published of late on the reproductive process in the *Basidiomycetes*, and the writers of these papers have noted (with me) the diverse mycelia seen amongst the germinating spores; but I am convinced that the threads which produced antherozoids, as seen by Van Tieghem, came from the cystidia and not the spores. When a mass of spores and threads are seen in a solution of horse-dung, nothing is more difficult than to decide for certain whether the threads really come from the spores or not, and the spores of *Coprinus radiatus* could not possibly have been "sown" on any decoction without cystidia likewise falling upon the liquid. The "rod-like" bodies described by Van Tieghem read remarkably like an illustration of the well-known fact of the threads of many fungi breaking up into Bacteria at the tips. In the many *Agarics* I have examined, it invariably happens with me that after a spore is pierced it discharges a single cell, which develops directly into a new individual, exactly as is seen in *Chara*, but an unpierced spore may produce a thread of indefinite length; now if this thread is attacked by the spermatozoids, it will in its turn produce this single primordial cell of a new fungus, or if the mere undifferentiated liquid contents of the cystidium should pass over the thread from an unpierced spore, it there gives rise (at the point of contact) to the primordial cells.

The persistence of form in spores, and especially the pierced spores, under extraordinary conditions, is something remarkable; for instance, repeated violent boiling has no effect on the form and color of the spore, and in the cases where the spermatozoids are attached no amount of boiling disengages them; if any thing, the boiling seems to make the piercing more distinct to the eye. I do not find conidia or gemmæ resist boiling in this manner, and certainly no bulbil or bud from a flowering plant would maintain its form under these conditions. As for the unpierced spores, I believe their life to be of the very shortest duration (not twenty-four hours), but after piercing the life remains, just as we find life slight in an ovule but enduring in a seed.

As dung swarms with fungi and infusoria of all sorts, the greatest care is necessary in experiments, or Bacteria will be mistaken for Spermata, and Sphærobacteria for Spermatozoids. To clear up some little of the confusion which might arise as to all these bodies, the infusoria are engraved in Fig. 2 to the same scale as the spores and bodies referred by me to spermatozoids; A shows the globular spermatozoids as they are at first produced within the cystidium, or within the end of a thread more or less long, protruded from a cystidium; these bodies are at first motionless, but after being kept in liquid for a few hours they begin to slowly revolve; this movement keeps on for several days, and is all the time accelerated, whilst the body (formerly spherical) now becomes slightly elongated, and a single turn of a spiral is seen in the larger figures shown at the bottom of the plate; the dotted lines indicate the whirling motion of the spermatozoids. For comparison with this, various tailed and tailless monads are engraved at D, which figures may be compared with the spermatozoids to the same scale on the upper part of the plate. These latter bodies have been long known, and Mr. Berkeley, in the "Micrographic Dictionary" (last edition), p. 20, says: "The bodies called cystidia or pollinaria are globular or oval cells, found associated with the basidia, containing granular matter, exhibiting molecular motion when discharged." In common with many other botanists, I am inclined to think this movement to be other than molecular, as the eddy clearly seen round the revolving bodies indicates the presence of cilia. It is impossible to believe that life can spring from no-life, the mere molecular movement of lifeless particles must

be of a totally different nature from the life-movements of spermatozoids and infusoria. The revolving bodies might be referred to Micrococcus, one of the Sphærobacteria, but it must be remembered that they are the differentiated contents of the cystidia. It may be answered that if the mycelial threads of fungi will break up into true Bacteria, why may not the cell contents reappear as one of the Sphærobacteria? The question then arises, What is the nature of the obscure bodies referred to Bacteria? Are Dr. Eidam's spermata no other than Bacteria? they are sufficiently like Bacteria. Are certain so-called Bacteria and antherozoids of some cryptograms one and the same? When once produced they are both very persistent, and they are both produced under somewhat similar conditions. The spermatozoids of *Agarics* only appear as the fungus has just passed maturity, and as decomposition is setting in; and as soon as the spermatozoids are set free, then the material from which they spring has vanished.

It is just worthy of note that certain spermatozoids and bacteria resemble each other in size and external form, that they are alike in being furnished with flagella or cilia, that they are produced at a time when decomposition is just setting in, and that they are seen in semi-milky fluids. The spherical bodies when they become attached to the spores, or even threads, rupture at one end and discharge a fine thread, as at E, where they are shown germinating in a free state, which is no uncommon occurrence. Vibriones are shown at F, Fig. 2; these bodies, with the bacteria, show at G and H, are well known to all microscopists, and it has appeared to me that they are mere differentiated states of the old cells of the fungus which have broken up and (as a collection of cells) totally disappeared. The dotted lines indicate the movements of the bodies, whether straight, zigzag, or revolving like a wheel.

Whether the fertilized spores are able to withstand great vicissitudes of temperature and still not lose their vitality is unknown, but if other living atoms are able to live in boiling fluids, or be frozen and still live, it should be a matter of no surprise to find fungus spores passing unscathed through similar ordeals. It is reasonable enough to imagine life to be

FIG. 2. COPRINUS RADIATUS. Fr. Spores pierced by Spermatozoids: unpierced Spore germinating at both ends; infant plant and infusoria enlarged 1000 diameters; Spermatozoids at bottom of plate further enlarged 3000 diameters.

maintained under extraordinary conditions, but to me most unreasonable to imagine life to spring from that which is without life.

As the cells of decomposing fungi disappear, various infusoria at the same time swarm into being and take the place of the collapsed cells. On violently boiling these infusoria in test-tubes along with fragments of putrid fungi, for five minutes, and sealing at the highest point of ebullition, I find the infusoria, after one or two or three months, to be still alive. During this time putrefaction is arrested in the test-tubes, the decayed fragments of fungi remain the same, and the infusoria remain inert. On opening the flasks the infusoria are at first motionless, and life is apparently at a low ebb, but the individual infusoria, as watched under the microscope, rapidly regain their accustomed activity, and in a few hours are as full of life as if they had never been boiled. The decomposing fragments of fungi now rapidly give the fluid an offensive odor.

Whilst able to resist the heat of boiling-point, these same infusoria are equally well able to resist cold, for during the frosts of the past month I froze decoctions of fungi containing infusoria into solid blocks of ice for a whole week. On thawing after this time the bacteria, vibriones, monads, etc. were as full of life as before freezing.

EXPLANATION OF PLATE.* (preceding page.)

FIG. 1. Vertical section through gills and stem of *Agaricus (Hypholoma) lacrymabundus* (Fr.). E, Veil, F, drops on gill. G, drops on upper part of stem, natural size.

FIG. 2. Vertical transverse section through edge of gill, with drops *in situ*. Enlarged 100 diameters. A, Trama. B, Cells forming Hymenium. C, Cystidium. D, Basidium. E, lowermost drop, containing cystidia and spores.

FIG. 3. Cells of Hymenium. C, Cystidia, containing sper-

matozoids. D, Basidia, with spores attached, and tringular flat crystals from the juices of the plant. Enlarged 500 diameters.

FIG. 4. Spore unpierced by spermatozoids, enlarged 3000 diameters.

FIG. 5. Spore pierced by spermatozoids, from a drop which had dried on the surface of the Hymenium. Enlarged 3000 diameters.

[Journal of the Society of Arts.]

IMPORTATION OF FRESH MEAT FROM AMERICA IN REFRIGERATED CHAMBERS.

ATTENTION having very recently been called in these columns to the various methods projected and experimented on for bringing meat to this country by the aid of refrigeration, it is not necessary to recapitulate them. Though some have been partially successful when tried on a small scale, it is unhappily a fact that, till within the last few months, or it might be said weeks, no method has been practically adopted in the way of a commercial enterprise likely to appreciably augment our meat supply and affect our home market. The cost of producing and maintaining by ice itself, or by some chemical process, a sufficiently low temperature for the transportation of fresh meat to this country, has been the main obstacle in the way of a realization of the much desired end; and it will, therefore, be interesting to put on record some account of a recent attempt to supply our Metropolitan meat-market with fresh meat from America, by aid of the means alluded to.

It was in October last that the experiment was first made, and this having proved successful, it has been followed up with a succession of importations since Christmas, the total weight imported having reached as much as 400 tons, and it is stated that the consignment which arrived last week exceeded all previous ones. The meat hitherto received, consisting chiefly of beef, has been shipped in steamers at New-York, landed at Liverpool, and forwarded to the Metropolitan Meat-market by train, where it has been openly sold, and in as good condition as that in which it left New-York. Of course there is nothing new in meat being kept in a temporary state of preservation by means of refrigeration. At the International Exhibition at South Kensington, in 1873, when a variety of refrigerators or ice-safes were exhibited in the Food Section, Mr. Kent, of High Holborn, exhibited a ventilated ice-safe constructed on this principle. A large number of the leading butchers in London and the provinces have had chambers constructed on this principle in their establishments, and found them to answer admirably; at the Alexandra Palace the refreshment contractors keep tons of miscellaneous provisions in such a chamber during the hottest weather; several first-class ocean steamers, the *Walmer Castle* and *Windsor Castle* for instance, have been furnished with them, so that passengers and crew can depend on a supply of fresh meat for more than two months; while the principle has been adopted at the House of Commons, and the draught of air over tons of ice keeps the house at a comfortably low temperature during the dog-days. But this "draught" principle of refrigeration has not been yet applied in reference to the transport of fresh meat from foreign countries, unless, indeed, the patented system by which the recent importations of meat have been effected is more or less an adaptation of it, thereby affording another instance of an American borrowing English ideas and getting the credit of originality.

The result of the experimental importations has been reported as satisfactory as far as the condition of the meat is concerned. The quarters of beef, being tightly stitched up in canvas and suspended in a chamber on board ship, are constantly surrounded by a temperature of about 37 degrees, this atmosphere being kept in constant motion by the action of a fan worked over the ice-tanks by steam. Hence the meat is kept not only cool but dry, in fact so much drier than ordinary meat that it is not in the least affected by its transit from Liverpool to London. But, according to the old adage, the best proof of the pudding is in the eating. The imported American beef has been entrusted to several salesmen in the Metropolitan Market by Mr. J. D. Link, also of the Metropolitan Market, and sole agent of the American consignors, and the salesmen have made no secret of the matter at all, but sold it for what it was, and at a price but a few pence per stone below that realized by good Scotch and English meat. Those who have tasted it in a critical spirit, and among them may be mentioned the Lord Mayor and other civic authorities, have declared it to be in no way inferior to sound Norfolk beef or even "prime Scot."

What will be the future of this experiment, the extent of its development, and its effect on our markets, are interesting questions. It is said that it has already caused a fall in Scotch meat of 1d. per lb., though the foreign meat imported is, comparatively speaking, inappreciable in quantity. But the consignors, it is stated, are ready to send over 500 tons per week "if the market will bear it." If this were carried out, prices "all round" must be affected, but with them the price of the imported meat itself. And then what will happen? The question is asked, because the price of meat in New-York does not differ substantially from the price here. Perhaps the success of the experiment will lead to its being tried in reference to meat obtained in some cheap locality, and shipped at other ports than New-York, and thus the cost of present carriage, or of driving the cattle from Western districts to New-York, would be saved. But, then, arises the question of a longer voyage, increased consumption of ice, and so increased cost in these particulars. Again, the recent experiments have been made mainly in the winter months. How will the system work in the summer months? There will be a greater consumption of ice, of course, and not so much to sell of that unconsumed on the voyage as has been sold in Liverpool during the winter months. And further, the transport of the meat from Liverpool to London by railway will be a hazardous thing in the summer, unless refrigerating and ventilated wagons are used—and this would, of course, add to the expenses.

It is probable that the Food Committee of the Society will shortly make a trial of some of the American meat, and report upon the result.

THE Academy of Medicine and Surgery, St. Petersburg, is attended by 171 women and girls, 103 of whom are of titled birth. They mix harmoniously with the other sex in the dissecting-room.