being placarded. This results in an apparent mortality rate of from 25 to 30 per cent., a figure very little, if any, less than that which existed before the introduction of antitoxin.

Second, the fact that physicians who only occasionally use antitoxin administer it in small doses or too late in the course of the disease.

The writer gives the notes of cases occurring in his own practice before the introduction of antitoxin, showing the uncertainty of treatment at that time. He compares this with the specific action of antitoxin and the consequent reduction of the mortality rate which he has observed in his own experience under its use.

In conclusion he gives the following rules, which should be observed in every case:

A culture should always be made in throat cases.

Give antitoxin at the time of making culture.

Give 2000 units, and if there is no improvement in six hours, repeat the dose. Every twelve hours thereafter administer 2000 units until improvement begins.

The Arthritides of Scarlatina.—SZONTAGH (*Jahrbuch f. Kinderheilkunde*, 1902, Band v., S. 702) records fifteen observations of scarlatinal arthritis. Except in two cases, joint manifestations did not occur before the fourth week, several times five, six, or even seven weeks intervening after the onset of the scarlatinal attack.

In six cases a single articulation was affected; in four, two joints; in four, three; and once, four. The articulations of the left side were involved twice as frequently as those of the right side. The joints affected were the ankle in seven, the wrist in six, the carpo-metacarpal joint of the index finger in six, the knee in five, the shoulder in two, the elbow in two, the hip in one, and once the sternoclavicular articulation.

In all these cases there was at the same time a nephritis, and in the majority this was quite grave. In thirteen of the cases the nephritis had preceded the joint symptoms; in one case renal and articular manifestations appeared simultaneously; and in one arthritis developed first. In the author's opinion there existed a sort of connection between the nephritis and the arthritis, in several cases the appearance of uræmia apparently being prevented by the development of joint symptoms.

The author considers the joint involvement not as a secondary infection, but as a localization of the scarlatinal poison. True scarlatinal rheumatism, therefore, is not attended with suppuration. If suppuration occur it should be regarded as an expression of a general pyæmia, and not as a specific manifestation of scarlet fever.

Cream for the Home Modification of Milk.—CHARLES W. TOWNSEND (*Boston Medical and Surgical Journal*, April 16, 1903, p. 414) discusses the accuracy of results obtained in securing creams for modification by the several methods in vogue. These are: (1) Cream by the centrifugal process; (2) gravity cream or top-milk obtained by siphonage; (3) gravity cream obtained by dipping off the top; and (4) gravity cream obtained by pouring off the top.