It is to me of profound significance that the one who hopes most in all these cases, and who is most loth to give up a patient, is Dr. Trudeau himself. He, more than any man in this country, has studied the possibilities of treatment and of cure in every form of pulmonary tuberculosis, and does not hesitate to-day to say that the more cases he sees, the less capable he feels of predicting what cases may recover.

THE SANITARIUM FOR CONSUMPTIVE SAILORS ESTABLISHED BY THE U. S. MARINE-HOSPITAL SERVICE AT FORT STANTON, N. M.*

J. O. COBB, M.D.

Passed Assistant-Surgeon U. S. Marine-Hospital Service.

There is a prevalent belief that the sailor does not have tuberculosis. Just because he is at sea breathing the purest air, we naturally think of him as having the greatest chance of escaping that dread infection. The public mind loses sight of the fact that the average sailor spends a great part of his life ashore, participating in drunken brawls, gambling in the vile holes underground in New York, San Francisco, Baltimore and all the large maritime commercial centers. The little time he spends in sleep when ashore is in sailors' boarding houses, usually filthy dives. In lieu of beds, there are rough bunks built in like a set of shelves clear up to the ceiling. Into these a little straw is thrown, and this is the bed of one man one night, and another the next, and so on. The straw is rarely, if ever, changed, and this is the only bed these poor fellows can get in these gruesome places. They need no covering, for the room is heated by the red-hot stove, around which some sit and smoke and play and drink the long night through. Such a place is awful to think about, and it is small wonder these men get sick; it is the greater wonder that every man so exposed is not rapidly infected with tuberculosis, for it is only reasonable to believe that all such places become infected in time. Then, too, it is easy to understand that one case may infect an entire crew at sea, on a long voyage, when the men are packed together like animals. The sailor breathes pure air only when he is on watch, but when he "turns in," his sleeping quarters are exceedingly bad—so bad that it is impossible to describe them adequately without subjecting one to the accusation of exaggeration.

We of the Service know that there are hundreds of consumptive sailors and our annual reports show that we have been losing on the average more than one hundred a year. Our experience has also shown us that all the cases that remain, eventually die; they drift in and out of the different hospitals, better to-day and worse to-morrow, and sooner or later serve as a necropsy study.

So it has gone for years. No movement which is purely humanitarian can be brought about without much talk and some sober writing. Officers talked and thought, but unfortunately this is usually what a non-political government officer is compelled to do—take it out in talking or writing, if he is so inclined. It is to the credit of the officers of this Service that they have ever been alert to the prevention, spread and proper control of the disease. The present surgeon-general, Dr. Walter Wyman, was the first in the Service to undertake to bring about the establishment of a sanitarium

for consumptive sailors. This was many years ago, when he was a junior officer, and his efforts met with little encouragement, and although unsuccessful he can properly lay claim to being a pioneer. He did not forget his early ambitions after he became surgeon-general, and it fell to his lot to be the instrument to consummate the establishment of a sanitarium for consumptive sailors.

In December, 1898, I was sent on an inspection tour through Arizona and New Mexico for the purpose of recommending a site for the contemplated sanitarium. It was thought that as the Army had been first in the arid region—and of course had selected the most desirable site—abandoned posts would best serve our purpose as fulfilling the requirements of large tracts of land, good water-supplies, and that the data published by the officers of weather and other conditions would be very valuable as a guide.

After careful examination in both territories, Fort Stanton was decided on as best suited for our purposes, specifically and generally. The weather reckonings made by the Army, together with special reports by its officers serving here in early times, showed mild winters with delightfully cool summers. It was necessary to have a station combine these two features or else have two stations, one at a low level for winter and one higher for summer, for otherwise patients at low levels would lose in summer all that had been gained in winter. With its undoubted advantage of climate, a generous water-supply, a large reservation, and with substantial stone buildings, it is an ideal site.

Fort Stanton is situated in Lincoln county, N. M., about six miles from Capitan, the terminus of the El Paso and North Eastern railroad. The elevation is 6126 feet. There are 38 buildings of various sizes, most of them being of native rubble stone. There is room for 200 patients, besides quarters for officers and attendants. Repairs are going on, and new machinery is to be introduced shortly. A modern steam laundry and a one-ton ice plant are now contracted for, and the plans and specifications are being prepared for an electric-light plant and extensive repairs, which will be commenced this fall. All the buildings will be made modern, and the appointments will equal any in the country. The laboratory will be conducted especially in the study of the disease. The apparatus for the purpose is already on the ground and is temporarily set up in an old building awaiting the repairs on the executive building, where there is to be a room set apart for this purpose.

There is a large farm and garden attached to the station. On the farm is being raised all the forage and most of the grain for the horses, milk cows, cattle and chickens. In the garden we have a source of pleasure not often found at such institutions. We have all the fresh vegetables needed and such vegetables as can not be bought in this vicinity.

Our herd of Jersey milk cows supply all the milk used at the station. While speaking of the cows it may be of interest to describe the method of dealing with the milk. We try to have all the patients follow as nearly as possible the method of taking milk described by Dr. Bulkley. To make this a little more perfect, it is necessary that the milk should always have a given percentage of fat, and to obtain this our milk is cooled immediately after milking and run through a separator. The cream will turn out about 20 per cent. (volume), and then this is added to skim milk to make it 20 or 25 or 30 per cent. of cream, as is prescribed by

* Presented in a Symposium on Tuberculosis, to the Section on Hygiene and Sanitary Science, at the Fifty-first Annual Meeting of the American Medical Association, held at Atlantic City, N. J., June 5-8, 1900.

the medical officer. In this way there can be no marked variation in the percentage of fat, as there would be on cold, stormy days, or in change of food or other well-known conditions, which cause a cow's milk and fat percentage to vary. We make some butter and buttermilk, and intend to make kumyss and kefir. The patients have increased so fast that we are not altogether prepared for them, as it was the original intention of admitting only twenty the first year.

The reservation is very large and it is the intention to fence it for pasturage and to raise our own sheep and beef cattle. This will be not only a paying operation, but will give us fine meats. Fort Stanton was set aside for a sanitarium by executive order dated April 1, 1899, and the station was officially opened April 27, 1899. To date, there have been 70 patients admitted, 6

was a good case when he came. This man had a small cavity, which apparently began to heal. He began to improve after a time, and would probably have recovered but for contracting tubercular pneumonia on his good side. Of the other cases which died, not one was considered, by the officers of this station, as good subjects for transfer and all died shortly after reaching here.

To get the patients constantly in the open air has been a most vexatious problem. Most of them came here from hospitals which were heated by steam and would have rebelled had they been put in the open air to sleep immediately after arriving. Knowing the sailor as I do, it was necessary that all this be brought about indirectly and gradually, and now they are very enthusiastic for the open air. All sleep with the windows

Fig. 1.—Looking Across Parade Ground, Showing Capitan Mountains to the Northeast. Fort Stanton, New Mexico.

of whom have died, 17 have been discharged, 13 improved and 4 apparently recovered. Seven of those discharged improved are still under observation as attendants, of whom all appear to be well except one. This one, however, was a bad case, but is now doing well. Three cases reported as improved are working in the vicinity. One of these had both lungs badly infected, but as he was strong, had no fever and felt well, he obtained employment at the mines near here and is working every day. He reports that he still expectorates, but otherwise is still improving. Another has just gone to work. His trouble originally was a tubercular empyema, for which several ribs were partially resected. The other man is well.

Of the cases remaining we shall probably lose a few, but seven seem to be entirely cured and will be sent back as soon as the hot weather in the East is over. Twenty-one seem to be arrested, though the evidence of the disease still persists. Of those who died, only one

and doors open and in the day they walk about, swing in hammocks here and there under the cottonwood or box-elder, ride the native ponies, which are for their use, or recline in easy chairs or amuse themselves by doing odd jobs. One man bought himself a tent and sleeps out every night.

To find means of diversion or amusement has been a hard problem and will continue to be so. While they are running fever or otherwise doing badly, I have no trouble, but just as soon as they begin to get well they become restless for the water again.

In the matter of treatment, dependence has been put in keeping the men constantly in the open air and giving the best of food. However, we have had several cases with severe complications, malaria being the most frequent. Many of the men have come from localities where they had previously contracted malaria and I have several under observation whose blood is full of the organisms. Dyspepsias are exceedingly troublesome and

loss of appetite in some cases hopelessly discouraging, these complications compelling us to give some medicine. Just here it is well to warn those who have not had much experience with heroin to be cautious in its use; the habit is easily contracted.

We have been making some careful observations of the effects of altitude. If mere clinical experience is worth anything at all it leads me to believe that most cases become anemic shortly after coming here. Of course, the

We have noticed, too, that the majority of patients lose weight at first, notwithstanding an improvement in appetite and general symptoms. The urea percentage is markedly increased and has continued so with cases under observation.

The method of handling the sputum and preventing infection will be of interest to the general practitioner. There are no half-way measures. The theoretical part is nearly perfect and its execution will be almost so

Fig. 2.—Group of Buildings Looking North, Showing the High Mesa on the North, Which Affords Protection from the Winds. Fort Stanton, New Mexico.

Fig. 3.—Looking up South Avenue, with Cottonwood Limbs Which Were Set Out This Spring, and Which Are Growing, Showing White Mountain to the Southwest and Mesa to the South. Fort Stanton, New Mexico.

blood-count and hemoglobin percentage are increased, but I feel certain, without being able to prove it, that it is not actually true. After a time the proper relative standing of the blood-count and hemoglobin is re-established and the increase is actual, though the readings of the instrument may not show an increase. I have about decided that the blood tests can only be of interest as a guide to each individual's relative condition and progress. Collectively the tests are misleading.

when all the sterilizers are put in place and in operation. I am happy to say that there is no such thing at this station as a spittoon, which demonstrates that a large institution can be run without one. To have a spittoon lowers the standard of effectiveness at once, for no one will deny the dangers of using one. Any method of sterilizing sputum except by incineration or steam heat is not perfect. Any method of collecting sputum by the process of ejecting it from the mouth

at a distance into the receptacle is unsafe. The receptacle must be brought close to the mouth. The patient must have his own receptacle, for it is disgusting to see the sputum of others. To secure this desirable end we are using sputum-cups and pocket sputum-flasks. Each patient has a flask and cup, but I find that they dislike the cup and nearly all of them want to use the flask only. They carry the flask wherever they go, and use the cup in their rooms. Once a day at a given time every cup and bottle used is carried to the sputum room and there checked off, by an attendant, from a complete list of the patients. It is necessary to keep close watch on each individual, and take note whether he brings his cup and bottle for sterilization, for if we did not he might empty the sputum in the sinks and closets.

exasperatingly narrow in matters pertaining to public health, unless his pocket or the pockets of his constituents are affected. Let a wide and destructive epidemic break out among animals and then how prompt legislation is introduced to suppress it! There is money! It is a crying shame that this country, one of the foremost in all grand evolutionary movements, should have waited till the closing year of this century before joining in the sanitarium movement against the worst foe of mankind. What excuse can there be? None. The results were well known to be good and it was not an experiment. The way had been marked for us by Germany as a leader in this crusade with her first sanitarium in 1859 at Goerbersdorf, or 1854 according to Von Leyden. The impetus given to the sanitarium move-

Fig. 4.—Showing Officers' Quarters. Fort Stanton, New Mexico.

The cups and flasks are placed in the sterilizers and the temperature maintained at 245 F. for one hour. This heat thoroughly liquefies sputum and makes it easier to cleanse the receptacles. Our large sterilizer is not yet set up, but when it is, all the bed clothing and individual clothing will be sterilized by steam or formaldehyde at regular intervals.

We, as a nation, lead in all matters for the social and political welfare of man; but we have been very strangely conservative in helping him in his sickness, especially in the inner circles of his home, there to combat the ravages of disease which are undoubtedly caused by the greed of owners and keepers of unhygienic tenements, where men, women and children have less care than given domestic animals. The average legislator is

ment by the patronage of the Kaiser has done much to stimulate still greater effort. Even the life insurance companies of Germany have built sanitariums for their insured. This has been found by the insurance companies to pay, and it is to be hoped that the companies of this country will soon follow their example. The mutual aid companies should, by all means, do something for the many consumptives that they undoubtedly have in their membership. I have heard that the question had been seriously presented to the A. O. U. W., as this order has hundreds of consumptive members.

The philanthropy of our large cities has long done much harm in building large hospitals for this class of patients. Wealthy men and women have subscribed immense sums to erect places where the poor consumptive

could go—and die. Make them comfortable has been the cry. The end was inevitable. A new sense of responsibility is now thrust upon the man or woman with means, for this frightful death percentage must not be. The case must be diagnosticated early, and that is the awful and fearful responsibility of the physician. And look to it, you who make snap diagnoses with the unaided ear through the patient's clothing and who never make thorough chest examinations of women at all; and you who stain one specimen of sputum and finding no bacilli declare the trouble is only bronchitis; or you, the worst offender of all, who withhold from the patient the true nature of his disease, allowing him to go on expectorating at random, spreading the infection broadcast and to the members of his own household! And I say to these, and to the vampire who holds his patients, drugging them to death, finally abandoning them when

of the sufferer. Man of wealth! Woman of wealth! Do not put your money in grand edifices in the overcrowded city where your philanthropy only serves to make a doomed consumptive's life comfortable for a few weeks, at most a few months. Go to the woods all about you and put these people where there be a chance for recovery and where they can return to their homes as useful citizens. This much may be hoped for, but even if the patient should die, we will have gained by removing the focus of further infection. This to me is the most important side of the subject.

All grand movements begin first with suggestions, and nothing is ever accomplished except by discouraging set-backs at every turn. Men like Trudeau can not help but look back with pride on their accomplishments under the very great drawbacks they must have had. What a sensation must the founder have had when, in 1884,

Fig. 5.—Patients Ready for a Ride. Fort Stanton, New Mexico.

they are in a dying condition by sending them to health resorts, that the public is awakening to the realization of the danger of delay and in wrath will have none such as you.

The phthisiologist can not make a successful issue of this undertaking without the aid of the conscientious general practitioner. I am told that in Germany every medical man is alive to this question and the most careful examinations are being made to detect the disease in the first stage. Tuberculosis is not such a grave disease if taken in hand early and sent into the pure open air.

Medical men have no money, and if they diagnosticate their cases early, then the public at large, through its wealthy citizens or its municipal, state or governmental authorities, becomes obligated morally as well as socially and economically, to give the patient a chance for his life. It must not be care for the doomed and dying—the reach must be greater and extend out to the home

the first little cottage at Saranac Lake came into existence! His was no selfish ambition; what must have been his hopes and fears? He would have been laughed at then if he had gone to Congress or his state with a bill to help erect this same little cottage. His patients were not animals and there was no loss to the pocket-book, therefore no need of legislative help.

The public is awakening to the danger. Towns, cities and states have become aroused, and now that our Government, first through the Marine-Hospital Service, and then very shortly after by the Army, has given its official cognizance of the importance of stamping out the white plague we may expect in the next few years a forward move of which we may be proud. Of course the work done by the government may, in a way, be comparatively small; yet of all the patients admitted were not one to recover, were all to die, still the good done to the community at large is inestimable. So many foci of infection removed from our ships in their world-

encircling courses; from the barracks, camps and hospitals of the Army; from the sailors' boarding houses; from our general hospitals, where we have had such a discouragingly large percentage of deaths; to bring all these patients away from thickly populated centers, where they undoubtedly spread the infection, is an accomplishment itself of no mean importance.

With such opportunities for study afforded by the well-equipped laboratories, the ever-increasing clinical material, it is not too optimistic to feel that we may help to clear away one by one the many obstacles yet confronting the world in effecting immunity and cure.

THE DIAGNOSIS OF PULMONARY TUBERCULOSIS.*

ROBERT H. BABCOCK, A.M., M.D.

Professor of Clinical Medicine and Diseases of Chest, College of Physicians and Surgeons.
CHICAGO.

This may be a matter of considerable ease, or one of extreme difficulty, depending on the stage of the disease and the nature of the pathologic conditions. Let us first consider its diagnosis at an early stage. Tuberculosis is an infection, and as is the rule with infections, it occasions a rise of temperature even at its incipency. Except very rarely, when the infection is violent or the disease assumes the acute pneumonic type, the elevation of temperature is slight, and may even escape detection. If the thermometer shows a constant elevation ranging between 99.5 and 100.5 F. it is in a suspected case strongly suggestive of tuberculous infection. The temperature should be registered not merely night and morning, but four or five times daily, as, for instance, at 9, 12, 3, 6, and 9 o'clock.

The pulse is as a rule accelerated and to a rate out of proportion to the temperature. But it should be remembered that an individual who is under examination for suspected tuberculosis is apprehensive, and therefore likely to manifest acceleration of the pulse. Nevertheless, the pulse-rate will be found to range in the neighborhood of 100, not rarely 120 or thereabouts. It is usually a pulse of low tension, and this is more characteristic than is an increase in the frequency. Undue rapidity together with lowness of tension forms, in the absence of signs of cardiac disease, a very suspicious combination. Its value is enhanced if conjoined with increased body temperature between the extremes noted above. The respirations may or may not be increased. As a rule in incipient cases there is neither subjective nor objective dyspnea.

Loss of weight is a suspicious circumstance and is an invariable accompaniment, I believe, of incipient pulmonary tuberculosis. It is so often slight that it escapes the patient's notice, and therefore his statements are not always to be relied on. The value of this symptom depends largely on its association with other symptoms and signs; without them it can scarcely be held suggestive of tuberculosis. Decline in strength and a falling off of appetite are also to be considered in doubtful cases, but are of small evidential value.

Cough is usually the first symptom to attract the patient's attention and to arouse apprehensions in the mind of the physician as well as of the patient. Although it is frequently symptomatic of this disease even at its invasion, still it is not invariably so, and taken

alone has but small diagnostic value as regards pulmonary tuberculosis. In every suspected case in which physical signs do not settle the question of diagnosis, the upper respiratory tract and the heart should be carefully examined for an explanation of the cough. Aurists will say that even the ears should not be left out of the field of exploration.

Sputum is, for a time, absent in cases of beginning tuberculosis, but when present exhibits nothing characteristic, unless, of course, it shows bacilli. Even hemoptysis is not pathognomonic, although highly suspicious of pulmonary tuberculosis. When due to a deposit of tubercles in the lungs, physical signs may usually be detected, or there will be some of the suggestive symptoms already considered. If not, then the gums, nares, pharynx, larynx, trachea and heart should be carefully examined for a possible source of the hemorrhage. In conjunction with a slight rise of temperature, loss of weight and cough, hemoptysis is very significant of tuberculosis.

Having thus briefly considered the symptoms attending incipient pulmonary phthisis, let us now take up the physical signs.

Inspection.—It is not necessary to describe the contour of the thorax with its narrow apices, weak musculature, deficient expansion, etc., which so often characterizes the tuberculous diathesis, and therefore I shall pass at once to such evidence of early tuberculosis as may be gained by inspection. One of the earliest ocular signs is lessened expansion of one apex, observed oftentimes in the supraclavicular region. This area may be slightly depressed or retracted, as compared with the other side. In rather more advanced cases, or those in which the tubercular deposits are not so limited, these same changes may be noted in the infraclavicular region as well. Another sign, which by Carcassonne and others is said to be a very early and significant one, is atrophy more or less marked of the scapulothoracic muscles on the affected side. That drooping of the shoulders, which forms a part of the phthisical habitus, may be seen at an early stage, but is generally present when the disease is more advanced.

Palpation does not furnish very reliable or positive information, except as corroborative of that obtained by other means. It is so largely influenced by individual peculiarities. Nevertheless, when in a suspected case there is a marked difference in the pectoral fremitus of the two apices, that apex at which it is distinctly increased is apt to be the seat of tubercular deposits. This is particularly likely to be the case in the suprascapular region, for a reason which is about to be stated.

Percussion is the means generally relied on for the detection of incipient pulmonary tuberculosis, yet it is not always so trustworthy as is auscultation. Dulness at either apex is usually regarded as indicative of the disease in question, and justly so. Still, one should always bear in mind that it is normal for some individuals to show slightly less resonance at the right apex. Therefore, in estimating the importance of trifling dulness at the right summit of the chest one must also consider the presence or absence of the symptoms of constitutional disturbance already spoken of. It is common for physicians to look for impaired resonance below the clavicle, whereas in my experience it is far more frequently found above and in the suprascapular region. This is owing to the fact that, as pointed out by Fowler, deposits of tubercles take place earlier and more frequently at the tip of the lung, along its borders and on the posterior aspect of the apex. Consequently a small

* Presented in a Symposium on Tuberculosis, to the Section on Hygiene and Sanitary Science, at the Fifty-first Annual Meeting of the American Medical Association, held at Atlantic City, N. J., June 5-8, 1900.