were the seat of a chronic endocarditis upon which the acute process implanted itself; in one of these cases the bacteriologic examination failed to reveal any bacteria and it would seem that the inflammatory and necrotic changes in the endocardium had fallen into temporary or permanent quiet suggesting the probability of recovery from the acute symptoms; this corresponds well with the clinical facts in the case which show that the patient died from the effects of an uncompensated valvular lesion rather than from an acute infection (Group I, Case I). In one of these instances of acute, destructive changes developing upon sclerotic valves was a history of previous attack of rheumatic fever obtained to which the chronic endocarditis could be traced.

Otherwise the anatomic and histologic changes presented by these specimens are so typical and characteristic of ulcerative endocarditis as to require little elucidation; they show vegetative and necrotic changes in profusion with the production of typical valvular aneurisms and subsequent rupture. It is quite noticeable that in only four of the cases were there evidences of embolism and infarcts and here the embolic changes were quite limited; as a rule the showers of infected emboli from the endocardial vegetations and ulcerations constitute a striking feature in the clinical as well as the anatomic manifestations of this form of endocarditis.

A number of facts and considerations in regard to the etiology and pathology of acute ulcerative endocarditis may be summarized as follows:

1. The disease is caused by the localization of microbes in the blood upon the endocardium. The streptococcus pyogenes, the micrococcus lanceolatus (pneumococcus), the staphylococcus pyogenes are found most frequently and in the order named. Various other microbes have been found also, such as the pus microbes, the bacterium coli commune, the gonococcus, the bacillus typhosus, as well as bacteria that have not been found in other diseases and are in so far as yet peculiar to malignant endocarditis. Recently the bacillus diphtheriae was found in the endocardial lesion (Howard, *J. Hosp. Bull.*, April, 1893). Experimentally the disease can be produced by first causing some mechanical or chemical injury to the valvular endocardium and subsequently injecting cultures of various bacteria into the blood, or by intravenous injections of cultures of staphylococcus pyogenes aureus on potato suspended in water with scrapings from the potato when the micrococcus agglomerations attach themselves to the endocardium more readily than the single cocci (Rosenbach, Wyssokowitch, Weichselbaum, Prudden, Fränkel and Sängner, Netter, Gerarieu, Ribbert).

2. Acute ulcerative endocarditis is met with as a secondary lesion in the acute infectious diseases, notably and most frequently pneumonia, also in meningitis, acute articular rheumatism the specific fevers, gonorrhoea, dysentery and so on. It is frequently incorporated into suppurative, septic and pyæmic processes. Finally it may occur as a primary or cryptogenetic affection, developing without any known or demonstrable infection atrium. From the mycotic endocardial foci microbes and infected fragments may be carried away by the blood current and give rise to embolism in the various parts of the body, local necroses, and secondary pus accumulations.

3. In over one-half of the cases (three-fourths according to Osler), the acute ulceration is implanted on an old valvular lesion, the absence of endothelium and the roughness presenting assailable points for microbic invasion. This is the only competent reason for the occurrence of an overwhelming majority of the instances in the left heart but the explanation is not complete, because of the cases affecting primarily healthy valves, over two-thirds occur in the left side and about one-third in the right half.*

EXPLANATION OF PLATE.

FIG. 1. Acute ulcerative endocarditis of tricuspid valve with rupture of valve aneurism through which match has been passed. Group I.

FIG. 2. Acute ulcerative endocarditis engrafted on sclerotic aortic valves; perforated aneurism; necrotic spot in center of anterior mitral flap. Chronic mitral endocarditis. Group II, Case 1.

FIG. 3. Acute ulcerative endocarditis of mitral valves following lobar pneumonia. Group IV, Case 1.

FIG. 4. Acute ulcerative endocarditis of aortic valves; ruptured aneurism; necrotic spot in center of anterior cusp. Group IV, Case 3. Acute leptomeningitis.

THE INFLUENCE OF HIGH ALTITUDES ON THE ARREST OF PULMONARY PHTHISIS.

BY BOYD CORNICK, M.D.

KNICKERBOCKER, TEXAS.

A generation ago, in prescribing change of climate for consumption, the usual medical advice was that the patient should take a sea-voyage, should spend a period of time in travel, or should resort to some locality with a mild and equable temperature; little regard being paid to other factors of climate than temperature alone. But with the appearance in 1864 of Dr. Hermann Weber's "Climate of the Swiss Alps," began a change in medical opinion in favor of high altitudes, a rarefied atmosphere being esteemed the most important climatic factor. The latest phase of this changed opinion has been well formulated by James Alex Lindsay, M.A., M.D., in his work on "The Climatic Treatment of Consumption." Says Dr. Lindsay:

"The characteristics of mountain climates which are fairly constant, are as follows: Increased rarefaction, directly proportionate to the elevation; great purity of the air, and freedom from organic contamination; much ozone; free solar radiation; abundance of light, and freedom from fog. The tendency of mountain climates is towards dryness rather than humidity." . . . "There can be no doubt that the most essential of all is the rarefaction of the air, the only important factor common to all the mountain sanatoria of proved efficacy."

Dr. S. E. Solly, author of the article on climate in Hare's System of Therapeutics (H. C. Lea & Co., Philadelphia), arrives at the conclusion that: "In prescribing a particular resort for a given case of consumption" . . . "the broad truth remains that of all climatic factors, in the treatment, as well as in the prevention of phthisis, elevation is by far the most powerful of them all."

* Out of seventy-eight cases of ulcerative endocarditis of healthy valves, twenty-four were right sided and fifty-four left sided.

NOTE.—In Hamilton Text-Book of Pathology, Vol. I, will be found references to the more important literature on ulcerative or malignant endocarditis.

I quote at second hand from a reprint of an essay in The National Magazine, February, 1892: "Can Consumption be Cured?" by A. J. Meuer, M.D.

Fig. 1.

Fig. 2.

Fig. 3.

Fig. 4.

THE JOURNAL OF THE AMERICAN MEDICAL ASSOCIATION.

And Dr. C. Theodore Williams, in his essay entitled "Influence of Climate on Health," (v. a Treatise on Hygiene and Public Health, Stevenson and Murphy, editors; P. Blakiston, Son & Co., Philadelphia), after enumerating and recommending different mountain climates for phthisis, summarizes his views as follows: "Though these groups vary considerably in climate as regards temperature and moisture, they all agree in the reduction of the barometric pressure, and this is the essential feature of the whole series."

But no climatologist, probably, has exerted a greater or more widespread influence on the English-speaking medical world, in advocating the claims of high altitudes, than has Charles Denison, A.M., M.D., whose wonderfully able, though rather too condensed little pamphlet: "The Preferable Climate for Consumption," has been reprinted from the "Transactions of the Ninth International Medical Congress," for free general distribution, by order of the legislature of Colorado.

In discussing the comparative importance of different climatic attributes in the arrest of chronic pulmonary disease, while cataloguing five principal, and five confirmative factors of climate, Denison gives first rank to "*dryness* as opposed to moisture." Second, "*coolness* or cold preferable to *warmth* or heat," and third, "*rarefaction* as opposed to *sea level pressure*." And under caption third he says:

"That altitude is not made to precede dryness and coldness may need to be explained, in view of the great prominence given to elevation by Jaccoud in his excellent work on pulmonary phthisis, which, by the way, contains a better appreciation of the benefits of climatic treatment than any other similar book extant. The difference is this: In the present classification *the qualities which altitude produces* (*italics mine*) are given their due prominence without special reference to their causes, while rarefaction is considered as an individual quality, with special reference to its mechanical effect upon the respiratory function."

It would seem that no other construction can be placed on this language than that in the author's opinion various other climatic factors, in addition to aerial rarefaction, are caused by altitude; and that of the factors so caused, *dryness* and *coolness* or *cold* are of first and second relative importance respectively. Indeed the author states explicitly in a footnote on page 8: Upon temperature, elevation has a constant effect in the production of cold," and again (p. 4), "Cold is probably the most important factor in the production of dryness, and elevation is not far inferior." But we shall farther on see that the extremest degree of dryness may be found almost at sea level and in connection with a high mean temperature; while it must be evident, to whoever will consider the matter a moment, that the terms *coolness* and *cold* are purely relative and, aside from their implication of an influence on the absolute humidity of the atmosphere, are utterly indefinite in their significance.²

² Without going out of the way to propound the counter-thesis (as being beside the question) that, the degree of relative dryness being the same, a comfortable degree of coolness is always preferable to uncomfortable heat; and that *warmth* is, in a majority of cases, not only more agreeable but also more conducive to the welfare of feeble and anemic invalids than is *cold*, it will suffice in this connection to advert to the fortunate fact that there are not a few other factors besides altitude which contribute to the production of agreeable temperatures, among which not the least important is a constant and brisk air movement in summer in some very dry localities, which, from their only moderate

And to buttress the chain of reasoning of thinkers whom the medical world delights to honor, statistics may be had in sufficient abundance to demonstrate that actual results observed at high (and dry) altitudes are incomparably better than at low (and humid) altitudes; citing for comparison Austin Flint's 670 cases, with but seven cured and no statement of number benefited, in a compilation of 1,809 cases sent to "*various low climates*," of which number 1,000 had been reported by one other authority.³

But while statistics may be marshalled to prove or disprove any debatable proposition, although when collated from a past generation they are too often valueless as bearing on the climato-therapy of the present day, yet a most valuable summary is furnished in juxtaposition with these cases, demonstrating that in the Swiss Alps and at Colorado resorts with an altitude of 4,500 feet and upwards, results have been attained such as have certainly never been reported from the "*various low climates*" to which the 670 consumptives resorted as instanced by Dr. Flint, or to which the unfortunate 1,000 repaired, as quoted from Dr. C. J. B. Williams.⁴

The fact being admitted then, that about 37 per cent. of the patients sent by several reputable observers to the Swiss Alps and to Colorado have been reported cured, while about 74 per cent. have been benefited, it does not seem remarkable that the view expressed by Dr. Lindsay should have become a widely accepted medical doctrine, viz.: "There can be no doubt that the most essential of all (the characteristics of mountain climates) is the rarefaction of the air, the only (?) important factor common to all the mountain sanatoria of proved efficacy." But having just enumerated six different "characteristics of mountain climates which are fairly constant" and having further stated that "the tendency of mountain climates is towards dryness rather than humidity," this conclusion by Dr. Lindsay of the unimportance of all other factors excepting rarefaction alone, would look to an unprejudiced reader to be hardly so free from doubt as it evidently appears to his mind.

Let it not be forgotten in this connection that so acute an observer and thinker as Dr. Denison, in his catalogue of the comparative importance of different climatic attributes, after asserting as quoted above, that "*dryness*" and "*coolness* or *cold*," (to which he gives first and second rank respectively) are among the qualities which altitude produces, gives only *third* rank in order of prominence to "*rarefaction* as opposed to *sea level pressure*." But if Denison be correct in classing "*dryness*" and "*coolness* or *cold*" among the qualities, and indeed, as the very most important qualities which altitude produces, it follows that Dr. Lindsay must be in error when he claims that "the rarefaction of the air (is) the only important factor common to all the mountain sanatoria of proved efficacy."

A logical result of the enthusiastic claims made for a generation past in behalf of the influence of the rarefaction of the air on the arrest of phthisis, was manifest in the line of treatment pursued by a noted and capable clinical lecturer of Louisville, Ky., who, in 1877-78, during my term of service as

altitude and southerly latitude, one might expect to find uncomfortably warm or, indeed, depressingly hot.

³ Hare's System of Therapeutics, Art. Climate. H. C. Lea & Sons, Philadelphia.

⁴ Ibid.

one of the house physicians to the city hospital, put his patients, in the acute progress of pulmonary phthisis, through a series of calisthenic exercises with a view, by deep and forced inspiration and expiration, to obtain the same benefits of expansion of the acutely inflamed and infiltrated lung tissue, which are claimed as due to the rarefaction of the air in the Swiss Alps and in Colorado.

The same influence, no doubt, contributed not a little to the invention and quite widespread use of the pneumatic cabinet and kindred apparatus, which by utilizing the expansive tendency of compressed air brought the mechanical advantages generally claimed for the aerial rarefaction of high altitudes into the inner sanctuary of the doctor's office, thus placing "the only (?) important factor common to all the mountain sanatoria of proved efficacy" within the reach of every patient, without the necessity of leaving home. Judging, however, by the absence for a year or two past of all reference in current medical literature to the pneumatic cabinet, as a factor in the arrest of phthisis, the casual observer would suppose its use has failed to fulfill the hopes of its advocates.

Dr. Denison's claim, as quoted above, that the prime factor, *dryness*, is one of "the qualities which altitude produces," qualified as it is by Dr. Lindsay's assertion that "the tendency of mountain climates is towards dryness rather than humidity," may not, however, be accepted without further qualification. For in "Denison's seasonal chart, showing combined atmospheric humidities for winter, 1883," (op. cit.) Pike's Peak, with an elevation of 14,134 feet, appears as a humid blue oasis in the midst of the lurid red which marks the otherwise dry and arid Rocky Mountain plateau. But this very dry area, designated on the chart by a uniform red color, is so extensive as to demonstrate that whether or not elevation be an important factor in the production of dryness, it certainly is not essential thereto. For, according to the chart, this arid red region includes not merely the noted dry table-lands of Colorado, situated more than a mile and a half nearer to sealevel than is the humid summit of Pike's Peak, but it extends south and west without a break in the uniformity of its tint to nearly the borders of salt water, including Yuma, Arizona, where the altitude is some 13,993 feet lower than the summit of Pike's Peak, and but 141 feet above the sea.

Of course, in calling attention to this fact I must not be understood as claiming that so capable an authority as Dr. Denison has ever advocated the resort to an altitude such as that of Pike's Peak for the arrest of consumption, though he does mention "10,000 feet as a possible extreme in the southern portion (of the United States) in summer." (Op. cit., p. 14.) My purpose, on the contrary, is to show from the authority of a painstaking statistician, who has gone to the prodigious labor of compiling "some eight million separate signal service observations into the Annual and Seasonal Climatic Maps of the United States," (op. cit. p. 2) that, while dryness and altitude may be and often are coincident, the one is by no means a necessary consequence of the other. And that, while "the tendency of mountain climates is towards dryness rather than humidity," at least in those altitudes where the arrest of pulmonary complaints may be hoped for, yet as a matter of fact, the very greatest degree of possible dryness

may also be found at altitudes within a few hundred feet of sea level. And not only is this true, but there are elevations also, well within the limits recommended by our advocates of high altitudes, such as those along the seaward slope of coast mountain ranges, which very often have an exceptionally high percentage of humidity by reason of the impact against them of the prevailing humid sea-breezes,⁵ and for that very reason, there seems little room for doubt, are specially ill adapted for consumptives as a health resort.

A consideration of this fact seems to have led Dr. A. J. Meuer, while discussing the cognate question of an altitude of immunity from phthisis, to the conclusion, after enumerating the points of superiority in Colorado's climate, that (op. cit. p. 5) "in the mountain region between the eastern slope of the Rockies and the Pacific coast the conditions vary to such an extent that *altitude alone can scarcely be said to enter into the account at all*, unless accompanied by the long list of favorable factors above enumerated. *Accompanied by these conditions, however, altitude is everything!*" (Italics mine.)

It is a fact beyond question or cavil that both in the Swiss Alps and in Colorado, at altitudes ranging from 4,000 to 6,000 feet above sea level, a larger percentage of consumptives is benefited by the climate than is recorded of any of the popular resorts of a generation ago, most of which were characterized by not merely sea level pressure but also by great humidity of the atmosphere. What are the factors of climate which bring about these later so gratifying results? The coincidence that a high altitude, and consequent rarefaction of the air, is common both to the Swiss Alps and to Colorado is so evident that, on first glance, many otherwise cautious observers have jumped to the conclusion that aerial rarefaction is *the* curative influence; or is, at least, *a* curative influence of paramount value. And much labor has been expended in the production of many, not always harmonious, and often unphysiological explanations of how the attenuation and lessened pressure of the air produce the beneficial results that have there been observed.

The Swiss Alps, like Colorado, are mid-continental in their situation. The sea breezes which penetrate so far into the interior have been robbed of their moisture by every chain of hills and every range of mountains over which they blow. The driest atmosphere of all Europe, at least in winter, is doubtless in the Alpine region of Switzerland; although Davos, its most noted health resort, with nearly the same altitude as Denver, hardly more than approximates

⁵ Prof. E. A. Parkes, in his work on Practical Hygiene, in discussing the climate of India (Vol. ii, p. 336. Wm. Wood & Co., N. Y., 1883) says: "The humidity of different parts of India varies extremely; there are climates of extreme humidity—either flat, hot plains, like lower Sindh, where, without rain, the hot air is frequently almost saturated, and may contain ten or eleven grains of vapor in a cubic foot; or mountain ranges like Dodabetta in Madras, 8,640 feet above sea-level, where during the rainy season of the year the air is almost saturated; a copious rain at certain times of the year may make the air excessively moist, as on the Khayab Hills, where the southwest monsoon parts with its vapors in enormous quantities."

"When the southwest monsoon, loaded with vapor, first strikes on high land as on the Western Ghats, on the Malabar coast, or on the mountains of Tenasserim, and especially on the mountains of the Khayab Hills, at some points of which it meets with a still colder air, a deluge of rain falls, as for example, at Cannanore (Malabar) 121 inches; Mahableshwur, 253 inches; Mouleim (Tenasserim) 180 inches; Cherrapoonjee (Khayab Hills) 600 inches." On the succeeding page the statement occurs that "in inland districts the southwest monsoon having lost most of its water as it passed over the hills, may be comparatively dry."

And Dr. Denison, in a foot note to page 8 (op. cit.) referring to the favorable location of the eastern slope of the Rocky mountains, mentions "the protection of mountain ranges which drain western humid air currents of their moisture."

to that degree of low relative atmospheric humidity which characterizes the Rocky Mountain plateau.

During the winter months while this Alpine valley is clad in a mantle of crisp, dry snow, consumptives from the lower and damper levels resort thither on the advice of the most capable physicians of every nation of Europe. And though advanced or acutely progressing cases of phthisis seldom do well, the most gratifying improvement is often noted in the incipient stages. But with the breaking up of winter and the melting of the snows in spring, there is a hegira from Davos. A uniform consensus of medical authority opposes a prolongation of the invalid's stay into the spring of the year. Why?

If a rarefied atmosphere be the factor of climate which arrests the progress of phthisis at Davos, why must invalids who have made favorable progress during the winter months fly for their lives on the approach of a milder temperature? The altitude is not appreciably diminished by the melting of the snow. The atmospheric pressure remains "fairly constant" in spring, as in winter, at about two and one-half pounds per square inch less than at sea level. The act of respiration continues just as frequent and just as deep as before, and the infiltrated pulmonary tissue, inflamed and sensitive though it be, is no less thoroughly expanded by the deeper and more frequent respirations, and no less permeated by the attenuated air, than when the thermometer was below zero and the snow was crisp and dry.

Dr. Denison's explanation of this noteworthy and most important fact is (op. cit. p. 14) "that we in America have an immense advantage over most European high climates, in that we keep up the curative effect by suitable increase of altitude in summer. Instead, they are compelled, as at Davos, St. Moritz, etc., to give up the chosen climatic treatment during warm weather." And on page 7 the statement is made: "It is to secure the cooler temperature in summer time that some of the phthisical patients from the plains are sent higher up to the parks and divides of the Rocky Mountains." Dr. Denison, be it remembered, ranks "coolness or cold" as the most important attribute of climate next after "dryness."

But when regard for their welfare compels consumptives to abandon Davos by reason of the rise in temperature caused by the approach of spring, were Denison's explanation the correct one they would hardly resort, with the unanimous approval of the best medical advisers, to Badenweiler and Freiberg in the Black Forest, and to other widely esteemed continental resorts where the snow has melted earlier, and where the ground is dry and the air is relatively warm. Then too, there are habitable Alpine heights but little more elevated than Davos, where the summer temperature is cooler far than in the mountain parks of Colorado, and the air is yet more rarefied than at Badenweiler. But these heights are close to the line of perpetual snow and, in summer, to almost perpetual thawing.

In the absence of any systematic meteorological observations continued at Davos throughout the year, whoever will dispassionately consider the facts in the case will, I think, be forced to recognize that, with the melting of the Alpine snows, the erstwhile dry atmosphere of the Swiss Alps must necessarily become relatively charged with moisture.⁶

⁶ The following extracts from the Encyclopædia Britannica, Vol. xvi, p. 125. Stoddart, Phila., will illustrate the point: "Thus the maximum

If, as Dr. Denison teaches—and the doctrine receives my fullest endorsement—if *dryness* be the factor of prime importance in the climatic arrest of phthisis, then the reason why invalids are restored to health at Davos and in Colorado is not because of the altitude and consequent rarefaction of the air, for, as Dr. Meuer well says (op. cit. p. 5), "altitude alone can scarcely be said to enter into the account at all, unless accompanied by the long list of favorable conditions above enumerated"—chief of which is *dryness*. And the reason why consumptives are compelled to abandon Davos on the approach of spring, in spite of the altitude remaining "fairly constant," and notwithstanding they are forced to resort for continued climatic treatment to lower elevations where the temperature is not so cool, is because a damp temperature at high altitudes during the blustery season of the year when raw winds prevail, if of sufficient duration will, in cases of incomplete arrest, almost infallibly occasion a fresh extension of the disease.⁷

There are many elements of uncertainty entering into statistics designed to show the percentage of cases of phthisis actually cured by a given climate. A prolonged period of observation is a condition which may not be eliminated from consideration in such patients as may seem from their general appearance, and even from a careful physical examination, to have fully regained their former health. How long this probationary period must last hardly any two observers are agreed. Many cases too, appear fully recovered only so long as they remain in the more favorable climate; while a return to more humid atmospheres—the only positive test—is very frequently followed by a renewal of the old symptoms of disease. A knowledge of this truth has probably contributed to the paucity of positive statistics. But the fact remains, which may be verified by every traveler, that there are many thousands of former sufferers from pulmonary phthisis contracted in damper climates, who are now living throughout the drier regions shown on Denison's seasonal chart, at altitudes ranging from a few hundred to over 7,000 feet above sea level, and whose ailment is in a state of apparently complete arrest.⁸

If, then, it appears as a logical deduction from established facts that altitude plays no part or, at best, an unessential part in the arrest of the phthisical process; that relative dryness of the atmosphere, and not rarefaction, is the invariable and indispensable climatic factor of all regions of *proved* efficacy, be they elevated or not; there yet remains for consid-

humidity occurs from midnight to 4 A. M., or when the daily temperature is at the minimum; and the minimum humidity at 2 P. M., when the temperature is at the maximum, the curve of humidity being thus inverse to that of temperature." Two exceptions are cited to this law, of which one is as follows: "In the climate of Central Asia in winter, the amount of vapor is very small, and the increase to the relative humidity during the day is probably occasioned by the more active evaporation from the snow during the day." (Italics mine.)

⁷ Dr. C. T. Williams, in discussing the influence of atmospheric humidity on phthisis, well expresses the idea when he says (op. cit. p. 190): "The combination of moisture and heat, as is seen in the sirocco wind, is felt oppressive by most people, but it is doubtful if the combination of cold and moisture be not more harmful."

⁸ It is a pardonable error for a medical man, whose whole vision is focused upon the Swiss Alps and Colorado, while searching for the most prominent climatic factors common to these elevated resorts, to lay the utmost stress upon rarefaction in accounting for their observed climatic benefits. But it would be also pardonable, in accounting for the climatic benefits observed in the dry Rio Grande valley and contiguous territory of West Texas, and in portions of Arizona within 1,000 feet of sea level—benefits apparently quite equal with those obtained in the somewhat cooler and sometimes drier altitudes of New Mexico and Colorado—should a medicine man of the Zuni Indians, with inadequate information as to meteorological conditions, claim that *rattlesnakes* are "the only important factor common to all" the Rocky Mountain slope, and that to their presence chiefly is due the proved efficacy of all this region.

eration the question whether atmospheric rarefaction be a purely negative factor of climate, simply coincident with dryness in several well known mid-continental mountainous regions; or whether it may not, in advanced and in acutely progressing cases at least, be positively detrimental.

There have been several ingenious, not a few labored, and some positively unphysiological explanations offered by medical writers who have witnessed the beneficial influence of *dry* climates at high altitudes in their effort to account for the *assumed* benefits of a rarefied air and of lessened atmospheric pressure to the square inch. But whoever will read the list of general reasons why a given invalid may not go to an otherwise preferable high climate, as enumerated, with evident solicitude and after ample experience, by Dr. Denison, agreeing substantially with other authorities on the subject, will be apt to conclude that while the average patient who is in condition to travel may tolerate an elevation a mile above sea level it is nevertheless, when viewed with expert judgment, to very many a matter of serious hazard. The possible contra-indications are (op. cit. p. 14): 1. The coldest season of the year, *intensifying the effects of altitude too much* (italics mine) for those coming from warmer climes. 2. Advanced age. 3. A very excitable nervous temperament. 4. Some women, for a like susceptibility, and less adaptability to the change and to out-door life than men. 5. Valvular lesions, with rapid action of the heart. Diseases of the great vessels such as aneurisms. 6. Marked and extensive emphysema, pneumothorax and hydro-pneumo thorax. 7. *Active* pneumonia or *existing* hemoptysis. If the pneumonia or hemorrhage is recent the contra-indication amounts to little; if remote, to nothing. If there is reason for some doubt in any such otherwise favorable case, a gradual rise in elevation should be chosen (!). 8. High bodily temperature whether it be rather constant . . . or whether it be regularly vacillating . . . especially with suspicious laryngeal complication, or in so-called "catarrhal phthisis." 9. Extensive involvement of lung tissue in diseased action. 10. The stage of softening, if accompanied by high daily fever, or in one of decided hemorrhagic diathesis. "Quick consumption" with or without intestinal ulceration or albuminuria.

After reading such a list of exceptions from an advocate of high altitudes, who justly ranks among the ablest climatologists of his generation, it would seem indeed, a work of supererogation to argue that elevation *may be* highly detrimental. Indeed, accepting the wisdom of the 7th and 8th contra indications quoted, it would appear superfluous to argue that aerial rarefaction must really be regarded a positive obstacle, and not an aid, in the arrest of the active progress of phthisis. My effort, then, shall be the far more moderate task (after briefly presenting a personal experience as an invalid) of maintaining the doctrine that *when the progress of pulmonary phthisis is arrested at a high altitude, this fortunate result is in no degree attributable to, but occurs in spite of the altitude.*

In the summer of 1890, being thirty-four years of age and engaged in active general practice, I suffered several pulmonary hemorrhages. The last, being quite profuse, was followed by fever and the formation of a small cavity in the apex of the right lung. Bacilli were found to be abundantly present in the

sputa. In October, a month later, I left my home, Mascoutah, Ill., and went to southwest Texas. During the first six months I remained seldom longer than two weeks in any one locality. During this period I traveled over the greater part of Texas lying west of the ninety-eighth degree of west longitude and south of the thirty-second parallel of latitude, and also over portions of southern New Mexico. East of the Pecos river I had regained my normal weight at the end of three and one-half months, but feeling sensitive to every cloudy day, though the sun is obscured in this region as little as in Colorado.* I then went to El Paso for the sake of its still sunnier sky. The air there was ideally dry, and improvement in strength continued with each day. All cough had long since ceased, and expectoration was reduced to a minimum. Early March found me at Marfa, Texas, with an altitude of 4,700 feet. Here, as everywhere in my journeying, were to be found invalids, who, prompted by the benefits received by others, were also seeking a climatic cure. But here one day there was an overcast sky. A gusty March wind was blowing. Prudential considerations kept me within doors all day except for a fifteen minutes stroll, well wrapped up, in the course of the afternoon. Though at lower altitudes and sometimes at a lower temperature, I had, in the course of the preceding five months, repeatedly driven in the open air all day long and without any detriment whatever, I had the misfortune now, though exercising all reasonable precautions, to contract a cold with a tickling cough, due to a recurrence of irritation in the right apex; and during my remaining stay of a week this cough showed no indication of subsiding. Then I returned to El Paso, a thousand feet lower—3,700 feet above the sea. But this locality, which had seemed on my first arrival six weeks previously and which yet seemed, so far as dryness, temperature and sunshine were concerned, to be almost ideal in its winter climate, afforded no relief to the teasing, dry cough. Twelve days of thoughtful solicitude passed, when, after much hesitation, I resolved to return to the considerably less rarefied, if not quite so kiln-dried atmosphere east of the Pecos river and west of the 100th meridian. My train left El Paso in the afternoon, and halted for supper 4,400 feet above sea-level at Sierra Blanca. It is there begins the gentle downward incline towards the Pecos river, which is crossed at Pecos City about midnight. Falling asleep early, though still coughing, I knew nothing more till daybreak. But with the new day the cough was entirely gone.

I have not been able to account for the complete subsidence in one night while traveling down hill at twenty-five miles an hour, of an irritative cough of three weeks duration, on any other hypothesis than that it was due to a diminution of aerial rarefaction—to a restoration of the accustomed atmospheric pressure on a circumscribed acutely congested area of the respiratory tract. Here is the theoretical explanation:

A cupping glass applied to any surface of the body will produce a diminution of superficial air pressure corresponding in degree to the rarefaction of the air remaining within the cup. This diminished air pressure manifests itself by a suction exerted on the surface circumscribed by the rim of the cup. The force of this suction corresponds directly with the

* Denison's Climatic Chart, op. cit.

diminution of the pressure of the surrounding atmosphere by virtue of the rarefied condition of the air within the cup. When the suction exerted by a cupping glass amounts to two and one-half pounds per square inch, then is manifested over one or two inches of surface the tremendous diminution of atmospheric pressure which, over every square inch of both the cutaneous and respiratory surfaces of the human body, is encountered at an elevation of about 5,000 feet above sea-level.

The phenomena consequent upon the application of a cupping glass to the surface of the body are evident and well marked, but they are limited in area. The same phenomena in kind would be produced over the entire surface, both cutaneous and respiratory, by a corresponding diminution of superficial air pressure on the human body, were it enclosed within an air-tight cabinet, the air being partially exhausted therefrom by means of a pump; and are often actually produced by ascending to a sufficient height in a balloon, or, as is more commonly the case, by rapid transit within a few hours or days from low levels to a higher elevation. The most marked phenomena observable on first glance are:

a. A dilatation and engorgement, sometimes to the point of rupture and consequent hemorrhage, of the capillary system of blood vessels, both on the cutaneous and respiratory surfaces.¹⁰ This diminution of superficial air pressure even though of only moderate degree, will at any point of localized irritation from whatever cause, produce congestion, and if long enough continued would necessarily bring about a transudation of serum into the surrounding tissues. This process may be cut short, if it has not gone too far, by a restoration of the accustomed pressure of the air.

b. Quickened and deeper respiration consequent on the diminution of oxygen per cubic foot of rarefied air, this diminution corresponding directly with the degree of rarefaction.

Man possesses wonderful adaptability to his surroundings. But his endurance of heat and cold, of hunger and thirst, of variations in atmospheric pressure ranging from fifteen and one-half pounds per square inch at sea level to less than ten pounds in the Andes and Himalayas, or less than five pounds in balloon ascents as experienced by Glaisher and Coxwell,¹¹—such extremes are endurable only as the individual approximates to a perfect physique and in direct proportion thereto. The fittest alone survive unaccustomed hardships of such character if suddenly imposed and carried to an extreme; the unfit, *i.e.*, physically weak, die.

Fully appreciating how much easier it is to point out the defects of an obviously erroneous theory than to build up another which may successfully withstand criticism, I shall nevertheless venture to formulate two propositions based on the foregoing data: the one explanatory of what I conceive to be the chief detrimental influence of markedly diminished air pressure in cases of unarrested pulmonary phthisis; the other illustrating the antagonism of pro-

nounced atmospheric rarefaction to nature's process of arrest as shown by pathological evidence *post-mortem*:

1. The healthy human body soon adapts itself to high altitudes if not too suddenly reached. But a superficial ulceration, with or without marked infiltration of the adjacent tissues, whether on the mucous surface of the respiratory tract or on the skin, is congested, and its normal tendency to repair is consequently hindered by any considerable diminution of atmospheric pressure. Hence (it must be) those writers who most enthusiastically praise rarefaction as a curative agent (instancing, however, in proof altitudes which are also very dry) nevertheless advise urgently against sending to high altitudes such invalids as are suffering from "*active pneumonia*," "*existing hemoptysis*," or with "*high bodily temperature*," or "*during the stage of softening*." Despite the recognized hazard, however, it is astonishing how many doubtful cases do well if only the air be sufficiently dry and the temperature sufficiently mild. As in morals, charity, so in climate, dryness covers a multitude of sins. For the rapid evaporation by a relatively dry air of the serum transuding from the capillary vessels of a superficial congested area, is nature's best antidote to the irritation (affording a safety-valve, so to speak, to the consequent infiltration) which a considerably diminished air pressure causes, more especially when coincident with an approximate saturation of the air with aqueous vapor; and this saturation occurs occasionally even in the driest regions.¹²

2. In his discussion of the mechanical effects of diminished air pressure Dr. Denison says (*op. cit.* p. 9): "The writer has always maintained and still holds that there is no cause for fear if proper precautions are taken as to extreme elevations and the character and stage of the disease existing in the lungs; *i.e.*, that there should be no spots of softened lung tissue (especially near the root of the lungs), which are insufficiently protected by nature's great conservative process—the deposition of fibrous tissue." A little farther on he adds: "Of course, in very acute conditions we are 'on the other side of the fence,' and it is *rest* which is needed."

It seems hardly necessary to formulate the evident and logical deduction from these statements. If "in very acute conditions it is *rest* which is needed"—and rest is certainly a condition indispensable to the speediest cessation of all inflammatory processes, how can "an increase both in frequency and in the depth of respiration" which "first" occurs "on the arrival of a healthy individual in a high altitude" (*op. cit.* p. 9), a phenomenon which is yet more marked in invalids with diminished breathing capacity—how can such an increase in respiratory activity fail to be detrimental *per se* in its effects on a case of pulmonary phthisis the progress of which has not been previously arrested? And if "the depo-

¹⁰ J. Mitchell Bruce, in discussing the Therapeutics of Air, (*Quain's Dictionary of Medicine*, p. 19) says with reference to air which has been mechanically rarefied: "Inspiration of air which has been rarefied by one two-hundred-and-fortieth to one one-hundred-and-twentieth, . . . of an atmosphere, immediately causes the phenomena of inspiratory dyspnea: the thoracic viscera are congested and hemoptysis may result, for the effect may be regarded as that of dry cupping the pulmonary alveoli."

¹¹ C. Theodore Williams, *op. cit.* p. 193.

¹² The annual mean relative humidity of Denver, compiled from twenty years observations is quoted in the Ninth Annual Report of the Denver Chamber of Commerce (1892) as 49.5 per cent. of saturation. The warmer months—June to October inclusive—show a lower, and the colder months—November to May inclusive—a higher per cent. of saturation than the remarkably low annual mean. We may find in this fact an additional possible explanation of the contra-indication, previously quoted from Dr. Denison, that a given invalid coming from a warmer clime may not resort to an otherwise preferable high altitude during the coldest season of the year. The personal experience of the writer (though from a colder climate) and Dr. C. T. Williams' opinion previously cited, would go to show that not merely cold and altitude, but these two in conjunction with a high per cent. of humidity, even if of only brief duration, form a combination calculated to be specially injurious to consumptives.

sition of fibrous tissue" be "nature's great conservative process," are we justified in limiting its purpose to the protection "of spots of softened lung tissue?" Its invariable deposition in all cases of either partial or complete arrest of tuberculous pulmonary processes, as shown *post mortem*, can receive no other adequate explanation than that it is nature's means of securing rest to the inflamed tissues—of placing them in splints to limit motion. But an increase in the frequency and in the depth of respiration, which immediately occurs to a much more marked degree than is the case with a healthy individual, on the arrival in a high altitude of an invalid with a relatively reduced breathing capacity, directly antagonizes nature's effort, by the deposition of fibrous tissue, to secure rest by limiting motion.

The very ingenious and logical line of argument advanced by Dr. Denison in favor of *cold* and of *rarefaction*,¹³ both of which are produced by elevation, is based chiefly upon his opinion as to the *modus operandi* of dry air, viz.: "In the writer's opinion the chief argument in favor of atmospheric dryness is based upon the *increased transpiration of aqueous vapor from the lungs*, in a degree according with the dryness of the air breathed" (op. cit. p. 3). And since, the percentage of saturation with aqueous vapor (relative humidity) being the same, a given volume of cold air contains a smaller weight of aqueous vapor (*i.e.*, a lower *absolute* humidity) than the same volume of warm air; and since, under the same condition, rarefied air will contain a smaller weight of moisture than an equal volume at sea level, it therefore follows, *the percentage of saturation being the same*, that cool and rarefied air contains less absolute humidity than warmer air at lower levels. And the breathing of such an atmosphere must consequently increase the transpiration of aqueous vapor from the lungs more than is possible at higher temperatures and lower altitudes.

The fact that on Pike's Peak, at least in the winter of 1883, the air was both relatively and absolutely more humid than at Denver (Denison's seasonal chart, op. cit.), despite its greater elevation by some 9,000 feet, and notwithstanding its lower temperature, demonstrates that the "tendency of mountain climates towards dryness rather than humidity" may be overcome by counter-tendencies, the causes for which it is not necessary here to discuss. So that were the premise beyond criticism from which is drawn the conclusion just cited, in favor of high altitudes by reason of the low absolute humidity claimed for them, the deduction would not apply in favor of *all* high altitudes, but only in favor of those elevations where the relative humidity is likewise low.

But the premise quoted, from which so important conclusions are drawn, is offered only as "the writer's opinion." As such it is entitled to the very highest consideration. There is indeed no writer on climate known to me whose unsupported opinion is entitled to more weight. But in the absence of demonstration its correctness may (and should) be questioned, if the conclusions logically deducible therefrom fail to harmonize with facts which are well established.

Let us hazard *per contra* the different opinion, which alone appears to harmonize with all the known facts

bearing on the subject, viz.: that the acknowledged influence of a dry air in retarding or arresting the progress of pulmonary phthisis be due *not* to "*the increased transpiration of aqueous vapor from the lungs*," which is but the exaggeration of a vital, physiological function pertaining to normal physiological lung tissue; but that, on the contrary, it be due to the rapid evaporation of serum transuding from the engorged superficial capillaries of certain circumscribed inflamed pulmonary areas, which rapid evaporation promotes nature's tendency toward limiting the infiltration and the extent of inflamed pulmonary tissue. And whether such circumscribed engorgement were, or were not, exaggerated by the reduced air pressure of a high altitude—equivalent on first arrival to the mechanical effects of a cupping glass—its diminution by reason of the absorbent properties of dry air could certainly not be regarded as a vital physiological process affecting chiefly the normal pulmonary tissue, but as *a process due to physical laws only—one affecting pathological tissue alone*.

And, upon such a supposition as a premise, it would follow that "*coolness or cold* as opposed to *warmth or heat*," and also "*rarefaction* as opposed to *sea level pressure*," would both be eliminated from the list of important climatic attributes, in so far at least as they affect the prime factor, *dryness*; for it is a fact thoroughly well established that the absorbent power of the air bears no definite relation to its *absolute* humidity, in the calculation of which cold and rarefaction are always important factors; but does correspond definitely with its *relative* humidity, taken in connection with the temperature, wind movement and the approximation to sea level. For, *as temperature rises the evaporative power of an atmosphere, whose per cent. of saturation remains the same, increases faster than the rise in the thermometer*.¹⁴ And it is equally true that, *relative humidity and temperature remaining the same, a given volume of air will absorb and retain more moisture at or near sea level than at any higher elevation whatever*, its capacity to absorb moisture diminishing as the altitude increases. Evaporation is also far more rapid during a brisk breeze than in calm weather, so that Parkes has laid down the rational proposition, acted upon by every weather bureau throughout the world, that "the degree of dryness or moisture of an atmosphere should be expressed in terms of relative (and not of absolute) humidity, and should always be taken in connection with the temperature, movement and density of the air, if this latter varies much from sea level."¹⁵

The important deduction then logically follows that though a particular high altitude resort may have a low relative humidity, yet the low mean temperature and the atmospheric rarefaction may be such as to afford a smaller capacity for the absorption of moisture by a given volume of air than would be the case in some locality nearer to sea level with a relative humidity somewhat greater, but with a mean temperature considerably higher and with a total annual wind movement materially greater.

It is not within the scope of this paper to institute comparisons between particular health resorts at high and low altitudes, or to make extended reference to any other factors of climate than those which are so intimately associated with the growing

¹³ W. Steffen of Davos, seems to have argued along the same line; his argument being combated by Dr. Julius Hann (v. Davos, in Buck's Reference Handbook of the Medical Sciences), but I have failed to secure his original discussion of the subject.

¹⁴ E. A. Parkes, op. cit. Vol. ii, p. 88.

¹⁵ Op. cit. Vol. ii, p. 87.

professional advocacy of high altitudes that they may not be eliminated from consideration. But it may not be out of place after presenting a table summarizing the weather observations at three tolerably widely separated localities of moderate altitude in West Texas to offer a few closing suggestions. Let it be remarked, however, in passing, that while it would appear from the evidence already cited that there is no causative relation whatever between absolute elevation and relative humidity, it can not be doubted that the air is relatively drier on a hill top than in the adjacent valley, or even very frequently on a house top than at the level of the ground. This difference is greater in proportion to the rain or snowfall, and to the impermeability or bad drainage of the soil. And since men occupy habitations located on the soil, and usually at a much lower level than the psychrometers of the Weather Bureau, there is a wide range for selection of a place of dwelling in a given region, and also of regions with no other apparent climatic differences than those pertaining to the rainfall, drainage and porosity of the soil.¹⁶

Altitude, Mean Temperature, Wind Movement, Precipitation, Cloudiness and Mean Relative Humidity of stations named, from 1877, for 10 years, 6 years and 8 years respectively.

Months.	San Antonio. Elevation 781 feet.				Eagle Pass. Elevation 800 feet.				San Angelo. Elevation 1,900 feet						
	Mean temperature.	Wind movement, miles per month.	Precipitation in inches.	Cloudiness in tenths.	Mean relative humidity	Mean temperature.	Wind movement, miles per month.	Precipitation in inches.	Cloudiness in tenths.	Mean relative humidity	Mean temperature.	Wind movement, miles per month.	Precipitation in inches.	Cloudiness in tenths.	Mean relative humidity
October.	70	4,276	1.99	4.3	69	72	3,717	1.92	3.8	64	65	5,986	2.99	4.4	69
Nov. . . .	58	4,587	2.09	5.0	68	59	3,859	0.60	3.7	64	51	5,698	1.31	3.9	66
Dec. . . .	53	4,832	1.76	4.9	70	53	3,316	1.19	4.6	67	45	6,442	1.47	3.9	65
Jan. . . .	50	4,960	1.58	5.1	72	52	3,427	1.12	4.6	66	42	6,206	1.14	4.0	68
Feb. . . .	56	4,803	2.13	5.4	68	57	4,199	0.97	4.2	61	48	6,385	1.14	4.1	64
March.	63	5,199	2.22	5.3	67	65	4,256	1.80	4.2	63	57	7,231	1.30	3.8	60
Six cooler months.	58	28,657	11.77	5.0	69	60	22,774	7.60	4.2	64	51	37,948	9.35	4.0	65
April. . . .	69	5,149	3.38	5.2	65	73	4,922	1.21	3.6	53	65	7,959	2.22	3.3	58
May. . . .	75	4,820	3.31	5.3	70	80	5,518	3.61	4.2	63	72	8,024	4.32	4.2	63
June	81	4,582	2.70	4.5	69	86	5,431	2.13	3.2	59	80	7,253	2.67	3.6	60
July	83	4,172	2.84	4.5	64	87	5,075	3.34	2.8	55	82	6,565	3.47	3.4	58
August . . .	82	3,767	3.65	4.4	67	84	4,025	3.54	3.5	62	80	5,527	3.55	3.7	62
Sept. . . .	77	4,000	4.21	4.4	70	79	3,550	3.54	3.0	64	73	5,607	4.36	4.3	66
Six warmer months.	78	26,490	20.09	4.7	67	82	28,521	17.37	3.4	59	75	40,935	20.59	3.7	61
Annual . . .	68	55,147	31.86	4.8	68	71	51,295	24.97	3.8	62	63	78,883	29.94	3.8	63

It is apparent from the foregoing table that of the three localities cited the first two are preferable in winter so far as temperature is concerned; but temperature being equal the drier the air the better. In summer on the other hand, the greater wind movement and lower temperature of the third locality commend it to favor. The writer does not wish to appear as instituting invidious comparisons between health resorts; for the climate of no city or town is comparable with that of the open country. It is living and, if possible, sleeping out of doors in the suitable climate which is of the greatest benefit. It is

¹⁶ The acting chief of the Weather Bureau in his letter transmitting the data above given says: "The averages given are for rather short periods and are therefore not to be considered as having a fixed or permanent value." As showing the correctness of this view with reference at least to one of the localities mentioned, I will only cite the report of the chief signal officer for 1885, Part I, p. 181, giving the mean annual relative humidity at Ft. Concho (San Angelo) from Sept. 1872 to Nov. 1879 as being only 54 per cent, while the table shows 63 for a different period. The precipitation for fifteen years and one month at the same place averaged but 19.02 inches annually according to signal service report, 1888, on the rainfall of the Pacific Slope and Western States and Territories.

not "roughing it," nor exposure to inclement weather and to hardships, which indeed must be carefully avoided by the invalid; but living every hour possible in the open air and avoiding so much exercise as to cause fatigue—this it is which enables the climate to arrest the disease and with equal step, build up the patient's strength again. Sleeping on a cot on the open veranda of a country house is specially to be commended. The invalid who has tried it for one summer will know how to appreciate its manifold charms and benefits.

Finally, when every symptom of disease has subsided and the strength is fully restored, it should be thoroughly understood by every patient that his return to a humid climate will, not *possibly* but *probably* rekindle the processes of disease with far more than their former intensity and far less power of resistance on the part of the patient. I am cognizant of too many instances of this kind not to lay special emphasis upon so important a matter. The restored patient if his condition had ever been a grave one must make up his mind philosophically to reside in a dry climate. But he may then if so disposed, remove to a higher altitude, always with an eye single to a dry atmosphere as the *sine qua non*. For it is the writer's opinion, which he contemplates personally acting upon, that when the progress of phthisis has been completely arrested the restored invalid may then with impunity remove to any very dry altitude, provided only the temperature in winter be not disagreeably low, and the air pressure be not so reduced as to excite fresh irritation in an incompletely healed focus of disease.

MENTAL ABERRATIONS.

Read before the Southeast Missouri Medical Association, at Charleston, Mo., May, 1893.

BY L. T. HALL, M.D.

FIRST ASSISTANT, ASYLUM NO. 1, FULTON, MO.

"Man is a harp, whose chords elude the sight,
Each yielding harmony, disposed aright.
The screws reversed (a task which, if He please,
God in a moment executes with ease),
Ten thousand thousand strings at once go loose,
Lost, till He tune them, all their power and use."

Thus sadly sang one of England's purest, sweetest poets, out of the sad experience of a life of alternating light and shadow, harmony and discord, now reveling in the loftiest flights of poetic fancy, the profoundest depths of spiritual experience; now groveling in darkness and despair, driving him to a desperate attempt upon his life, the light of reason finally going out behind a cloud of impenetrable gloom. This comparison of the human mind to a musical instrument, susceptible when attuned aright of the most delightful harmonies, but with "the screws reversed" its "ten thousand thousand strings" giving forth only jarring discord, is as strikingly true as poetical; of which Cowper's own life and experience furnish the most striking illustration. Between the extremes presented in this beautiful figure we find degrees of almost infinite gradation; sometimes but a single chord of the "ten thousand thousand strings" seeming to have lost its tension, the harmony of the instrument remaining unbroken to the most skillful observation until the pressure of some untoward objective circumstance or subjective condition sends forth the discordant vibrations of the string "gone loose."