Schreiben des Herrn *Henderson*, Directors der Sternwarte in Edinburg.

Observatory Edinburgh, April 11, 1843.

On the Comet of 1668.

There appears to be great probability in favour of the supposition that the late Comet and the one which appeared in 1668 are the same. I have in my possession an engraved Map representing the apparent track of the latter Comet among the stars from the 9th to the 21st. of March as observed at Goa in India. The title of the Map is „Observationes Goæ habitæ circa Phænomenum Cœleste, quod apparuit Mense Martio Anno 1668 Romam missæ ad P. Aegidium Franciscum de Gottignies in Collegio Romano Matheseos Professorem.“ I think it probable that it formed part of the work entitled „Aegidius Franc. de Gottignies de figuris Cometarum qui an. 1664, 1665, 1668 apparuerunt cum animadversionibus & tabulis in æs incis. Romæ 1668,“ mentioned by *Weidler* (*Historia Astronomiæ* p. 508) and *La Lande* (*Bibliographie Astronomiæ* p. 271). I have not seen this work, but it will probably be found in some Library, and as it may contain important information about the course of the Comet it should be enquired for. The Map is rather a rough one, as I find on examining the relative positions of the stars which are laid down, errors amounting sometimes to a degree. The comet is represented by straight lines, to each of which is affixed the day of the month, when it was observed in the position indicated. The dates extend from the 5th to the 21st. of March; but to the 8th inclusive the western terminations of the lines are cut off by a line marked „Horizon,“ evidently intimating that on these days the body of the comet set before being visible. I assume that the western extremities of the lines indicate the observed positions of the nucleus. From the 18th to the 21st. inclusive portions of the lines towards both extremities are represented by dots; which I suppose are intended to denote that those portions of the Tail had become very faint and that the positions assigned to both extremities are doubtful. For this reason the positions of the supposed nucleus in the Map must be very uncertain for those days. The New Moon was on the 12th of March, and there is therefore reason for believing that from the increasing Moonlight, and the rapidly diminishing light of the Comet, the body was invisible after the 17th, and it was only the brighter part of the tail which could be seen.

On comparing the positions assigned in the Map to the supposed nucleus with those of the neighbouring stars reduced to 1668, I find for the Comet's observed places as follows:

Date New style 1668.	Right Ascens.	Daily differ.	Declination.	Daily differ.
March 9.	18° 49'	2° 2'	—10° 50'	+43'
10.	20 51	2 17	10 7	+44
11.	23 8	2 2	9 23	+24
12.	25 10	2 43	8 59	+22
13.	27 53	2 29	8 37	+14
14.	30 22	2 30	8 23	—16
15.	32 52	1 53	8 39	+17
16.	34 45	3 12	8 22	+7
17.	37 57	3 29	8 15	—24
18.	41 26	3 42	8 39	—10
19.	45 8	3 26	8 49	—1
20.	48 34	3 34	8 50	+9
21.	52 8		8 41	

The times of observation may be assumed to be 7^h or 7^h 30^m at Goa or 2^h 5^m or 2^h 35^m at Greenwich.

From the columns of daily differences it is evident that the places of the Comet must be affected with considerable errors; and those of the 18—21 appear to be too great in Right Ascension and too southerly in declination, which may be explained by the cause already mentioned.

Last year I made several attempts to determine an orbit which would represent these observations; and not paying attention to the doubtful character of the latter observed places, I employed the whole indiscriminately in the investigation. The apparent path of the Comet is nearly in that situation with regard to the Sun, when the problem of determining the orbit is of an indeterminate nature, and when errors of observation have an enormous influence on the results. The orbit which I obtained had the following elements; and I communicated them at the time to Professor *Encke*.

Time of perihelion passage Feb. 24^d 782 Greenwich time.

log q 9.39990

π 40° 9

Motion direct.

Ω 193 26

i 27 7

The computed places in this orbit are as follows

Date 1668.	Right Ascens.	Obs. R. A. = comp.	Declination.	Obs. D. = comp.
March 9.	18° 21	+28'	—9° 37'	—73'
10.	20 37	+14	9 41	—26
11.	22 59	+9	9 44	+21
12.	25 26	—16	9 45	+46
13.	27 58	—5	9 44	+67
14.	30 36	—14	9 42	+79
15.	33 19	—27	9 37	+58

Date 1668.	Right Ascens.	Obs. R. A. = comp.	Declina- tion.	Obs. D. = comp.
March 16.	36° 7'	-82'	-9° 30'	+68'
17.	39 0	-63	9 21	+66
18.	41 58	-32	9 10	+31
19.	45 2	+6	8 56	+7
20.	48 10	+24	8 40	-10
21.	51 24	+44	8 22	-19

When I heard of the appearance of the late Comet, I was struck by the similarity between it and this one of 1668; but I had no accurate intelligence of the observed positions of the former till I received Professor *Schumacher's* first Circular, and the Comet was not seen in Edinburgh. From the information contained in the Circular the probability that the Comets were the same appeared to have considerable strength. I was engaged in an attempt to reconcile the observations of the one first in date with the orbit of the latter, when I received the second Circular, and following in the steps there pointed out, I have assumed the following Elements in 1668 taken (excepting the time) from some of the recent observations.

Time of perihelion passage 1668 February 28^d 8 Greenw. time.

$\log q$ 7,68000

π 277° 2'

Ω 357 17

i 35 58

Motion retrograde.

And from them I computed the following places:

Date 1668.	Right Ascens.	Obs. R. A. = comp.	Declina- tion.	Obs. D. = comp.
March 9.	16° 46'	+2° 3'	-10° 16'	-34'
10.	19 59	+52	10 10	+3
11.	23 1	+7	10 2	+39

Date 1668.	Right Ascens.	Obs. R. A. = comp.	Declina- tion.	Obs. D. = comp.
March 12.	25° 53'	-43'	-9° 52'	+53'
13.	28 36	-43	9 42	+1° 5
14.	31 11	-49	9 28	+1 5
15.	33 38	-46	9 15	+36
16.	35 58	-1 13	9 1	+1 39
17.	38 10	-13	8 46	+31
18.	40 15	+1 11	8 30	-9
19.	42 12	+2 56	8 14	-35
20.	44 3	+4 31	7 57	-53
21.	45 47	+6 21	7 40	-1 1

The differences between the observed and computed places from the 9th to the 17th inclusive may perhaps be diminished by a better determination of observed places from the Map, and a better assumption of elements. The great differences from the 18th to the 21st are evidently owing to the uncertainty of the observations; and they may be accounted for on the supposition that the nucleus was then invisible.

This comparison and the great similarity in the appearances of the Comets seem to render it very probable that the Comets are the same. However the agreement within tolerable limits of the observed and computed places in 1668 is not sufficient of itself to establish the identity; as orbits widely different may be found to represent the rough observations made at that period.

I will send the Map to the Royal Astronomical Society of London to be preserved in their Library.

T. Henderson.

Verkäufliche Instrumente.

- 1) Ein Cometensucher von *Fraunhofer*, 34 Pariser Linien Oeffnung, mit 10 und 15maliger Vergrößerung, und einer Vorrichtung zum Anschrauben stärkerer Oculare. Das Objectiv verträgt noch sehr gut 100malige Vergrößerung. Das Instrument ist so sorgfältig erhalten, daß es als ganz neu betrachtet werden kann. Preis 50 Rthl. Pr. Crt.
- 2) Ein Cometensucher mit achromatischem Objectiv von 37 Par. Linien Oeffnung, Brennweite 2½ Fufs, Vergrößerung 12mal, und auf dem Fußboden stehenden Stative. Pr. 50 Rthl. Pr. Crt.

Das Objectiv dieses Instruments ist nicht von *Fraunhofer*, giebt aber an Lichtstärke, Präcision und Farbenlosigkeit dem *Fraunhoferschen* nichts nach, und hat noch den Vorzug, daß noch am Rande des etwa 5° fassenden Gesichtsfeldes die Bilder fast dieselbe Präcision, wie in der Mitte, haben.

Man kann sich wegen dieser Instrumente an den Herrn Superintendenten *Hülsmann* in Elberfeld wenden.

Inhalt.

Observations et éléments de la seconde Comète de l'année 1840. Par Mr. *Plantamour*. (Beschluss.) p. 321.

Schreiben des Herrn *Henderson*, Directors der Sternwarte in Edinburgh. p. 333.

Verkäufliche Instrumente. p. 335.