ANNALS OF OTOLOGY, RHINOLOGY AND LARYNGOLOGY

INCORPORATING THE INDEX OF OTOLARYNGOLOGY.

VOL. XXX.

JUNE, 1921.

No. 2.

SOME POINTS IN THE ANATOMY OF THE HUMAN TEMPORAL BONE.

By JOSEPH GOLDSTEIN, M. D.,

DEPARTMENT OF ANATOMY OF THE STANFORD UNIVERSITY
MEDICAL SCHOOL,

STANFORD UNIVERSITY, CALIF.

I.—THE WEIGHT AND DENSITY OF THE TEMPORAL BONE.

The weight of the bones of the skull in both sexes is greatest from twenty to forty-five, according to Gurriere and Massetti, 1895. Hence the maximum weight of 63.5 grams found in No. 119, among one hundred cleaned, dried temporal bones, probably approaches the maximum weight of the dry disarticulated adult bone quite closely. The lightest adult bone, No. 133 of this series, weighed but 22.5 grams, giving a difference of 41 grams.

Specimen 119 displaced 43 grams of water, its density hence being 1.48, but specimen 133 displaced but 10.5 grams of water, indicating a density of 2.14. Hence the lighter bone apparently had a far greater density. But the weight and density of bones, even when dried by years of exposure in a dry atmosphere, depends upon a great number of factors. One

must consider the age and possibly racial characteristics of the individual bone and especially the method and the extent of the cleaning. Hence for this and other reasons the results obtained from these two specimens probably are only roughly indicative of the true density.

The average weight of fifty intact right temporal bones in Table I, regardless of sex and including the ear ossicles and the styloid and zygomatic processes to the point of articulation of the latter, in the dried state is 41.5 grams. Fifty left temporal bones average 41.6 grams. Hence asymmetry in form, even though present to a marked degree as described by Cheatle in 1913, in certain skulls, does not give one side a preponderance in average weight, even if one takes only a relatively small number of temporal bones.

TABLE I.

Right Side		Left Side	
No. of Specimen	Weight	No. of Specimen	Weight
1	39 grs.	7	52 grs.
44	47 "	15	29 "
10	48 "	18	31 "
16	52 "	44	47 "
19	27 "	45	47 "
22	46 "	55	47 "
24	57 "	60	38 "
25	51 "	63	34 "
27	33 "	71	43 "
32	53 "	75	51 "
38	34 "	76	43 "
40	32 "	79	49 "
49	49 "	80	53 "
51	36 "	85	37 "
52	30 "	92	43 "
58	61 "	94	31 "
59	48 "	109	35 "
64	30 "	116	37 "
65	39 "	119	63 "
66	43 "	120	48 "
68	38 "	121	47 "

TABLE I.—Continued.

Right Side		Left Side	
No. of Specimen	Weight	No. of Specimen	Weight
69	52 "	130	52 "
70	62 "	145	50 "
105	47 "	146	47 "
106	32 "	148	42.5 "
107	37 "	153	38 "
108	30 "	156	25 "
114	52 "	157	36 "
115	39 "	163	27 "
122	35 "	166	31 "
124	27.5 "	167	36.5 "
125	37 "	168	36.5 "
131	36 "	323	43 "
133	22 "	325	38 "
134	31 "	327	38 "
142	46 "	328	34 "
147	57 "	333	38 "
152	35.4 "	341	47 "
154	38 "	345	33 "
164	37 "	347	36 "
322	39 "	349	40 "
326	32 "	352	49 "
329	50 "	353	40 "
331	52 "	357	44 "
335	40 "	359	30 "
340	50 "	361	48 "
342	48 "	364	36 "
343	40 "	369	40 "
344	41 "	372	55 "
348	32 "	373	59 "
Average wt., 41.5 "		Average wt., 41.6 "	

II.—TORSION OF THE PETROUS PORTION.

As is well known, the temporal bone represents three elements—the petrosal, squamous and tympanic elements—which usually can be separated at birth. The external acoustic

canal is largely unformed. At this time the tympanic ring is circular in outline and the tympanic membrane directly continuous and lies in the same plane as the external surface of the squamosa. Canal formation begins by the rapid growth of the anterior and posterior margins of the tympanic element, which are much thicker than the rest of the tympanic portion of the canal at this time. In adult bones, as we shall see later in the discussion of the osseous external acoustic orifice, we sometimes find a thickening at these very points, although this border usually becomes very thin in the adult condition.

At birth the petrous portion seems to show no torsion. In cross section it assumes the shape of an isosceles triangle—the anterior superior and the posterior superior surfaces forming straight sides. The inferior surface is quite irregular due to its articular surfaces, to the numerous foraminæ and the jugular fossa. The superior angle of the triangle is slightly rounded.

In the adult bone certain changes have occurred in the contour of the petrous portion. The medial extremity which articulates with the basilar portion of the sphenoid is directed anteriorly. This gives the superior border of the petrous bone a slightly bent direction, because it remains in juxtaposition with the sphenoid. But the superior margin, viewed from a median plane, also is rotated, anteriorally or clockwise, looking medially. The change that has taken place during this period of torsion has modified somewhat the structures underlying and contained within this portion of the bone, which forms the superior surface of the external auditory canal and envelops a part of the tympanum. The torsion which modifies the petrous portion modifies the form of the canal, as we shall see also, that of its external orifice.

In calculating the amount of rotation that the medial portion of the petrous element makes, I used the styloid process to fix the bone. It is impossible to determine the angle that the petrous portion makes with the styloid process in the newborn, on account of the absence of the process, but in the adult bones this process, when intact, is usually very definite and constant in direction.

The method used in calculating the angle of torsion is as follows: The bone was fixed by placing the styloid process

in a vertical position in a board with vertical holes graded as to size. The processes which are of variable size were fitted very snugly into suitable holes. In this manner, one can eliminate error to a great degree and can be relatively sure that all angles are measured from a fairly definite base line. The surface of the board is at right angles to the process, and the superior margin of the petrous portion is adjusted parallel to the edge of the board. By using the plane surface of the board as a base line, one can measure the angle of rotation of the petrous portion quite accurately with a protractor. In order to reduce error to a minimum, the angle of rotation was measured on a line passing through the center of the orifice of the internal auditory canal. This surface of the bone sometimes is irregular, due to a ridge just over this orifice. In these cases the arm of the protractor was placed as nearly parallel to the entire surface as possible. Such a procedure was rarely necessary, however, for usually the arm of the protractor could be rested on the side of the petrous bone, placed parallel to the surface with little difficulty and the angle read.

TABLE II.

Specimen No.	Angle of Rota'n	Specimen No.	Angle of Rota'n
3	118°	122	117°
7	113°	126	105°
16	117°	127	118°
17	113°	131	116°
20	118°	134	110°
22	116°	139	115°
24	122°	145	119°
28	116°	149	115°
31	114°	154	113°
35	125°	159	114
46	115°	163	113°
59	115°	166	120°
65	113°	170	114°
66	112°	322	115°
67	116°	343	112°
75	117°	345	115°
76	115°	346	113°

TABLE II.—Continued.

Specimen No.	Angle of Rota'n	Specimen No.	Angle of Rota'n
78	122°	354	121°
82	115°	355	117°
89	112°	359	113°
93	121°	365	114°
117	118°	369	116°
118	120°	374	114°
120	113°	1169	112°
121	113°	1350	120°

It was interesting to find that there is a correlation between the angle of rotation of the petrous portion, measured as above, and the direction of the long diameter of the osseous external auditory orifice. When the long axis of the external orifice approaches a vertical plane—that is, when it is almost parallel to the styloid process, the anterior superior surface as measured above makes an angle of approximately 120° with the plane surface of the board. In specimens with the long axis approaching a horizontal position, the anterior superior surface makes an angle of approximately 110° with the plane surface of the board. In by far the commonest form, the long axis lies between the two extremes, the angle being about 115°. An average of the fifty specimens in Table II, calculated by this method, gave an angle of 115.7°. The maximum amount of rotation as measured above was 20°, which is the difference between 105° in specimen No. 126 and 125° in specimen No. 24. As the accompanying photographs show, one can determine the amount of rotation of the petrous portion with considerable accuracy by noticing the direction of the long diameter of the osseous external acoustic orifice. Furthermore, if the styloid processes of any series are placed parallel and opposite, the lines indicating the direction of the long diameter of the external orifice will be parallel.

Specimen 149, Fig. 1, the opposite surface of which is represented in Fig. 2, illustrates the usual condition of rotation. In specimen No. 126, represented in Figs. 3 and 4, the direction of the string or the long diameter of the external acoustic orifice is almost parallel to the styloid process, thus repre-

senting a totally different angle of rotation. In specimen No. 24, represented in Figs. 5 and 6, on the other hand, the direction of the long diameter of the external acoustic foramen forms almost a right angle with the styloid process.

There are numerous factors to be considered as to the cause of this rotation, which is always in the same direction—the superior margin being directed anteriorly, but it occurs after birth. At birth and shortly afterwards, no torsion was observed in looking over 125 fetal and infantile specimens. It will be recalled that the basilar portions of the occipital bone also is more vertical and only slightly grooved at this period, the deep hollowing found in adult bones apparently being coincidental with the torsion of the petrous element.

By the sixth month of fetal life, the skull, though smaller, is in much the same condition as at birth, except that the occipital region is relatively larger. The most striking differences are the insignificance of the facial portion and the flatness of the base. In the cranium the frontal region is relatively small. According to Dwight, '07, Merkel divided the growth of the head into two periods, with an intervening one of rest. The first period ends with the seventh year and is followed by inactivity till puberty, when the second period begins. Our interest lies in Merkel's first two stages of the first period, which are said to end at the fifth year. In the first stage, reaching to the end of the first year, the growth is general, but the face gains on the cranium. At six months the basilar process rises more sharply, which, with the downward growth of the face, may have an important effect on the torsion of the temporal bone, since the greatest degree of rotation occurs at this time. In the second stage, to the end of the fifth year, the vault grows more than the base, assuming a more rounded and finished appearance. The face still gains relatively, but grows more in breadth than in height. It is quite probable from my observation that with the termination of this period the maximum degree of torsion has been attained.

As stated above, the torsion begins after birth and apparently continues quite rapidly, for as far as I could observe, in specimens a few years old, the maximum amount of torsion

already had been reached. Hence torsion occurs during infancy at a time the brain grows most rapidly. We have a rapidly growing brain and growing bones which envelop it. We also know that the brain molds the skull to a certain extent and that the skull also modifies the shape of the brain. Since these two conditions appear at the same period in development, the brain factor may have a great deal to do with these modifications of the petrous element at this time. It is possible that the cerebellum and the medulla oblongata exert a pressure by their rapid growth, forcing the basilar portion of the sphenoid and the superior margin of the petrous portion of the temporal bone anteriorly, thus accelerating the growth of the anterior portion of the skull which responds to the marked development of the frontal lobes.

The rapid growth of the cerebrum after birth also may exert a pull upon the dura at its close attachment to the superior margin of the petrous portion of the temporal bone, thus deflecting the medial extremity forward. Not only is the direction of this portion changed during this period, but the superior margin is molded somewhat, for this border is sharper in the adult form than it is in the infantile condition.

III.—THE ANGLE BETWEEN THE PETROSA AND SQUAMOSA.

The angle that the superior margin of the petrous portion makes with a tangent parallel in the main, with the squamosa also, is greater in the adult condition than it is in the newborn. The squamous portion of the temporal bone has the general direction of the lateral surface of the calvarium—the squamosa itself occupying about one-half of the lateral surface in length. The suture between the parietal and temporal bone immediately over the mastoid process is generally very low, and one can readily measure the angle that the petrous element makes with the squamosal by means of a protractor, upon disarticulated bones. Using the external surface of the squamosa immediately above the posterior root of the process as a base line for one arm of the protractor, the angle which the petrous element makes with this arm can be measured directly.

About the time of birth there apparently is a very slight variation in the angle. It varied from 66 to 70 degrees in a

small series of 20 infantile and newly born temporal bones. In the adult condition there is a greater variation, however, for an angle of 52 degrees was found as a minimum and one of 64 degrees as a maximum. The average of the angles between the petrous and squamous portions in 50 specimens in Table III is 57.2 degrees.

TABLE III.

No. of Specimen	Angle	No. of Specimen	Angle
1	59	35	56
2	53	39	53
3	55	40	60
4	60	42	58
6	60	43	58
7	52	44	60
9	55	45	60
14	54	46	64
15	58	47	60
16	58	48	56
17	55	49	60
18	60	51	58
20	55	53	58
21	58	56	60
22	59	58	60
23	58	61	55
24	56	71	60
25	53	152	55
26	60	159	55
27	60	321	62
28	60	326	61
29	60	341	56
30	60	374	57
31	57	1114	55
34	56	1168	58

IV.—VARIATIONS IN THE MASTOID FORAMEN.

The mastoid foramen, when present, has a greater uniformity in position than one infers from a textbook description. Piersol, 1907, gives a typical description in saying that a

small canal, the mastoid foramen, transmitting a vein, runs from the sinus to the outside of the bone, and sometimes reaches as far back as the suture between the temporal and the occipital. Cheatle, 1906, described the vein as leaving the lateral sinus in its descending part and usually running upwards, backwards and outwards with a curved course, to emerge at the posterior edge of the base of the mastoid process. I believe it would be more nearly correct to limit the area of emergence, for after looking over more than 250 adult temporal bones I find that it usually emerges directly over the digastric groove, on a level with the upper margin of the osseus external acoustic orifice. When the lambdoidal suture at its lower extremity forms a convex surface on the mastoid portion of the temporal bone, the foramen is found slightly medial to the digastric groove. If the lambdoidal suture is straight at this region, which is the more common form, the foramen may even appear in the suture, or within a centimeter of the suture, thus appearing directly above the digastric groove. It would appear from this that the suture between the temporal and the occipital bones had a great deal to do with determining the position of the foramen.* One very rarely finds the foramen lateral to the digastric groove.

By tabulating the occurrence of the common forms of canal, we find some interesting data, as judged by 250 specimens in Table 4, chosen at random from our collection. In 68 specimens, or 27 per cent of the bones, there was no indication of a canal from the lateral sinus to the outside of the bone. One can always find emissary foraminæ, variable in number and size, over the mastoid process and at its base, but I do not include any canals that cannot be traced to the lateral sinus. It also is relatively common to find a canal apparently for a vein, entering and leaving on the lateral surface of the base of the process without penetrating deeply into the bone.

In 167 specimens, or 67 per cent of the 250 bones, the canal is single and usually can be traced with very little difficulty from the sinus to the outside of the bone. It is common to find one opening of the canal on the inner table, which divides within the spongiosa and emerges by two openings. This was observed in 11 specimens, or 4 per cent. Four specimens had two separate canals traceable from the sinus to the outside.

In one case the canals were fully two centimeters apart, and both appeared above the digastric groove, being due to two separate veins leaving the lateral sinus.

When the foramen divides within the diplœ to emerge as two canals, one finds them emerging almost parallel. In a few cases, the canal is directed straight out from the lateral sinus, and the distance between the lateral sinus and the external surface of the bone is no greater than the thickness of the bone.

Cheatle, 1906, in discussing the size of the canal described it as being as "big as a pencil" in some cases, and so small that it would not admit a pin in others. Myer, 1913, reported an abnormally large unilateral foramen mastoideum with a diameter of 6 mm. on the inner surface, and a sulcus emissarium on the exterior with an oval mouth with a long diameter of 8 mm. In the 1500 adult temporal bones examined I found none with an abnormally large canal, thus indicating that large canals are very uncommon.

TABLE IV.

	No.	Per Cent.
Absence of emissary canal	68	27
Single emissary canal.....	167	67
Bifurcating emissary canals.....	11	4.4
Double canals.....	4	1.6
Total.....	250	100

V.—THE PROCESSES IN THE MASTOID REGION.

The pars mastoidea, which forms the posterior part of the temporal bone, is continued below into a conical projection, the mastoid process, the size and form of which varies somewhat and which is said to be larger in the male than in the female. The mastoid process is relatively small at birth and contains no air cells except at the antrum. It becomes distinct about the first year, coincident with the obliteration of the petrosquamous suture. The suture is frequently found not wholly fused in the adult temporal bone. Toward puberty

the process becomes pneumatic. This process, which serves for the attachment of the sternocleidomastoideus, splenius capitis and the longissimus capitis muscles, is sometimes divided by a deep groove, the digastric, for the attachment of the posterior belly of the digastricus. Just medial to this groove is a small shallow furrow which lodges the occipital artery.

So much diversity of opinion exists among authorities as to the correct names of the processes of the mastoid region that one is hardly safe in describing this region without first referring to matters of terminology. Gray, 1905 edition, makes no reference to paramastoid or paroccipital processes in man, but in the edition revised by Lewis, 1913, the paramastoid is described in connection with the occipital bone, as an eminence which sometimes projects downwards from the rough under surface of the jugular process, where the rectus capitis lateralis muscle and the lateral atlantooccipital ligament are attached. It is stated that this eminence may be of sufficient length to reach and articulate with the transverse process of the atlas. The last edition of Gray does not mention the par-occipital process, and neither Piersol, 1907, nor Allen, 1884, describe either process. Morris, 1899, says that a process of bone projects rarely from the under surface of the quadrilateral shaped jugular process "homologous to the par-occipital process present in many animals." Thomson, 1917, also described a rough or smooth elevated surface, or else a projecting process springing from the under surface of the extremity of the jugular process, the extremity of which may articulate with the transverse process of the atlas. When this process is met with, he terms it paramastoid or par-occipital. According to Corner, 1896, Macalister, regarded the jugular process and its extension as the paramastoid or par-occipital. Bryce, 1914, described a distinct ridge occurring normally on the medial lip of the groove for the digastric muscle, which may be developed into a process of considerable dimensions and may even be rendered bullous by containing an air cell. This process he calls paramastoid. The term par-occipital is not mentioned in this text.

Corner, 1896, examined 304 mastoid processes and called attention to the existence of a process of the temporal bone

on the inner side of the digastric groove which was present in some form in 93 per cent of the specimens he examined. He also found a similar crest, ridge or process present on the temporal bones of 20 monkeys and in the skulls of 3 lemurs. He proposed in the consideration of the constant presence of this process, that the name "par-occipital" should be confined to the downward expansions of the occipital bones, and the name "paramastoid" should be applied to the process at the inner lip of the digastric groove.

Waldeyer, 1909, in discussing processes of this region, wrote that "Der Processus retromastoideus ist diejenige Bildung, welche mich zu einer erneuten Untersuchung der Regio occipitalis und mastoidea des menschlichen Schädels veranlasst hat und den Kernpunkt dieser Abhandlung darstellt." He added that the retromastoid process is "einen meist stumpfen, bald mehr rundlichen, bald mehr länglichen Fortsatz von durchschnittlich 0.5 bis 2 cm. Höhe, der sich an der Stelle des Zusammenstosses des oberen queren Nebenschenkels der Linea nuchæ inferior mit der Linea semicircularis superior entwickelt."

Other processes also have been described in this region. Waldeyer, 1909, stated that von Haferland described a rounded process which develops on both sides from the angulus mastoideus of the parietal and which projects laterally directly above the parietomastoid suture. Von Haferland termed this projection the processus astericus, but Waldeyer preferred the designation tuberculous supramastoideum posterius.

Since the prefix *para* means beside, beyond, accessory to, apart from, and against, the paramastoid process should by definition be near the mastoid process. When one examines this region in a series of temporal bones, one can find the mastoid process with the digastric groove in all degrees of depth—from a mere slight sulcus in cases in which the process shows no indication of division, to that of a deep groove a centimeter or even more in depth, where the process is well divided. This seems to indicate that this process, though frequently divided by a deep groove, is in reality one and the same process developmentally. Hence it would seem that when the

process is bifid due to the presence of a deep digastric groove, the outer portion should preferably be called the lateral and the other the medial portion of the mastoid process. And in cases where there are two grooves, as described later, the terms lateral, intermediate and medial could be used.

In Table 5, 500 adult temporal bones showing indications of a mastoid-occipital suture are classified with reference to the appearance of the medial portion of the mastoid process. The lateral portion of the mastoid process is present in all. Specimen 32 has a definite retromastoid process, and in three others it was indicated. In 37 cases two grooves were observed, one the digastric and the other the occipital, thus splitting the mastoid process into three portions with the presence of an intervening ridge or tubercle.

It is stated that in rare cases the extremity of this process may articulate with the transverse process of the atlas. The corresponding atlas is not present in our collection, but the length of the jugular process is such that this could well have been the case in this specimen, although its character suggests that this was not so.

The triplicate condition or that in which the medial portion of the mastoid process is divided, may be confined entirely to the temporal bone; or as in Figure 7, it may be more extensive, occupying the adjacent lateral portion of the occipital. In this particular case the medial portion of the mastoid process exists as two marked ridges. The jugular process articulated with a superior articular process of the atlas obliterating the jugular notch entirely, the condyloid and jugular processes being continuous.

Fig. 8 shows an abnormally large jugular process, more than 1 cm. in length, extending from the quadrilateral shaped space on the under surface of the occipital bone. This process has been described by Thomson, 1917, and by others.

Other conditions within the process also may be of unusual interest. Fig. 9 shows a single large mastoid cell, more than 1 cm. in diameter, occupying the medial portion of the mastoid process, while another cell, Fig. 10, same specimen, occupies the entire lateral portion of the process. This latter cell was $1\frac{1}{2}$ cm. in length, 1 cm. wide and 1 cm. in depth.

TABLE V.

Character of the portion of the mastoid process medial to the digastric groove in 500 specimens:

Ill defined or absent.....	118	23.6%
Bullous	69	13.8%
Short crest	79	15.8%
Long crest	107	21.4%
Rough irregular prominence.....	21	4.2%
Tubercle	65	13. %
Divided (by occipital groove)	37	7.4%
Retromastoid process (1 definite, 3 indicated)	4	0.8%
Total.....	500	100. %

THE OSSEOUS EXTERNAL ACOUSTIC MEATUS.

Of all the variable features of the normal temporal bone, the osseus meatus reveals the greatest number. One finds so many grades of variations in the size and contour that the determination of the normal form is quite difficult. Harrison Allen, 1884, in describing the relationship of the tympanic ring to the antrum, stated that "the curve of the U is much roughened and produced downwards." In Morris, 1899, it is stated that the external auditory meatus assumes the form of an elliptical bony tube. Toldt, 1908, depicted as typical an antrum nearly circular in outline. Cheattle, 1906, who wrote extensively on various aspects of the temporal bone, did not describe the form of the antrum. Thomson, 1917, in Robinson's Revision of Cunningham's Anatomy, fifth edition, described the meatus as oval in form with its long axis almost vertical near the orifice. Neither Piersol, 1907, nor Bryce, 1915, mention its form.

The specimens examined by me include all the above forms—from those almost round in appearance to those of an extremely oval type. The round forms are of various sizes, some measuring no more than 0.5 cm. in diameter at the orifice, although the maximum diameter at this point was 1.5 cm., a variation in size of 300 per cent. From the round form the picture changes by degrees to that of a decided

oval. All gradations in form can be seen, even if one observes only a relatively small number of bones. The shortest diameter of an oval form was 0.5 cm., and the longest, a little less than 2.0 cm. Between these figures one can find all possible forms and dimensions.

Oval forms were observed extending in a vertical plane, practically parallel to the styloid process. The opposite condition with the long axis in a horizontal plane and almost at angles to the styloid also was observed. Specimen 126, Fig. 4, illustrates the former condition, with the string over the long axis, and specimen 24, Fig. 5, approaches the opposite form. The intermediate form, which is the more common condition, is seen in Fig. 1, specimen 149. No form was observed in which the superior margin of an oval orifice lay posterior to a line drawn vertically through the center of the orifice of the meatus.

The size of an orifice may be greatly modified by exostoses. Cheattle, 1913, said that these always occur on the tympanic portion of the bone. Welcker, 1864, was one of the earliest anatomists to describe this form of exostosis. He found it far more common in skulls from American Indians than in any other race. Although exostosis had been observed by otologists in Europe at that time, their presence was regarded as quite rare. According to Welcker, 1864, who described three exostoses of the tympanic portion of the ring, Seligman had noted their presence in five out of six temporal bones from South American Indians.

An exostosis of the tympanic portion of the antrum may develop to a state of almost total occlusion of the meatus, as is seen in the accompanying photographs, Figs. 11 and 12, of an Indian skull found on Blaylock's Island, Columbia River, near Umatilla, Oregon. In the left meatus, Fig. 11, the extremities of the superior margin of the ring have become so thickened that they almost touch. Fig. 12 of the right meatus, shows a third somewhat smaller exostosis appearing at the inferior border of the ring, midway between two marginal thickenings. It is very doubtful whether a patent canal was present during life, for it probably was completely occluded by the exostoses with the overlying soft parts.

The failure of otologists and anatomists in general to record these exostoses would indicate that they are quite rare. Yet specimen 100, Fig. 13, has an exostosis present on the posterior margin—the anterior being normal. This, too, came from an Indian. But on looking over 75 more Indian skulls I failed to find another specimen with exostoses at these points.

Another structure which tends to diminish the size of the orifice is the postauditory tubercle which develops downwards from the superior margin of the external auditory orifice and may reach a goodly size. Although frequently absent, it may be modified and project outwards and downwards as a shelf or laminum. Specimen 1228, Fig. 14, presents such a case. In specimen 331, Fig. 15, this tubercle has a base of almost one centimeter, and is in the form of an equilateral triangle, the apex of which is directed downwards and slightly outwards.

The tympanic portion of the meatus may be generally thickened, even to one-half a centimeter, thus occupying some of the orifice, as the petrous portion is the denser bone of the two and yields least, hence diminishing the size of the orifice.

SOME POINTS IN THE ANATOMY OF THE HUMAN TEMPORAL BONE.

1. The average weight of 100 clean, dry, temporal bones was found to be 41.5 gms. and approximately the same on both sides.

2. The average angle between the superior margin of the petrous portion and the squamosa is 67.2° in the adult and 78° in the infant.

3. The angle between a horizontal and the posterior superior surface of the petrous portion varied between 105 and 125 degrees and coincided with the long diameter of the osseus external acoustic foramen.

4. A mastoid emissary canal was absent in 27 per cent of 250 specimens; was single in 67 per cent and quite variable in direction and number in 6 per cent. The external foramen usually is located above the digastric groove on a level with the upper margin of the osseus external acoustic orifice. Tubercles and exostosis were observed in the meatus.

5. In view of the present confusion in use of the terms paramastoid and par-occipital, it is suggested that we speak

of lateral and medial portions of the mastoid, and also in cases of triplicate division, of an intermediate portion, designating the other paramastoid processes as jugular and retro-mastoid.

REFERENCES.

- Allen, Harrison: 1884. *Human Anatomy*, Philadelphia.
- Bryce, T. H.: 1914. *Osteogoly*, Vol. 4, Part 1. *Elementary Human Anatomy*, Philadelphia. Quain's edition.
- Cheattle, Arthur H.: *Surgical Anatomy of the Temporal Bone*. Hunterian Lectures, 1906.
- Corner, E. M.: 1896. "The Processes of the Occipital and Mastoid Regions in the Skull." *Journal of Anatomy and Physiol.*, London, Vol. 30.
- Dwight, Thomas: 1907. *Osteology*, Human Anatomy. Piersol, G. A. Philadelphia.
- Gray, Henry: 1899. *Human Anatomy*, London.
- Gurriere and Masetti: *Rivista speriment. di Freniatria e de Med. legale*, 1895. Quoted after Piersol, *Human Anatomy*, Philadelphia, 1907.
- Meyer, A. W.: *Spolia Anatomica*. *Journal of Anat. and Physiol.*, Vol. 48, 1913-14.
- Morris, Henry: 1899. *Human Anatomy*, Philadelphia.
- Quain's Anatomy. 1914.
- Thomson, Arthur: 1917. *Osteology*, Human Anatomy. Cunningham, D. J. New York, Edition 5.
- Toldt, Carl: 1908. *Anatomischer Atlas*, Suchste Auflage.
- Waldeyer: 1909. *Der Processus retromastoideus*. *Abh. der Königl. Preuss. Akad. d. Wiss.*
- Welcher, H.: 1864. "Unber knöchern Verengerung und Verschlussung des äussern Gehörganges." *Archiv. für Ohrenheilkunde*, Vol 1-2.

Fig. I. Specimen 149. Exemplifying the relation of the long axis of the external acoustic orifice to the styloid process, the usual condition.

Fig. II. The same specimen, the string indicating the direction of the posterior superior surface of the petrosa, the usual condition.

Fig. III. Specimen 126. Showing the direction of the posterior superior surface of petrosa almost parallel to styloid process.

Fig. IV. The same specimen, showing the long axis of external acoustic orifice almost parallel to styloid process.

Fig. V. Specimen 24. The long diameter of meatus forms almost a right angle with styloid process.

Fig. VI. The line across the posterior superior surface of petrosa forms almost a right angle with styloid process.

Fig. VII. Triplicate division of the left mastoid process.

Fig. VIII. An abnormally large jugular process.

Fig. IX. A single mastoid cell medial to the mastoid process.

Fig. X. A single large air cell lateral to the mastoid process.

Fig. XI. Marked exostoses of the tympanic element in a left external auditory orifice.

Fig. XII. Right meatus with three points of exostoses.

Fig. XIII. Exostosis of the posterior portion of tympanum.

Fig. XIV. A laminum projecting outwards and downwards from the superior margin of the external auditory orifice.

Fig. XV. A large post-auditory tubercle.