of these moneys. This attack came like a bolt from the blue. Had I not had the Medical Defence Union to fall back on I would have had to suffer in silence; as it was, the result speaks for itself.

I am, Sirs, yours faithfully,

JAMES KERR, M.A., M.D. Cantab.,
Honorary Surgeon, Bradford Eye and Ear Hospital.
Bradford, Jan. 24th, 1898.

THE MARKING OF INFECTED HOUSES.

To the Editors of THE LANCET.

SIRS,—The College of Physicians of Philadelphia will at its next meeting in February discuss the regulation of the Board of Health placarding houses in which are contagious diseases. A member of the College has informed me that such a custom would not be tolerated in England and I will be grateful to you for any information on the rules and regulations of the London Board of Health in this matter of specifying houses in which there are contagious diseases.

I am, Sirs, yours faithfully,

GEO. WOODWARD,
Member, Board of Health, Philadelphia.
Bureau of Health, Philadelphia, Jan. 11th, 1898.

* * If circumstances should ever arise making such placarding an advantage to the public weal we believe that Englishmen would be sufficiently good citizens to submit at once to the regulation. The practice is, however, unknown in modern England.—ED. L.

"HYDROCYANIC ACID AS AN ANTIDOTE IN CHLOROFORM POISONING."

To the Editors of THE LANCET.

SIRS,—It is true that Dr. Lauder Brunton in his excellent work on Therapeutics¹ states that hydrocyanic acid is "one of the most powerful and rapid poisons known" and that it has "a special paralysing effect upon the respiratory centre," but it is also well known that different doses of the same drug may produce entirely different effects. As Dr. Brunton very aptly puts it in his more recent work,² "A stroke from a switch will quicken any animal to increased exertion, but a blow from a bludgeon may knock it down and render it incapable of any exertion at all."

The use of hydrocyanic acid as an antidote to chloroform is an almost typical illustration of this. When given to a healthy animal in large toxic doses it speedily causes death, the latter occurring so rapidly that it requires a keen observer to note any very numerous train of symptoms. When, however, a smaller but still toxic dose is given definite symptoms (which are divided into three stages by Dr. Brunton) can be observed. With a still smaller dose—a dose which is the full medicinal or, perhaps, slightly over medicinal without being fatal—it is possible to produce certain results which are directly antagonistic upon the heart and respiration to those caused by over-doses of chloroform; this is the stage which is aimed at. In a healthy animal such a dose will within a few seconds produce respirations which are at first somewhat irregular and slow, but speedily become strong and powerful, whilst the pulse is at first slightly slower, then quickly increased in frequency and somewhat irregular, in a short time becoming full and regular.³ Upon the heart, when the animal is deeply anaesthetised, prussic acid in these doses seems to have a distinctly beneficial effect in relieving the congested, semi-paralysed condition into which that organ has been thrown. Another great advantage, too, that this antidote possesses is that when once respiration has recommenced the pharmacological action of the drug tells us that the latter will greatly assist in maintaining it until the centre has sufficiently recovered to be able to look after itself, as we know that the visible action of the acid will last for about half or three-quarters of an hour.

I was particularly struck when observing its action upon the first few cases with the rapidity with which animals recovered from prolonged operations under chloroform when a small dose of hydrocyanic acid was given to them. I formed the

opinion at the time that this was largely due to the stronger and deeper inspirations and expirations respectively causing a greater amount of fresh air to be inhaled and also more of the chloroform vapour to be expelled. This seems, too, to afford a reasonable explanation.

I am, Sirs, yours faithfully,

FREDK. HOBDAV.
Royal Veterinary College, London, N.W., Jan. 24th, 1898.

"TRANSMISSION OF SYPHILIS TO THE THIRD GENERATION."

To the Editors of THE LANCET.

SIRS,—Dr. George Ogilvie's conclusions on hereditary immunity in the third portion of his interesting paper, which is referred to in THE LANCET of Jan. 15th, are to a considerable extent founded on the discredit thrown by Professor Neumann upon Fergusson's observations on mitigated disease in Portugal in 1812. Dr. Ogilvie considers on those grounds that Mr. Henry Lee's views on individual hereditary immunity, dependent as they were to some extent upon Fergusson's observations, seem to "stand on a weak foundation"; that "the reports from different Russian, German, and English regiments adduced by Ferguson and Lee have, if possible, still less claim to trustworthiness"; and that the sad state of the British army at the present time in India (which I might say is a counterpart to the state of the army noted by Fergusson in Portugal) is "sufficient to considerably shake any theory of general immunity." The observations of William Fergusson (not Ferguson), for two years the Inspector-General of Hospitals to the Portuguese army, seem to me to be so conclusive and, moreover, verified at the time by Staff-Surgeon Jebb that it is impossible to accept Dr. Ogilvie's interpretation.

Everyone will agree that hereditary immunity is not the only cause of modified syphilis in the male. Neisser, while attributing the diminution of malignant syphilis (in his paper on Malignant Syphilis in the *British Dermatological Journal* for January, 1897) to hygienic, preventive, and therapeutic causes, which Dr. Ogilvie acknowledges, nevertheless says: "Nor have I noted during the recent great extension of syphilis in Russia that an unusual proportion of cases of malignant syphilis has been noted it is interesting in this reference to draw attention to the fact that Epstein has recently recorded a remarkable number of cases of malignant syphilis, amongst the Jews he is inclined to account for his observations on the hypothesis that the individuals of a community which has been free from syphilis tend to show the malignant type more frequently when at length attacked than the members of a population which has experienced the disease for generations." That is Fergusson's theory, which to be fully appreciated should be read in its entirety in the fourth volume of the Transactions of the Royal Medical and Chirurgical Society. I am, Sirs, yours faithfully,

JOHN A. SHAW-MACKENZIE.
Grosvenor-street, W., Jan. 18th, 1898.

"THE MECHANISM OF DEATH FROM CHLOROFORM."

To the Editors of THE LANCET.

SIRS,—I note in THE LANCET of Dec. 4th, 1897, a letter by Mr. Alexander Wilson called forth by my criticisms on his article published in your issue of Sept. 11th. His first point is that I did not read his paper carefully nor give his expressions their proper value. I read the paper sufficiently carefully to notice the immediate juxtaposition of the expressions "free respiration" and "breathing deeply," with the expression "some peculiarity about the colour of the face," this last denoting an attitude of mind so slovenly and inexact that it is impossible to consider the two former as necessarily accurate. Similarly in Case 3 the very point of my criticism is that with observation so inaccurate that the intense circulatory failure present passed entirely unnoticed by the anaesthetist and was only discovered on making an incision the further statement that respiration was going on freely cannot be taken on trust. As the facts in these two cases are not unimpeachable their value as the basis of a theory is gone.

As regards the question asked by Mr. Wilson with respect

¹ Pharmacology, Therapeutics, and Materia Medica. Third edition. Pp. 586 and 234.

² Action of Medicines, p. 34.

³ Journal of Comparative Pathology and Therapeutics, 1893, p. 107 et seq. Idem, 1896, p. 101 et seq.