nurse and told her she felt perfectly well. The tenderness had entirely left the face, the toxic symptoms had subsided and the pulse and temperature were nearly normal. At 8 P.M. the patient had a chill, which was followed by another exacerbation of the disease. The heart was overpowered and showed signs of extreme intoxication. The disease extended into the meatus of the ear, the temperature reached 103.5, which was followed by a mild delirium. At 1 P.M., the serum which I had ordered the day previously having arrived, I again injected 10 c.c. with most happy results. Within two hours the heart was again beating naturally and the symptoms gradually subsided. In twelve hours the temperature was normal, the pulse only slightly accelerated and the tenderness all gone. I then injected 5 c.c. as a prophylactic against further exacerbation and in twenty-four hours I again repeated the dose. The patient made an uneventful recovery, desquamation taking place within twenty-four hours from the last injection.

Within a few weeks I was called to attend a woman suffering from puerperal fever. From the clinical picture presented, as well as from facts surrounding her confinement, there was no question in my mind as to the diagnosis of streptococcic infection. Her temperature was 104.5, her pulse 150. She was a young woman of robust constitution, and afforded an excellent opportunity for the demonstration of the efficacy of antistreptococcic serum in this type of puerperal infection. Ten c.c. of Parke, Davis & Co.'s serum was injected, which was repeated in twelve hours. In twenty-four hours from the second injection the patient was absolutely well, no indications of the previous infection being in any way manifest. I have used the serum in a number of cases with results varying in direct ratio to the positiveness of my diagnosis. In all cases where I knew that the infection came from the streptococcus erysipelatus, the results were uniformly good.

Dr. Madden handed me the following report of a case which he exhibited before the Milwaukee Medical Society:

L. A. M., 40 years of age, subject to repeated attacks of facial erysipelas during the past four or five years, has had eight attacks of considerable severity. The disease generally began on some point of the face and extended over the whole head, on two or three occasions involving the neck and one shoulder. The length of these attacks varied from fourteen days to two and a half months. About February 1 a characteristic attack began on the right ala of the nose, extending upward and backward over the right cheek. The patient applied for treatment on the second day of the disease. At that time the parts bore all the characteristic signs of erysipelas. On the morning of the third day 10 c.c. of Parke, Davis & Co.'s antistreptococcic serum was injected in the subcutaneous tissue between the shoulders. Two hours afterward the patient declared that he felt relieved. In twenty-four hours the activity of the disease seemed to have completely ceased, and in forty-eight hours desquamation had begun. The progress of the case from this point was uninterrupted until complete recovery.

In conclusion, I would say that, owing to the fact that this serum has not yet reached a state of perfection and can not as yet be accurately standardized, we are apt to meet with some failures and be subjected to some disappointments in its use. A great many diseases of mixed origin are supposed to be benefited by this serum, but the results of its use must be determined by the extent to which the streptococcus erysipelatus infection acts as an etiologic factor in the disease. Any disease which is caused by streptococcic infection is certainly benefited, if not

cured, by antistreptococcic serum. Serum which is reasonably fresh and hermetically sealed, having been properly prepared and properly inspected, is never an element of danger and may be used with impunity, in reasonable quantities, where the diagnosis is not clear, but where streptococcic infection is suspected.

SOME OBSERVATIONS ON THE AURI ET SODII CHLORIDUM U. S. P.

Presented to the Section on Materia Medica, Pharmacy and Therapeutics, at the Forty-ninth Annual Meeting of the American Medical Association, held at Denver, Colo., June, 7-10, 1898.

BY DANIEL R. BROWER, A.M., M.D., LL.D.

Professor of Mental Diseases, Materia Medica, and Therapeutics, Rush Medical College, University of Chicago; and Professor of Diseases of the Nervous System, Woman's Medical School, Northwestern University, etc.

CHICAGO.

Auri et sodii chloridum is a mixture of equal parts of dry gold chlorid and sodium chlorid. It is very soluble in water and very easily decomposed by sunlight and organic matter. This drug is tonic, alterative, bactericidal, and antiseptic. It acts on the digestive tract, in small doses improving the appetite, stimulating the functions of the liver and stomach, promoting assimilation and general nutrition. The larger doses continued produce epigastric pain, nausea, and anorexia. Still larger doses produce violent gastro-enteritis without ulceration and salivation. Six and a half centigrams (gr. 1) have produced these symptoms in an adult, and this same dose has produced death in a dog in four minutes.

In anemia, in small doses, it increases percentage of hemoglobin and the number of red blood corpuscles; on the nervous system, in small doses, it is a cerebral stimulant, making the mind more active, and often inducing in the mentally depressed a sense of cheerfulness; upon the spinal cord it acts as a stimulant. In man it acts as an aphrodisiac, and in woman it augments the menstrual flow. In large doses it produces paralysis of the central nervous system, first in optic lobes and cerebellum, then in spinal cord, and then in cerebral lobes. In medicinal doses it increases urinary secretion and the percentage of urea. In large doses it will produce renal hyperemia and albumin in urea.

The bactericidal and antiseptic powers have recently been investigated by Prof. L. Hektoen, at my suggestion, and he writes as follows:

May 26, 1898.

Dear Doctor:—Sternberg, in his Manual of Bacteriology, made the following statement of the bactericidal and antiseptic properties of chlorid of gold and sodium:

	Destroys.	Restrains growth.
Anthrax bac.	1:8000	1:40,000.
Diphth. bac.	1:1000	1:40,000.
Bac. glanders	1:400	1:15,000.
Bac. typhoid	1:500	1:20,000.
Spir. Asiatic chol.	1:1000	1:25,000.

I have had the necessary experiments repeated by Mr. Danielson, who finds that with our cultures the above figures hold good with the bacillus of diphtheria and typhoid fever as to bactericidal powers, but that the dilutions mentioned do not destroy the other germs mentioned. He found that growth was delayed by the dilutions of Sternberg in the second column.

Yours sincerely,

L. HEKTOEN.

Shoemaker (Materia Medica and Therapeutics, 3d, Ed., p. 226), quoting Calmette, says that it is antidotal to the poison of the cobra de capello. Gibbs and Shurley (*Therapeutic Gazette*, April 15, 1891) demonstrated its bactericidal power against bacillus tuberculosis: First by inoculating guinea pigs and a

week afterward treating them with auri et sodii chloridum, limiting the lesion to the seat of the inoculation; second by putting healthy animals under treatment for one or two weeks before inoculation, and thereby preventing in these even the formation of an abscess at the seat of inoculation.

The elimination of gold and sodium chlorid takes place principally by kidneys and, to a small extent, by liver and bowels. There is an astonishing amount of skepticism in the profession as to this drug. Some of this, at least, arises from the fact that it is very unstable, being readily decomposed by light and organic matter. Sufficient care is not exercised in its combination with other drugs, or too large a quantity is ordered, so that decomposition has taken place before its ingestion, or its administration is not properly timed, so that destruction takes place in the stomach. The best vehicle with which to combine it in capsule is tragacantha or guaiac resina; neither of these decomposes it. The time for administration should be one hour after eating, or, better still, one hour before eating. The ideal method for administration is by hypodermatic injection, the solution used being made with equal parts aqua destillata and glycerina.

Syphilis.—The remedy is useful in syphilis, especially in those cases in which the iodid and mercurials do not agree, or in which there is a great amount of debility. Dr. E. F. Ingals (*Diseases Chest and Throat*, page 448) recommends it in syphilitic laryngitis. M. Martineau (*Bull. et Min. de la sac, de la Therap.*, April 30, 1883) recommends it in cases of inveterate syphilis. One of his cases, a woman in an advanced state of cachexia, broke her fibula, probably because of specific change in the bone. Though rebellious to other treatment, she recovered quickly and the fracture united readily under gold treatment. W. Hale White (*Mat. Med. and Therap.*, Wilcox, 2d Ed., p. 187) says it is a valuable remedy in the tertiary manifestations of syphilis, especially of the bones, and presents fewer disadvantages than does the corrosive mercuric chlorid. Shoemaker (*Mat. Med. and Ther.*, 3d, Ed., p. 226) and Potter (*Mat. Med., Pharmacy and Therap.*, 3d Ed., p. 123) both emphatically endorse it for the treatment of syphilis in its later manifestations.

The drug is useful in the treatment of connective tissue hyperplasia, as in cirrhosis of the liver, interstitial nephritis, arteriosclerosis, and the spinal sclerosis. Millard (*Bright's Disease*, p. 220) writes: "The chlorid of gold has proved of great value in interstitial nephritis. Under its use I have often known the albumin to diminish and disappear;" and again, "At all events its usefulness in chronic nephritis has sometimes been unmistakable, and it is likely to prove still more useful if the patient suffers, as is usual, from nervous symptoms, hypochondriasis, irritability, vertigo, etc." Bartholow (*Practice of Medicine*, p. 400) writes, anent topic interstitial nephritis: "Better results even, the author believes, are procured from the careful and persistent administration of the chlorid of gold, or of gold and sodium." Baumgarten (*Reference Handbook Medical Sciences*, Vol. iv, p. 282) writes (referring to Dana, *Boston Medical and Surgical Journal*, Oct. 24, 1883) under "Contracted Kidney": "Marked improvement and mitigation of distressing nervous symptoms have been observed from the use of chlorid of gold and sodium (0.005=1/12 gr.) in pill after each meal. Culbertson

(*Am. Jour. Oph.*, July, 1894) relates a case of albuminuric retinitis relieved by chlorid of gold and sodium. W. H. Walling (*Medical and Surgical Reporter*, Philadelphia, 1897, Vol. lxxvi), writing of sclerosis generally: "I will state my belief, founded on an experience of twenty years, that gold is far more efficient than any other drug I know of."

My first experience in the use of this remedy in interstitial nephritis was about fifteen years ago. The case came under my treatment because uremic symptoms had developed after severe exposure. When these acute symptoms disappeared he was given chlorid of gold and sodium, and he continued it in varying doses for five years, the albumin gradually diminishing and finally disappearing. Several examinations of the urine were made five years ago and no casts or albumin were found. I am sure that I have also repeatedly seen advantage from its use in the early stage of posterior and combined spinal sclerosis.

Diabetes mellitus.—J. A. Robison (*Chicago Medical Recorder*, May, 1891) reports two cases of diabetes mellitus cured by its use. A recent communication from Dr. Robison informs me that these cases continue well and that an equally favorable result has been reached in other cases.

Functional diseases of the nervous system.—Niemeyer (*Text Book Prac. Medicine*, p. 386) writes: "I have accidentally hit upon a nervine of great efficacy in hysteria and have made use of it with signal effect in many cases." The formula he recommends is:

R. Auri et sodii chloridi 0.30 gm. (gr. v)
Gummi tragacanth 4.00 gm. (dr. 1)
Sacch. alb. q. s. ft. in pil. No. 40.

Sig. Begin with two pills a day, increased to eight pills daily.

Mills (*The Nervous System and its Diseases*, p. 238) writes: "Sodium and gold chlorid, although it has attained an unmerited reputation in some directions, is a valuable tonic, and I not infrequently use it in the treatment of hysteria and other affections."

My personal experience is in accord with these great clinicians, and I especially desire to emphasize its value in that complexus of symptoms called neurasthenia.

Tuberculosis.—Gibbs and Shurley (*Therapeutic Gazette*, April 15, 1891) report twenty-seven cases of pulmonary tuberculosis treated with the drug, used hypodermatically, in doses 0.003 gm.—.001 gm. (1/20 gr.—1/6 gr.) during three to eight months, with gradual disappearance of temperature and cough, increase in weight, and disappearance of bacilli. Pepper (*Univ. Med. Mag.*, December, 1895) reports a case of phthisis apparently cured by the use of this drug.

GOLD CURES IN INEBRIETY.

Presented to the Section on Materia Medica, Pharmacy and Therapeutics, at the Forty-ninth Annual Meeting of the American Medical Association, held at Denver, Colo., June 7-10, 1898.

BY T. D. CROTHERS, M.D.

SUPERINTENDENT WALNUT LODGE HOSPITAL,
HARTFORD, CONN.

Any remedy introduced, surrounded by mystery and sustained by falsehood and deception, is unworthy of notice. Yet a number of physicians have expressed confidence in the value of gold as a remedy, both in inebriety and allied brain affections. This theory has been recognized by reputable druggists, who have prepared several preparations of gold for administration chiefly by the needle. The empiric prepara-