the work under notice, "We always receive with pleasure suggestions for the improvement of this publication, and within reasonable limits never allow either trouble or cost to prevent the adoption of all which in any way commend themselves to our judgment."

Your suggestion is as follows:—

"The publication of the monthly as well as the annual amounts of rain for the whole of the 1,700 stations is very desirable, and it is hoped that in an early issue of the 'British Rainfall' it will be done."

I shall be glad if you will allow me to supplement the data which were before you when the above paragraph was written by some other facts, and to learn from your pages whether or not this fuller information induces any modification of your views.

As I (whether fortunately or unfortunately I need not say) have to pay my own printer's bills, I always keep them as low as possible; hence, the publication being an annual one, statements made in one volume are rarely repeated in the next. Therefore, probably, your reviewer was not aware of the principles upon which the tables of monthly rainfall (pp. 140-145) are compiled, viz., to give one station in every county in the British Isles, and two in a few of the larger ones, such as York, Inverness, and Ross. I may add *en passant* that these tables give the monthly fall at 108 stations, while the Registrar-General of England is satisfied with forty-four, and of Scotland with fifty-five; so that my table exceeds both together. That, however, is of little moment. [For your own information, I enclose a map with these 108 stations plotted.]

In the next place, I must refer to "British Rainfall, 1871," pp. 135-138, where the question of publishing additional monthly returns is discussed at length, and the method of computing the monthly fall from the percentage tables (which are given every year) is explained and illustrated by a completely worked-out example.

To this let me add that returns from 150 other stations are published monthly in my *Meteorological Magazine*, and that up to the present time another very large series (143) has been printed biennially in the Reports of the British Association.

If it is the opinion of yourself and of others competent to judge that still more is necessary, more shall be done; but it must be borne in mind that the accurate (and without accuracy figures are worse than useless) printing of 20,400 values involves a great expenditure both of time and of money. I do not quite know whence either the one or the other is to be obtained.

G. J. SYMONS

[It was just because of the inadequacy of one station in each county of the British Isles, and two in the larger counties, to represent the rainfall, even though these be supplemented by Mr. Glaisher's forty-four stations, the Scottish Meteorological Society's two hundred odd, and by Mr. Symons himself in his *Magazine* and in the *British Association Reports*, that we stated it to be very desirable that the monthly as well as the annual amounts of rain for the whole of the 1,700 stations were published. The method of computing the monthly fall from the percentage tables referred to in "British Rainfall, 1871," pp. 135-138, does not supply what is desiderated. It is the capriciousness of the distribution of the rainfall and its important bearings on many practical questions which render so desirable a knowledge of the actual monthly amounts in particular localities. Since what is desired would be an invaluable contribution to British Meteorology, we earnestly hope that Mr. Symons will be induced to supply it, and that in that case he will receive substantial support in carrying on a work so important.]

Equilibrium of Temperature in a Vertical Column of Gas

I OBSERVE that Mr. R. C. Nichols, in his letter to NATURE (vol. xii. p. 67), admits that the mean energy of molecules "may" remain the same at all points of a vertical column. It is not difficult to show that it *must* do so if the velocities are distributed among the molecules according to the exponential law.

As I have never seen any direct proof of this in English I extract the following from Boltzmann.

In order not to take up too much of your space, we will take the simplest case, and suppose the molecules to be equal elastic spheres, moving in a vertical tube with elastic base and sides. Let them be acted upon by vertical forces, the potential of which

at height x above the base is $f(x)$. Assume first that no encounters take place between the molecules, and let the number of molecules at the base, the energy of whose vertical velocity

is v^2 , be $Ce^{-\frac{mv^2}{2k^2}}$ where C and k are constants. For each molecule the sum of the potential and kinetic energies is constant.

And as the horizontal velocities are constant, it follows that for each molecule the sum of the potential energy and the energy of vertical velocity is constant. That is, the energy of vertical velocity is diminished by $f(x)$ in the ascent from the base to x .

Therefore the molecules which at height x have u^2 for energy of vertical velocity are the same identical molecules which at the base have $u^2 + f(x)$ for energy of vertical velocity.

Their number is therefore $Ce^{-\frac{u^2 + f(x)}{k^2}}$ that is $\frac{f(x)}{k^2} Ce^{-\frac{u^2}{k^2}}$.

Therefore the number of each class at x is the same as the number of the same class at the base multiplied by the factor $e^{-\frac{f(x)}{k^2}}$.

Evidently the mean energy is the same at all points of the tube, and the density only varies, and is represented by $e^{-\frac{f(x)}{k^2}}$.

Again, still precluding encounters, let the velocities of the molecules in each of two horizontal directions at right angles to each other be distributed according to the same law as the vertical. And further, let the chance of a molecule having given horizontal velocity in either direction be independent of its velocity in the other horizontal direction or in the vertical. The same distribution and independence will be maintained throughout the tube. And we see that force has no tendency to disturb it.

Maxwell has shown that among such molecules as we have supposed encounters have no tendency to disturb the given distribution, which must therefore remain undisturbed though force and encounters both be present.

S. H. BURBURY

Primine and Secundine

WILL you allow me to avail myself of your pages as a means of pointing out to those who have purchased the English edition of "Sachs's Text-book of Botany" an unfortunate error which Prof. Oliver has been so good as to point out to me?

On p. 501 the inner coat of the ovule is identified with the "Primine" of Mirbel, and the outer with the "Secundine." The application of these terms is exactly inverted. The confusion easily arises from the fact that the secundine is developed first and the primine second. Mirbel, however, ignorant of, or disregarding that fact, numbered his structures from without inwards. The outer coat he termed the primine, the inner the secundine, the nucleus the tercine, and so on to quartine and quintine.

Except for the sake of accuracy the matter is of no essential consequence. Those who study the coats of ovules may well be indifferent to Mirbel's perplexing terms. But in these days, when students are expected for examination purposes to know about the names of things rather than about things themselves, it might lead to deplorable consequences, of which I hasten to relieve myself of the responsibility.

W. T. THISELTON DYER

American Indian Weapons

IN Col. Lane Fox's Catalogue of his Anthropological Collection he quotes Schoolcraft as saying, "There is no instance amongst the North American Indians in which the war-club employed by them is made of a straight piece, or has not a curved head." I send you a drawing (Fig. 1) of a club in common use among the Numas, or Indians of the Great Interior Basin, embracing Shoshones, Utes, Pueblos, &c., which will no doubt interest Col. Fox and others, not only on account of its extreme simplicity of form, but also of its method of use. It might be called appropriately a "face-masker," being grasped with the bulb next to the little finger, and thrust into the countenance of the foe. Major Powell sent a number of these to the Smithsonian Institution. They are of one piece of wood, generally mezquite, either very rude or quite smoothly polished, and are worn attached to the wrist by a leather thong. They vary in length from eight inches to fourteen. These same tribes use a simpler "slung shot" than the one described in Col. Fox's Cata-

logue, p. 65 (Fig. 2), the stone ball hanging loosely from the handle in a bag of buckskin. The Moquis of this same region use the boomerang; two of these (Fig. 3) are in the Smithsonian

Institution. I am not sure that it returns to the hand of the thrower.

On page 91 of Col. Fox's Catalogue he says: "In California

Fig. 1.—Pai-Ute War Club, for thrusting by a backhanded blow into the face of an enemy. Made from the wood of the Mezquite bean.

Fig. 2.—Pai-Ute War Club. Fig. 3.—Moquis Boomerang. Fig. 4.—Numa Reed Arrow, with hard wood foreshaft. Fig. 5.—Klamath River Pointed Arrow; soft wood shaft, hard wood foreshaft. Fig. 6.—Klamath River Arrow, without point; soft wood shaft, hard wood foreshaft sharpened.

and the greater part of the North American Continent the arrows are constructed either in a single piece or with a bone foreshaft; but in no case have I come across a foreshaft of hard wood." Among the Numas of the Great Basin, reed arrows with hard wood foreshaft are very common (Fig. 4). In Northern

California two kinds of arrows have hard wood foreshaft, those with and those without stone points (Figs. 5 and 6). The stripes on the feather end are rancheria marks, and the foreshaft is moveable.

OTIS T. MASON

Washington, D.C., U.S., May 19

Primroses and Cowslips

MR. FORDHAM (NATURE, vol. xii. p. 87) is quite right in conjecturing that it may be without foundation he has thought that primroses are not found in districts in which cowslips are common, and *vice versa*. In the north-east of Staffordshire, for miles round Denstone College, early in the spring, nearly all the hedges and many of the fields are covered with primroses. Later on cowslips abound; I might add that oxlips are also far from being rare.

I have watched closely, but have never found a trace of any destruction of the flower by birds. This, perhaps, may be accounted for by the fact that this being a pasture country, the sparrow, finding no grain, is a *rara avis* about here. I have noticed in Lord Bagot's wood, some twelve miles from here, where sparrows as well as many other birds are found in great numbers, that the primroses nearly always present a very ragged appearance.

D. EDWARDS

Denstone College, Uttoxeter

I COULD name half a dozen spots to the north of London (Mill Hill) where cowslips and primroses have abounded together in the same meadow, to my own knowledge, for the past twenty years. For at least five years I can say that neither the primroses nor cowslips were attacked by birds, though the crocuses were cut up by them more or less every season in the same locality.

R. A. N.

THE VISITATIONS OF GREENWICH AND EDINBURGH OBSERVATORIES

WE have before us the Annual Reports of the Astronomers Royal for England and Scotland, to their respective Boards of Visitors. The Report of Sir G. B. Airy consists mainly of the usual statements under the various heads of the state of the buildings and instruments, the constitution of the staff, and the amount of work done. In all these respects the Observatory seems to be in a satisfactory condition. One important change in the staff during the past year has been the resignation of

Mr. Glaisher, who has for so many years been connected with the Observatory, and which has rendered necessary a readjustment of the duties of the various observers.

Under the head of "Chronometers, Time-signals," &c., the Astronomer Royal refers to the supplemental mechanism which he himself has introduced into some chronometers in order to correct the perceptible defect of thermal compensation which occurs in nearly every case, even in the best chronometers. "There is," he states, "great difficulty in correcting the residual fault, not only because an inconceivably small movement of the weight on the balance-curve is required, but also because it endangers the equilibrium of the balance. To remedy this I have introduced small supplementary weights carried by means of a supplementary bar (rotating with stiff friction in the balance-staff), at whose ends are very light springs carrying the supplementary weights, and constantly pressing them to the interior of the balance-curve. When the supplementary bar is so turned that the supplementary weights are near the end of the balance-curve, the compensation is large; when they are near the root of the balance-curve, it is small. The movement from one state to the other is so simple that probably an assistant of the Observatory will be able to manage it, and it does not interfere with equilibrium. This arrangement has received the approval of some able chronometer-makers, and may perhaps with advantage be adopted generally."

The various time-signals and clocks connected with the Observatory have been worked with praiseworthy regularity and accuracy; the Westminster clock has been so well regulated, under check of automatic report to the Observatory, that in 83 per cent. of the days of the year its error is below one second. Proposals have been made for galvanic determination of the longitude of the Dublin Observatory, and the operation is delayed only for convenience in the arrangements to be made at Dublin. With the aid of a grant from the Treasury three computers are now steadily at work on the Astronomer Royal's New Lunar Theory.

The most novel and interesting part of Sir George Airy's Report is his concluding "General Remarks," in which