to individual sensitiveness. This was noted in every case, after injection, seeming to indicate the new serum to be antagonistic to the internal secretion of the pituitary body, as far as blood pressure is concerned. Thirty cases of diabetes mellitus were treated; of these, 8 did not respond with any noteworthy diminution of glycosuria, 14 had a complete disappearance of it while the intake of carbohydrates was notably increased; in 8, a diminution of glycosuria was seen. Polyuria commonly disappeared with rapidity, even in the most unfavorable cases. The favorable ones also showing improvement of the general condition and of special symptoms such as neuritis, retinitis, emaciation, headache, etc. Four cases are reported *in extenso*. [Author's abstract.]

Gordon, M. B. PINEAL GLAND IN PEDIATRICS. [Endocrinology, Oct.-Dec., 1919.]

The author does not consider that sufficient facts have been given in the literature to allow of any firm opinion as to the functional activity of the pineal gland. Does it possess an internal secretion? Nothing has been proved. Experimental work has failed to prove that it possesses a function, and no experimental studies are so complete as to allow comparison with the very striking syndrome seen clinically. Horrax and others maintain that the pineal gland controls the inhibition of sex growth, then pineal feeding should postpone adolescence, but by Dana, Berkeley and McCord proved the opposite. Yet, if the feeding results of McCord are correct, strong evidence of a pineal function ought to have followed the extirpations of Dandy and Horrax. Finally, Gordon says all knowledge of the pineal function is more problematic than accurate.

Krabbe, K. EARLY SYNOSTOSIS OF THE EPIPHYSIS WITH DWARFISM IN PUBERTAS PRÆCOX. [Endocrinology, Oct.-Dec., 1919.]

The case given by Krabbe is a girl thirteen and a half, who, when only a few months old, had bleeding from the vagina with recurrence every four weeks since. From the sixth to the seventh year the breasts were the most prominent, then there was a diminution in size; now they are like the breasts of a middle-aged virgin. For two or three years a growth of hair in the axilla and over the pubis was noticed. Her hips and thighs have always been large. Growth was rapid until she was seven, since that time it has stopped. The lower limbs are strikingly short in relation to the trunk. X-ray showed a normal sella turcica. The epiphyseal fissures of the lower and upper limbs were completely grown together. The patient's face is childish and gives no evidence of sexual knowledge in spite of the marked development of menstrual function and external evidences of sexual maturity. Seeing this, no intravaginal examination was made, so the possibility of an ovarian abnormality being the cause of the condition could not be eliminated. Thyroid