act on his own responsibility. This he will always attempt to shirk, and the fond parent will almost invariably finally, if not at first, accept his excuses. Particularly youths of this temperament should not be forced to comply with some arbitrary parental conception as to their life work. This ought to be decided late and only after a careful scrutiny of their talents and tendencies. It is in these cases, if not indeed, in all cases properly a problem of adolescence.

Unfortunately, many individuals with this temperament are impelled by undue activity and a weak judgment into speculative enterprises, which of course, exposes them more than anything else to the liability of the development of their latent malady. Practically, it is not nearly so difficult to decide on an occupation or course in life that would be desirable for this or that individual, as it is to secure it for him.

The marriage of two young people of this temperament furnishes some of the darkest chapters of domestic discord and disaster ever written. When they have once fallen in love, however, the mischief has been done, for they can rarely be persuaded even to postpone their nuptials, while violent and forcible measures of interference, even if effective, are liable to precipitate the development of insanity. Unless by strategy he can offer some plausible suggestion that would involve separation and delay, the physician must observe with patient composure the early scenes of a drama, apparently merry enough, but which he knows full well is likely only too soon to be fraught with passages of the most pitiful pathos and perhaps even tragedy.

100 State Street.

CARCINOMA OF THE BOWEL AND OF THE APPENDIX IN THE YOUNG *

J. SHELTON HORSLEY, M.D.

Professor of the Principles of Surgery and Clinical Surgery in the Medical College of Virginia; Surgeon to Memorial Hospital
RICHMOND, VA.

In no part of the body is the onset of cancer more insidious than in the bowel. As it is a local disease at first and becomes disseminated later on, an essential for successful treatment is early diagnosis, and anything tending to aid in this respect is of great practical advantage to the surgeon. It was formerly thought that carcinoma of all structures occurred in middle or later life, and a strong point in the differential diagnosis between carcinoma and an inflammatory mass, or a benign growth, has been the age of the patient. If the age were 40 years or more the chances seemed to favor cancer; under 40 years the likelihood of carcinoma apparently decreased with decreasing years, till its occurrence in childhood has been looked on as a pathologic curiosity.

Recent reports, however, seem to show that carcinoma of the bowel, and particularly of the appendix, is met with in the young more frequently than was formerly supposed. During the past year there has been a large amount of literature on cancer of the appendix. In most cases reported one is struck by the youth of the patient. Thus, Harte¹ collected reports of 92 cases of carcinoma of the appendix and finds 4 cases in patients under 10 years of age, 13 between 10 and 20 years of age, and 34 between 20 and 30 years of age. The age

of 4 patients is not given, so of the 88 patients with carcinoma of the appendix whose ages are known, 51 are under 30 years of age. Harte is impressed with this fact and explains it with the statement that "a large percentage of all carcinomatous appendices are found as a result of operations for appendicitis, and most operations of this character are done between the ages of 10 and 40 years." He believes that small foci of cancer may be found in other parts of the intestinal tract if carefully looked for.

If cancer of the appendix be discovered more frequently in the young because the age of the patient operated on for appendicitis is usually under 40, we would expect to find cancer frequently in other organs usually removed in the young for inflammatory conditions. No one has found any large number of carcinomata in the Fallopian tubes, though they are often removed in the young for inflammatory conditions. Probably the largest number of curettements is done during early child-bearing period, and though the scraping may be carefully examined, it is rather rare to find cancer of the uterus under 35 years of age.

The most probable explanation of the occurrence of such a large proportion of cancers of the appendix in the young is that it is due to the peculiar character of the appendix. Carcinoma is a malignant tumor of epithelial origin, and sarcoma a malignant tumor of connective tissue origin. We know that connective tissue is a simple and hardy tissue, requiring but little nutrition, and is able to overcome difficulties and live under conditions that would be impossible for the more delicate and highly differentiated epithelial tissue. Consequently, we find the connective tissue malignant tumor (sarcoma) can overcome the obstacle of the marked resistance of youth which the epithelial malignant tumor (cancer), being more delicate and a higher order of tissue, can not do. So cancer can gain a foothold, as a rule, only when the resistance of youth has been partly lessened by advancing years; it therefore usually begins in middle life or in old age. The appendix is a vestigial organ; its tissues are often irregularly arranged and its size and blood supply are very variable. It is a rule in anatomy that the more important an organ is the less liable are great variations to occur. A small difference in the size of an important organ like the liver or heart would be most noticeable, yet it is not uncommon to find appendices varying in length from one to six inches. The poor resistance the appendix offers to inflammation is a matter of common knowledge. It is but natural to expect, then, that the resistance to cancer of such an irregular and vestigial organ as the appendix would be very low even in the young, whose other tissues might be fully able to withstand the onslaught of this disease.

The reasons for the occurrence of cancer in the bowel or appendix in the young are these: 1. There may be a weakened resistance in patients whose tissues are prematurely old as a result of hardship or constitutional disease; this condition may, however, be purely local, as in the case of the appendix, in which, on account of the unimportance of this organ to the physiology of the body and its consequent irregular and uncertain structure, resistance to cancer is of a low type. 2. The other reason for the appearance of cancer in the young is that this disease may take a peculiarly virulent form. In other words, a weakened resistance may permit cancer of an average type to gain a foothold, or a cancer of unusual virulence may overcome a normal resistance.

* Read at the Tri-State Medical Association of the Carolinas and Virginia, Charleston, S. C., Feb. 17, 1909.
1. Carcinoma of the Appendix, *Ann. Surg.*, June, 1908.

It has often been noted that cancer in the young is peculiarly likely to be fatal and exceedingly difficult to cure. This is true as a rule, and will readily be appreciated when the cause is either a general loss of resistance on the part of the patient, or the great virulence of the disease, or probably a combination of these two causes. But when that lack of resistance is purely local and due solely to the structural weakness of the appendix, there is no reason to expect any greater degree of malignancy than when cancer develops in persons of more advanced age. On the contrary, if the patient is otherwise of good resistance it seems that the carcinomatous process would be limited by the normal resistance of the other tissues to that portion of the body where the local lack of resistance exists. This is borne out by the fact that, though cancer in the young is usually peculiarly likely to be fatal, in the large majority of cases operations for cancer of the appendix have been followed by no recurrence after a number of years. When cancer originates in an organ whose anatomy and physiology are fixed within definite limits, and whose important functions connect it closely with the rest of the body, it is not a question of a local lack of resistance, but this defect must be shared, to some extent, by other tissues. If the development of the cancer is due to a peculiar virulence of its cells the outlook is grave wherever the cancer originates.

Fig. 1.—Photograph of the tumor before it was opened.

The case reported below is an example of an unusual form of cancer which seems to be prone to originate in the young. It does not proceed with the same rapidity that would be expected from a cancer originating in an important structure in the young. In this instance the origin and course of the cancer may be due to the fact that in the life history of this peculiar cell an unusual change takes place which results in the formation of colloid material within its substance, but not in the surrounding tissues, and after the cell has gained a foothold the colloid which develops as the cell grows lessens its virulence. The tumor, on the whole, presents on palpation and gross inspection the ordinary characteristics of a scirrhus growth, being firm and hard to the touch.

Patient.—Mrs. W. C. W., white, aged 23. Admitted to the hospital Oct. 13, 1906. Discharged, Nov. 15, 1906. Referred by Dr. J. B. Bailey, of Clayville, Va., and Dr. E. W. Gee, of Richmond, Va.

History.—About four weeks before admission to the hospital the patient had been delivered, normally, of a healthy baby. She had suffered with constipation for a number of years. About three days before admission she had severe colicky pains in her abdomen, with absolute obstruction, and gradually increasing tympanites.

First Operation.—A median incision was made. Both large and small intestines were distended. Exploration with the finger disclosed a hard mass involving the lower part of the sigmoid. On account of the great distention, this mass could not be inspected. The patient's condition would not permit of a prolonged operation, so the descending colon was drawn into the wound, fastened with sutures, packed around with gauze, and opened.

She recovered promptly from the operation, discharging large quantities of gas and fecal matter.

Second Operation.—On Oct. 24, 1906, the patient was again operated on under ether, an incision being made in the left rectus, exposing the mass that had been previously felt. It was dissected loose and brought into the wound. The tumor was in the sigmoid, adherent to the broad ligament and abdominal wall, and easily separated. It was two inches in the longitudinal diameter of the bowel and about one inch thick. The tumor replaced the wall of the bowel and almost occluded the lumen. It was hard and firm, the line

Fig. 2.—Photograph of the tumor from the lumen of the bowel after section.

of demarcation with healthy bowel wall at each end was sharp but irregular, and the mucous membrane seemed to be preserved on the lumen side of the tumor. Above and below the tumor, in the wall of the sigmoid, were small nodules that looked like tubercles, pin-head size. Similar nodules were scattered over the mesentery and the peritoneum of the broad ligament. Because the mass was thought tuberculous, on account of the apparently tuberculous nodules and the age of the patient, it was excised. The irregularity of the bowel ends and its vascularity made it advisable to insert two rows of sutures. The inner row of chromic catgut in a round needle secured all bleeding points and approximated the bowel ends. Outside of this was placed a continuous right-angled suture of silk penetrating all coats. Through a stab wound drainage was carried down to the site of the operation and the wound was closed.

The patient made an uneventful recovery. The artificial anus gradually closed without any further operation. A section of the specimen showed that practically all of the bowel wall except the mucous membrane had been destroyed by the growth, which was thick and firm. The caliber of the intestine at this point had been reduced to about the size of a uterine probe. The immediate cause of the acute obstruction was the absolute blocking of this narrow canal by a clove. The patient left the hospital thirty-three days after admission in good condition and rapidly improved in health and weight. For

ten months she did well, but in August, 1907, further symptoms of obstruction developed.

Third operation.—Sept. 20, 1907, she was readmitted to the hospital and again operated on. As had been anticipated, there was recurrence, not only in the bowel wall, but on the parietal peritoneum and all the adjacent tissues. Nodules were scattered along the whole length of the descending colon. It was impossible to excise the neoplasm, so the upper part of the descending colon was brought into the wound and a colostomy performed.

The patient left the hospital after three weeks and returned to her home. She gradually failed and died about three months later.

Microscopic Report.—The essential cell of the neoplasm was one with which I was unacquainted, so the specimen was sent for examination and report to Dr. Joseph Bloodgood, of Baltimore, to whom I am indebted, not only for the following report, but for the photomicrographs which show strikingly the peculiar cell of this growth:

"Section from one of the nodules. There is the dense zone of fibrous tissue, with a zone of tumor tissue to one side, but on the other side of the zone of fibrous tissue

power some cells may be seen to resemble, in a general way, a section of the eye, the clear portion corresponding to the cavity of the eyeball and the eccentric displaced nucleus the lens and cornea. Of course, it will be appreciated that if the cell happens to be cut in this direction the nucleus will be found as described. Those cells whose nuclei appear to be in the center probably have an eccentric nucleus, but, the cell being cut in a different axis, it looks as though the nucleus were central. This appearance is frequently encountered in a spindle-celled tumor when, instead of the cell being cut longitudinally it is cut transversely, which makes the section of the spindle cell look like a round cell.

Aside from its pathologic feature, this case brings to our attention the possibility of cancer of the bowel originating in the young and shows that when dealing with neoplasms of the large intestine we should not exclude carcinoma because of the youth of the patient. As regards prognosis, when cancer in the young is not due to a lack of general resistance in an important organ or to a peculiar virulence of the disease, but

Fig. 3.—Photomicrograph of a characteristic field of the neoplasm herewith reported.

there are irregular areas of tumor cells. The morphology of the cell and its alveolar arrangement is better shown in the smaller areas than in the larger. The cell is of the cuboidal epithelial type of unusual morphology; the majority of the cells contain a material which does not take the stain—colloid or mucoid material. When the cell is entirely filled the nucleus is pushed to one side and becomes eccentric; there are areas where the entire cell is replaced by this material. Even in the larger tumor there is a definite, but thin fibrous-tissue mesh-work enclosing larger and smaller areas of cells. I believe the tumor is a carcinoma with colloid degeneration, a tumor which produces multiple metastases, and has been observed in young people. It can be called carcinoma cubocellulare colloides."

The cell appears edematous. Its nucleus is nearly always eccentric, which seems to be due to the fact that the colloid material accumulated in some portions of the cell more rapidly than in others and so shoved the nucleus to the periphery. On examining the photomicrograph of the higher

Fig. 4.—Photomicrograph of the same field as shown in Figure 3 under higher power. Note eccentric nuclei and dense connective tissue.

merely to the lowered resistance of a vestigial organ, as the appendix, the chances of permanent cure after excision are excellent.

303 West Grace Street.

Human Intestinal Juice.—Salzberg-Faifel (*Arch. internat. de physiol.*, 1908, vi, 81). From a human intestinal fistula relatively normal and pure intestinal juice was obtained. The intestinal juice was markedly alkaline. The alkalinity was due to the presence of alkaline phosphates rather than to carbonates. Expressed in terms of sodium carbonate the alkalinity ranged from 0.35. 0.46 to 0.56 per cent. The average specific gravity was 1.014; the average freezing point, 0.615 degrees C. On chemical analysis no sugars or other substances which reduce Fehling's solution were present. There were no coagulable proteids. The juice was 98.2 per cent. water and 1.02 per cent. non-combustible material. It contained the following ferments: Erepsin, amylase, invertin and enterokinase.