hand, are excreted exclusively by the tubular epithelium. Attempts to separate the functions of the glomeruli and tubules in mammals have not thus far led to decisive results. It is probable, but not proved, that the same conditions prevail as in frogs. There is no conclusive evidence that urea is secreted differently from the salts.

2. *Secretion or Reabsorption by the Tubular Epithelium.*—As stated, the tubular epithelium can secrete all the urinary constituents in the frog. It is highly probable that it secretes pigments in mammals. It is proven that it secretes sugar under the influence of phlorrhizin. There can be no doubt, therefore, that it is capable of secreting some substances, but there is no definite evidence as to whether any of the normal constituents are normally secreted by this epithelium. Reabsorption under normal conditions is neither proved nor disproved; however, its acceptance furnishes the clearest explanation of many phenomena. If it exists it must be selective, discriminating between the different constituents and varying with conditions.

3. *Filtration in the Kidney.*—Filtration through the glomeruli is physically possible, because it occurs in dead kidneys. The influence of saline diuretics, of changes of circulation, etc., on the urine flow corresponds closely in the living and dead kidneys. This indicates that filtration is probably an important factor in the formation of urine, but there is no direct proof. The composition of the urine can not be explained by filtration.

4. *"Secretion" by the Glomeruli.*—It is very probable, though not proved, that glomerular urine may, at times, have a much lower concentration than the plasma. The formation of such a fluid would probably require a force exceeding the available filtration pressure—in other words, it would constitute a secretory process.

5. *The Concentration of the Urine.*—It is very probable, though not proved, that the urine ordinarily becomes more concentrated as it passes through the tubules. At all events, the formation of a urine of greater concentration than the plasma requires an unknown force—whether secretory or absorbing—which must be classed as vital.

6. *The Regulation of the Urine.*—There is no satisfactory explanation of the adaptation of the urine to the needs of the body, none of the proposed theories resting on conclusive proof.

7. *The Action of Diuretics.*—There is no binding evidence as to the mechanism of any of the diuretics, but it is highly probable that salts act physically and mainly on the glomeruli, and that phlorrhizin acts on the tubules. The diuretic actions of water, urea and caffeine are even more obscure.

THE IMPORTANCE OF ICE IN THE PRODUCTION OF TYPHOID FEVER.*

WILLIAM H. PARK, M.D.
NEW YORK CITY.

We may endeavor to settle this question directly by determining whether epidemics or scattered cases of typhoid fever have been traced to ice, or, failing in this, we may try to estimate the probability of such infection by learning the duration of life of the typhoid bacillus after freezing.

The total number of instances of typhoid fever which

have been directly traced to ice infection are remarkably few. I have been unable to trace more than two or three. One was in France, where a group of officers placed ice made from water polluted by a sewer in their wine and afterward developed typhoid fever, while those of the same company not using ice escaped. A second case was in a small epidemic brought to my notice which occurred in those who used ice from a pond. It was found that water directly infected with typhoid feces had flowed over its frozen surface and been congealed there. If typhoid fever is communicated through ice, except under exceptional conditions, it is remarkable that so few cases are traced to it.

The fact that freezing kills a large percentage of typhoid bacilli makes it indeed possible to conceive that ice from moderately infected water contains so few living typhoid bacilli that only the exceptional person here and there becomes infected, and so the source of the infection remains undetected.

If this be true and scattered cases occur, there should be at least some increase on some or every year in March, April and May in such a city as New York, where four-fifths of all the ice consumed is from the Hudson River, which is known to be contaminated with typhoid bacilli. The people of New York use ice very freely and most of them put it directly in their water or place their vegetables on it. The new ice from the Hudson River is gathered in January or February and stored on top of the left-over ice, and thus shipments to the city are immediately begun. It is an established fact that typhoid bacilli in ice are most abundant during the days immediately after freezing. At the end of two months less than 1 per cent. of the original number survive.

If Hudson River ice produced an appreciable amount of typhoid fever, this would then be noticeable in March and in April and perhaps in May.

When we examine the records for the past ten years we find no increase of typhoid fever in Greater New York during those months, with the one exception of the present year, when we had in the Borough of Manhattan a sharp outbreak lasting four weeks. This outbreak did not occur at all in Brooklyn. As the people of Brooklyn drank different water, but received ice from the same places of the Hudson River as those of Manhattan, this directed attention to the water or milk rather than the ice. Examination of the Croton watershed at the time showed that a small epidemic of typhoid existed there and that pollution of the water was probable. This suggested still more strongly that the water and not the ice was the cause of the typhoid infection.

It happened that most of the cases occurred in those living in the section of the upper West Side, where only well-to-do people live. An investigation showed that the majority of the infected had used only artificial ice and several had used no ice in their water at all.

Let us now turn our attention to the life of the typhoid bacillus in ice in laboratory experiments. The first important investigation was that of Prudden, who showed that typhoid bacilli might live for three months or longer in ice. This experiment is frequently wrongly interpreted, as when a recent writer states: "It has been amply demonstrated that the germs of typhoid fever are not killed by freezing and that they have been known to live in ice for long periods of time."

It is true that in Dr. Prudden's experiment a few typhoid bacilli remained alive for three months, when the experiment was terminated, but these were but a small fraction of 1 per cent. of the original number.

* Read in the Section on Hygiene and Sanitary Science of the American Medical Association, at the Fifty-eighth Annual Session, held at Atlantic City, June, 1907.

Following Prudden's experiment Sedgwick and Winslow in Boston and I in New York City carried on independently by a series of experiments. These led to the same conclusions. A table summarizing a final experiment of mine in which twenty-one different strains, mostly of recent isolation, were subjected to the test is given below.

In these experiments twenty-one different flasks of Croton water were inoculated each with a different strain of typhoid bacilli. In one a little of the feces rich in typhoid was directly added. The infected water in each flask was then pipetted into thirty tubes. These tubes were placed in a cold storage room in which the temperature varied from 20 to 28° F. At first tubes were removed and tested twice a week, later once a week. The object of using so many different strains was because it has become evident that some cultures lived longer than others.

LIFE OF TWENTY-ONE STRAINS OF TYPHOID BACILLI IN ICE.

	Average number of bacilli in ice.	Percentage typhoid bacilli living.
Before freezing	2,560,410	100
Frozen three days	1,089,470	42
Frozen seven days	361,136	14
Frozen fourteen days	203,300	8
Frozen twenty-one days	10,280	0.4
Frozen twenty-eight days	4,540	0.17
Frozen five weeks	2,950	0.1
Frozen seven weeks	2,302	0.09
Frozen nine weeks	127	0.005
Frozen sixteen weeks	107	0.004
Frozen twenty-two weeks	0	0

At the end of five weeks the water infected with six cultures was sterile, at the end of sixteen weeks only four strains remained alive.

Two interesting investigations of Hudson River ice have just been completed. One by Dr. D. D. Jackson for the city and the other by Dr. C. E. North for a private laboratory.

There was noticed a considerable difference between the number of bacteria in the top, middle and bottom layers of ice. This is natural, since while water in freezing from above downward markedly purifies itself, 75 per cent. of the solids and a fair proportion of bacteria being eliminated, yet this can not happen in the case of the snow blanket which becomes flooded by rain or by cutting holes through the ice. Here all impurities, such as dust and leaves which have fallen on the surface and dirt which may come from the water, remain with the bacteria which they carry. The bottom ice, being the most recently frozen, has had least destruction by freezing. Dr. Jackson's report gives us a rather unpleasant impression of Hudson River ice. He states that most of the samples either at the top or bottom show intestinal germs. The worst portions of the ice showed from 1 to 10 bacilli of the colon group per c.c. These were taken above or less than five miles below Albany. Some of this ice might aptly be termed solidified sewage.

The analysis of the twenty-five samples shows, as one would expect, that the surface of the ice just at Albany was badly contaminated, one c.c. of melted ice averaging about 15,000 bacteria for four samples. Only one of the twenty-one samples taken at different distances below Albany showed over 1,000 per c.c. A sample from top of a cake of ice obtained two miles below Albany showed only 14. The other twenty samples had on the average 105 per c.c.

Dr. North, in his investigation, examined the ice from forty spots between Hudson and Albany. He took samples from the top, middle and bottom of each cake and the water of the river. He has kindly permitted me to use some of his figures.

The river water in the forty specimens averaged 1,800 bacteria per c.c., the top ice 306, the bottom ice 36 and the middle ice 14. Only four specimens of top ice had over 500 bacteria per c.c., none of the specimens of middle or bottom ice.

Thirty-three of the specimens of water had over 500 and 23 over 1,000. Colon bacilli were obtained from but one specimen of the middle ice, two from the bottom ice and twelve from the top ice.

The great destruction by freezing is noticeable in these figures. Even the top ice soiled by the horses and men gathering it contained but 16 per cent. as many bacteria as the water from which it was contained. The bottom ice, the last to be frozen, had but 2 per cent. of those in the water.

CONCLUSIONS.

The danger from the use of ice produced from polluted water is always much less than the use of the water itself. Every week that the ice is stored the danger becomes less, so that at four weeks it has become as much purified from typhoid bacilli as if subjected to sand filtration. At the end of four months the danger becomes almost negligible, and at the end of six months quite so. The slight danger from freshly cut ice, as well as the natural desire not to put even sterilized frozen sewage in our water, suggests agreement with the report of Dr. Jackson, that portions of rivers greatly contaminated, such as the Hudson within three miles of Albany, should be condemned for harvesting ice for domestic purposes—such ice alone to be used where there is absolutely no contact with food.

ROENTGEN RAY TECHNIC IN DERMATOLOGY.*

MIHRAN K. KASSABIAN, M.D.

Director of the Roentgen Ray Laboratory of Philadelphia Hospital.
PHILADELPHIA.

The value of Roentgen rays in the domain of dermatology has been thoroughly tested and cordially approved. Its place in medicine, as a therapeutic agent, is now established on a sound and rational basis. I have, therefore, attempted to prepare a brief paper on "Roentgen Ray Technic in Dermatology," because the technic, or, if you wish, the art, the skill of application, is the one great prime essential demanded of every operator, if he would attain the very best possible results. So diametrically opposed are the announcements made relative to the value, the uses and the results attained by the Roentgen rays in cutaneous affections, that an explanation for these varying opinions must be sought in the differences in technic employed by the great army of x-ray therapists the world over.

Needless to say, the Roentgenologist must be an adept at his work. The hand of the tyro holds no comparison with the experienced operator in this field of endeavor. In Roentgen diagnosis differences in technic result in excellent or defective skiagrams. What is true here holds equally well in the therapeutic application of the rays. In order to employ this agent so that satisfactory results may be obtained, it is primarily essential that its use as a diagnostic means should be perfectly mastered, and that the operator have more than an ordinary knowledge of the principles involved and likewise an

* Read in the Section on Cutaneous Medicine and Surgery of the American Medical Association, at the Fifty-eighth Annual Session, held at Atlantic City, June, 1907.