H y g e i a.

Set.	1851	Greenw. M. T.	R. A.	Exc. of Eph.	N. P. D.	Exc. of Eph.	No. of Comps in	
							R. A.	N. P. D.
1	Sept. 13	12 ^h 38 ^m 16 ^s .8	0 ^h 20 ^m 42 ^s .75	—1 ^s .59	82° 8' 44".5	+ 8".9	24	8
2	15	12 19 27,4	19 20,32	1,43	16 9,0	7,0	12	4
3	21	13 16 10,4	15 1,16	1,42	82 40 32,8	10,2	24	8
4	27	13 3 23,2	10 37,20	1,02	83 7 12,7	7,8	18	8
5	28	10 52 33,3	9 57,14	0,93	11 22,6	9,2	18	6
6	Octb. 2	12 1 57,6	7 1,03	1,07	30 19,6	8,4	6	2
7	6	9 54 52,1	4 15,14	0,94	48 48,4	11,5	9	3
8	7	10 30 42,8	3 32,65	0,81	53 41,0	10,2	24	8
9	7	11 55 57,7	3 30,67	1,27	53 57,7	10,3	21	7
10	8	10 16 27,3	2 52,50	1,13	83 58 21,5	10,6	16	6
11	10	12 43 30,5	1 28,03	0,84	84 8 16,7	9,2	24	8
12	12	10 50 4,5	0 0 13,27	—0,64	84 17 16,3	+ 7,2	24	8

Parallax and computed places taken from Suppl. Naut. Alm. 1854.

Mean places of Stars of comparison 1851,0.

Set.	Name.	R. A.	N. P. D.
1	Weisse 0 ^h 318	0 ^h 19 ^m 13 ^s .49	82° 15' 48".4
2	— 0 ^h 375	22 50,14	18 42,3
3	Greenw. T. Y. C. 16	12 55,98	82 38 15,3
4	Weisse 0 ^h 169	10 1,75	83 4 7,2
5	Greenw. T. Y. C. 22	18 1,11	7 58,7
6	Weisse 0 ^h 192	11 27,89	32 45,1
7	— 0 97	6 4,79	53 20,0
8	— 0 89	0 5 22,91	55 59,6
9	— 23 1222	23 59 26,52	83 57 9,9
10	— 23 1252	0 0 53,74	84 2 47,2
11 and 12	— 23 1251	0 0 46,54	84 12 46,6

N o t e s.

Hygeia appeared as a star of 10.11 magnitude.

Sept. 15 and Oct. 2. Interrupted by fog.

Oct. 6. Interrupted by clouds.

Oct. 10. Observed through thin clouds.

R. C. Carrington.

Beobachtungen auf der Liverpooler Sternwarte, von Herrn *J. Hartnup*.

Observatory, Liverpool 1851, November 6.

On the 15th of September last we had here a very fine state of atmosphere for astronomical observation. On pointing the 8½ inch. achromatic refractor to Saturn, I was struck with the great beauty of the image, and the new ring obtruded itself upon my notice immediately. It was certainly much broader than represented by Professor *Bond* and Mr. *Daves* at the time of discovery, extending over full half, or from half to two thirds of the space between the inner side of the inner bright ring and the ball of the planet. The obscure ring was well defined and equally well seen on both sides of the ball of the planet, but it was fainter near the inner edge than it was near its junction with the bright ring. The bright

and obscure rings, where they crossed the ball of the planet, were both beautifully defined, the primary division in the former was seen as a fine black line projected on the face of the planet. I never saw the primary division so perfect before. I examined the planet for three hours with powers from 170 to 1100 with all of which the new ring was distinctly seen, but I could detect no other than the primary division in the bright ring.

The following observations of Irene, *Brorsen's Comet*, Hebe, Iris and Hygeia, I have obtained since my last communication.

I r e n e.

Liverpool.

Equatoreal.

(Mr. Hartnup).

		Greenw. M. T.	R. A.	Log. $\frac{P}{P}$	N. P. D.	Log. $\frac{q}{P}$	Star of Comparison.
1851	June 10	12 ^h 34 ^m 40 ^s .4	15 ^h 43 ^m 42 ^s .60	+8,292	104° 12' 31".3	-9,9559	B. A. C. 5134
	—	12 58 36,8	43 41,86	8,367	12 23,2	9,9520	— —
	—	12 22 33,2	43 40,90	8,426	12 28,5	9,9475	— —
	15	10 29 46,2	40 14,12	7,333	28 10,2	9,9651	— —
	—	10 54 42,5	40 13,56	7,820	28 13,7	9,9642	— —
	—	11 19 38,4	40 12,79	+7,041	28 16,6	9,9628	— —
	16	9 55 17,2	39 37,18	-7,509	31 31,7	9,9652	— —
	—	10 20 13,1	39 36,24	+7,099	31 35,7	9,9653	— —
	—	10 45 9,2	39 35,65	7,758	31 39,4	9,9648	— —
	17	10 11 42,0	39 0,00	6,731	35 6,7	9,9636	— —
	—	11 1 34,2	38 58,66	7,975	35 14,4	9,9636	— —
	July 16	9 55 10,3	33 41,92	8,269	106 46 0,1	9,9634	— 5257
	—	10 37 5,8	33 42,08	8,398	46 7,8	9,9556	— —
	17	9 59 27,7	33 56,47	8,298	51 20,6	9,9624	— —
	—	10 19 25,3	33 56,43	8,361	51 23,7	9,9587	— —
	18	10 19 46,1	34 12,82	8,373	56 47,2	9,9580	— —
	—	10 39 44,5	34 13,16	8,423	56 47,8	9,9536	— —
	21	10 12 55,2	35 11,14	8,381	107 13 8,0	9,9581	— —
	—	10 32 53,9	35 11,46	8,430	13 13,2	9,9536	— —
	Aug. 1	9 57 7,6	40 41,53	8,439	108 15 22,2	9,9543	— 5176
	—	10 13 7,5	15 40 41,71	+8,472	108 15 25,7	-9,9486	— —

The observations are corrected for refraction. The corrections to be applied for parallax in time and arc, are represented by p and q . P is the Equatoreal horizontal parallax.

The following are the assumed mean places of the stars of Comparison for 1851,0 derived from the Greenwich 12-year Catalogue.

	R. A.	N. P. D.
B. A. C. 5134	15 ^h 27 ^m 11 ^s .83	104° 17' 19".49
— 5257	15 45 20,87	106 17 15,98
— 5176	15 33 22,20	109 11 30,54

Brorsen's Comet.

	Greenw. M. T.	R. A.	Log. $\frac{P}{P}$	N. P. D.	Log. $\frac{q}{P}$	Star of Comparison.
1851 Aug. 13	10 ^h 18 ^m 21 ^s .4	14 ^h 17 ^m 16 ^s .87	+8,690	53° 1' 35".4	-9,7558	B. A. C. 4812
—	10 38 20,3	17 18,29	8,695	1 18,3	9,7788	— —
—	10 58 19,5	17 20,21	8,698	5 52,4	9,8007	— —
15	10 24 47,4	21 37,28	8,699	52 4 26,8	9,7605	— —
—	10 39 46,4	21 38,28	8,702	4 14,0	9,7781	— —
—	10 54 45,5	21 39,49	8,703	3 53,7	9,7950	— —
25	9 14 18,0	46 23,59	8,700	47 11 56,5	9,6346	— 4958
—	9 34 17,4	46 25,45	8,714	11 31,8	9,6667	— —
26	9 49 11,8	49 18,85	8,726	46 40 56,7	9,6831	— —
—	10 4 11,5	49 20,68	8,732	40 40,6	9,7093	— —
—	10 19 10,8	49 21,93	8,736	40 17,0	9,7312	— —
Sept. 6	9 53 59,5	15 27 29,46	8,775	40 58 59,6	9,6416	— 5177
—	10 14 0,8	15 27 33,55	+8,781	58 38,1	-9,6784	— —

The observations are corrected for refraction.

The following are the assumed mean places of the stars of Comparison for 1851,0.

	R. A.	N. P. D.	Authority.
B. A. C. 4812	14 ^h 26 ^m 4 ^s .63	51° 2' 16".13	Greenwich 12-year Catalogue.
— 4958	14 56 20,07	49 1 9,78	— —
— 5177	15 33 30,00	42 42 32,01	Oxford and Edinburgh Observations.

Liverpool.

H e b e.
Equatoreal.

(Mr. Hartnup).

		Greenw. M. T.	R. A.	N. P. D.	Computed — Observed.		Star of Comp.
					R. A.	N. P. D.	
1851 July	17	11 ^h 14 ^m 51 ^s .6	19 ^h 13 ^m 37 ^s .57	99 ^o 58' 45 ^u .1	—1 ^s .15	—2 ^u .9	B. A. C. 6564
—	—	11 34 48,0	36,86	51,3	—1,24	—1,6	—
18	18	11 19 57,3	19 12 39,55	100 7 53,2	—1,32	—1,8	—
—	—	11 39 53,4	12 38,72	59,5	—1,29	—0,4	—

The observations are corrected for refraction and parallax. The computed places were taken from the Ephemeris contained in the supplement to the Nautical Almanac for 1854. The following is the assumed mean place of the star of comparison for 1851,0 derived from the Greenwich and Edinburgh Observations.

B. A. C. 6564 19^h 14^m 35^s.76 R. A. 98^o 11' 1^u.55 N. P. D.

I r i s.

		Greenw. M. T.	R. A.	N. P. D.	Comp. — Obs.		Stars of Comp.
					R. A.	N. P. D.	
1851 Sept.	14	11 ^h 16 ^m 0 ^s .8	0 ^h 1 ^m 24 ^s .04	77 ^o 45' 43 ^u .8	+1 ^s .62	—5 ^u .7	B. A. C. 26 and 8331
—	—	11 57 53,7	0 1 22,98	77 45 49,1	1,36	6,2	—
15	15	9 57 31,9	0 0 40,60	77 49 1,4	1,68	7,0	—
—	—	10 37 25,1	0 0 39,54	77 49 6,6	1,45	6,2	—
20	20	12 11 25,3	23 56 36,19	78 10 47,5	+1,48	—7,2	B. A. C. 26.

The observed places have been corrected for parallax and refraction. The computed places were deduced from the Ephemeris contained in the supplement to the Nautical Almanac for 1854.

The following are the assumed mean places of the stars of comparison for 1851,0 derived from the Greenw. Observations.

B. A. C. 26 0^h 5^m 34^s.05 R. A. 75^o 38' 41^u.02 N. P. D.
 ——— 8331 23 51 39,72 „ 83 57 41,82 „

H y g e i a.

		Greenw. M. T.	R. A.	N. P. D.	Comp. — Obs.		Stars of Comp.
					R. A.	N. P. D.	
1851 Sept.	14	11 ^h 34 ^m 36 ^s .9	—0 ^h 20 ^m 3 ^s .39	—82 ^o 12' 11 ^u .2	—1 ^s .57	+12 ^u .3	B. A. C. 26 and 8331
—	—	12 16 29,9	20 2,28	12 18,9	1,65	11,1	—
15	15	10 16 11,9	19 23,97	15 45,7	1,50	10,7	—
—	—	10 56 5,1	—0 19 22,83	—82 15 50,6	—1,51	+12,1	—

The assumed places of the stars of comparison are the same as given for Iris. The observed places have been corrected for refraction and parallax. The computed places were deduced from the Ephemeris contained in the supplement to the Nautical Almanac for 1854.

J. Hartnup.

Beobachtung des von Herrn *Brorsen* 1851 October 22 entdeckten Cometen,
auf der Berliner Sternwarte von Herrn *Luther*.

	M. Zt. Berlin.	AR. ☿	Decl. ☿	Vergl.
1851 Nov. 11	6 ^h 38' 3 ^u .2	227 ^o 44' 53 ^u .5	+43 ^o 17' 43 ^u .9	6
Angenommener scheinbarer Ort des Vergleichsterns:				
		227 ^o 16' 24 ^u .1	+43 ^o 9' 5 ^u .6	