meter, though embedded in the mass of ice, did not really indicate the true temperature of the letter. With the object, therefore, of proving whether the ice is hot or not, I have, at the suggestion of Professor Roscoe, made the following calorimetric determination:

The arrangement of the apparatus was so modified that the ice, after being strongly heated, could be suddenly dropped into a calorimeter containing a known quantity of water of known temperature. The resulting temperature, after the ice had been dropped in, was read off by a thermometer graduated so as to indicate a difference of 0.05° C. The weight of the ice was found by re-weighing the calorimeter.

So far I have only had the opportunity of completing the two following determinations, and in the second of these the weight of the ice could not be found, as a small quantity of water was lost out of the calorimeter, owing to a sudden jerk at the moment the ice entered it:

- (1.) Weight of water in calorimeter, including the value of the latter=185 grammes.

Weight of ice dropped in=1.3 grammes.

Temperature of calorimeter before=13.4°

after = 13.6°

Rise in temperature=0.2°

$M(\theta - t) + 80W = W(T - \theta)$

$(185 \times 0.2) + (80 \times 1.3) = 1.3(T - 13.6)$

$\therefore T = 122^\circ \text{C.}$ Where T =temperature of ice.

- (2.) Weight of water in calorimeter, etc.=185 grammes.

Temperature of calorimeter before=12.7°

after = 12.8°

Rise in temperature=0.1°

On weighing the calorimeter after the experiment, the increase in weight was only 0.15 gramme, but as a portion of the water had been jerked out during the operation the true weight of the ice, and therefore its temperature, could not be found. But since the calorimeter had slightly risen in temperature, the ice must have been above 80° C.

From the nature of the experiment, as carried out on the present scale, the weight of the ice which can be dropped into the calorimeter is only small, and therefore the rise in temperature is but slight. But since a fall in temperature of a much larger amount ought to have been obtained had the ice been at 0°, it is considered that the above experiments are conclusive. Great care was taken in order to obtain correct temperatures in the calorimeter. The latter was inclosed in several casings, and the water was allowed to stand in it for several hours before the experiment, so that it might first attain the temperature of the room, while the time which elapsed between the readings of the thermometer before and after the ice was dropped in would not be more than from ten to fifteen seconds.

In the course of the next few weeks I intend to make one or two more determinations, and, if possible, on a larger scale.

ADULTERATION OF FOODS.

ONE of the most interesting recent contributions to this branch is a report on an investigation to determine the prevalence of adulteration in food supplies, undertaken, in accordance with instructions from the National Board of Health, by Dr. Smart.

Tea.—Dr. Smart does not consider the so called "facing" of teas with Prussian blue and silicates as injurious. Mixing teas is very common in this country, but it is not harmful to health. His general results were as follows:

The teas examined numbered 117, but of these two which were grown in Georgia, and six sent specially from Japan, may be excluded from present consideration. Of the 109 remaining, 90 were obtained from sources which presumably should have furnished a pure article, while 19 were purchased at such stores as might be expected to furnish an adulterated article, if adulteration was at all prevalent. In no case was a leaf observed which was not a true tea leaf. Of the 90 the only cases of debasement were those already mentioned as having probably originated before importation, while 5 of the 19 were so deteriorated that an analyst would have been warranted in reporting them as fraudulently adulterated—one, gunpowder (Y 2), from an excess of lime sulphate facing; one imperial (W 2), from admixture with sand, and three, imperials (Y 1, 2, and 3), from the presence of exhausted leaves. Whether these debasements were effected in China or in this country is an open question; but from the character of the English importations it is probable that exhausted leaves are dried and recolored in this country, for sale among the poorer classes of the community.

Coffee.—In this popular beverage something more positive was reported; but, on the whole, the examination did not yield serious results. The investigator says:

Most of the dealers with whom I have conversed have been strong in their belief that adulterated coffee is not to be found in this country, as the beans are either home-ground or ground by the supplying grocer. The results of H. B. Hill's examination for the State Board of Health of Massachusetts sustains this opinion, as ten samples of ground coffee, obtained in bulk, were found to be pure, while seven package samples consisted of one with no coffee, two with very little, and four with from fifty to seventy per cent. of coffee, the rest being roasted wheat, peas, beans, and chicory. But the supplying grocer is not to be trusted in all cases, as out of the few purchased samples of loose ground coffee which have been examined during the present investigation, one contained a small percentage of chicory, while two consisted largely of chicory and roasted beans. Only one sample of package coffee was received, and it sustained the accuracy of Dr. Hill's results. Chicory, corn, wheat, and rye were noted on the microscopic field as adulterants; but adulterants of what? for no coffee had been discovered, and half a dozen successive slides carefully examined failed to show the presence of the nominal article.

Three extracts or essences of coffee were examined, one of which consisted of chicory and the two others of roasted starches. Two of these bore on the label an offer of a reward for proof that any other extract of coffee was as pure as they, which does not say much in a general way for the purity of extract of coffee. Two samples of chicory were examined, both of which were found pure.

Sugar.—Cane sugar seems widely adulterated by glucose manufactured from starch. On this point the result was as follows:

Among 47 brown sugars, most of which were furnished by dealers who knew that their samples would be examined, there were found 3 which contained glucose, while among 38 samples purchased for analysis no less than 9 were thus

adulterated. The glucose varied from a small admixture of thirty per cent. A remark by one of the dealers who furnished a thirty per cent. sugar is worthy of mention in this place: "Oh! they all have it, but they don't send any of it to you." Another storekeeper from whom a degraded sugar was obtained remarked, as he was putting up the samples, that I was welcome to anything he had, as he did not suppose he was any worse than his neighbors.

The white sugars, powdered, granulated, etc., were free from this adulteration; 24 of them were furnished dealers and 15 were purchased. The sirups, of which 21 were sent in for examination and 12 purchased, were also pure.

Flour.—This important staple furnishes gratifying results. Of 58 samples examined, only 1 had any adulteration, and this was merely corn meal, probably introduced by accident. The same cannot be said of bread.

Bread.—Of 18 samples, 8 contained alum. Although it is by no means decided that the substance, as used by bakers, is hurtful, it is advised that it should be prohibited.

Cream of Tartar.—This substance is one of the articles which are subject to gross adulteration. Only 18 samples were examined, but these were considered to be enough, in view of the character of the results. Six were of satisfactory purity. Eleven contained sulphate of lime, varying from seventeen to ninety per cent., 3 having nearly the latter figure. Two contained no cream of tartar, but consisted instead, the one of sulphate of lime, alum, and acid phosphate of lime, and the other of alum, acid phosphate, and potato starch. Corn starch was also found in large proportion in one of the lime sulphate powders. Considering the use of cream of tartar in baking, its impure condition is a serious evil.

Baking Powders.—Contained alum and sulphate of lime, and were often decidedly impure.

Ground Pepper.—As ground spices in general are quite impure, Dr. Smart observes:

The examination of the ground black peppers and spices shows to what extent adulteration may be practiced when its detection by the public is a matter of difficulty. The dealer himself appears to have lost the knowledge of the characters of the pure article, as out of 4 samples sent in by respectable houses in this city for the purpose of being examined, only 1 was pure. The others contained baked flour and rice with sand enough to prove the unclean condition of the pepper itself.

Out of 18 samples of ground cayenne pepper only 2 were pure. Out of 23 specimens of commercial ground mustard, 21 were adulterated. Out of 36 samples of ground cinnamon, only one was the genuine article! The usual adulterants are colored clays, roasted wheat, beans, rice, sawdust, etc., but, fortunately, no red lead.

In confectionery the red colors are generally harmless cochineal, but the yellows are often from salts of lead. Some yellow lemon balls yielded seven grains of lead chromate to the pound of candy.

THE LIQUEFACTION OF OZONE AND ITS COLOR IN THE GASEOUS STATE.

By MM. HAUTEFEUILLE and J. CHAPPUIS.

MM. HAUTEFEUILLE and J. CHAPPUIS have recently published some notes on this subject, which supply interesting information. Ozone, such as one usually prepares, possesses so little tension in oxygen, at most 53 mm., that the physical properties of this body are hardly known and distinguished from those of oxygen. The difficulties which Soret has so ably overcome in determining the density of ozone by operating on weakly ozonized oxygen are known. One of the fixed physical properties of this body is its heat of formation, which was accurately determined by Berthelot, in spite of the state of dilution in which it is found on leaving the apparatus for ozonization. The preparation of a mixture very rich in ozone is then the first condition to be fulfilled, in order to acquire new notions of this curious body. We have already established the fact that the isomeric transformation of oxygen submitted to electricity obeys simple laws, and that the proportion of ozone increases but very little with the pressure for every temperature, whereas, in passing from 20° to -55°, the proportion of ozone increases fivefold. Removed from the action of electrical discharges, the mixture of oxygen and ozone ceases to be a homogeneous and balanced system; in spite of that, the mixture is preserved without appreciable alteration as long as a constant temperature is maintained, if we operate on it below 0°. This relative stability of ozone allows us to compress the mixture, and to obtain the tensions of ozone of several atmospheres. As it is necessary to prepare ozone, destined for these experiments, under the strongest possible tension, one must ozonize oxygen at a very low temperature. Consequently the oxygen remains a quarter of an hour in an apparatus of alternating discharges, whose concentric tubes of thin glass are dipped into chloride of methyl, then one passes it into a test tube, terminated by a capillary tube of the Cailletet apparatus. This reservoir, of about 63 c.c., at first empty and kept at -23°, not being able to fill itself at one time under a pressure approaching 760 mm., is put rapidly in communication, five times successively, with the electric apparatus, the capacity of which cannot exceed 20 c.c. In a quarter of an hour one succeeds by this method in filling the test tube with a mixture of oxygen and ozone, very fully charged with the latter gas. The test tube is then taken away from the chloride of methyl and separated from the electric apparatus, and the gas which it contains is driven slowly back by mercury, cooled to 0° in a capillary tube maintained at -23°. The mercury which transmits the pressure of the hydraulic press does not impoverish the gaseous mixture as quickly as might be feared; a solid glaze is formed on the surface of the metal, which rapidly limits the action; the heating of the gas during the compression is more formidable. In spite of these difficulties one succeeds in increasing the tension of the ozone to a considerable degree. From the first working of the piston, the capillary tube becomes of an azure blue; this coloration is intensified in proportion as we reduce the volume of gas; and if the tension of the ozone is brought by the compression to be one of several atmospheres the gas is indigo blue, and the meniscus of mercury, seen through the gas, is then of a steel blue. The blue color of the gas becomes less intense, and the mercury regains its usual metallic appearance as soon as the tension of the ozone diminishes. The preceding mixture contains enough ozone for one to observe a thick white mist at the moment of the expansion which succeeds a compression of 75 atmospheres. It is not then necessary to compress the ozonized oxygen as much as the pure oxygen—300 atmospheres—in order to determine by a sudden expansion the momentary

formation of a mist, a certain sign of liquefaction or even of solidification.

A comparative study of the mixtures of oxygen and ozone, and those of oxygen and carbonic acid, shows that in very comparable conditions the expansion ought to be sensibly stronger with the ozone than with the carbonic acid for one to begin to perceive a mist. The ozone would then be a little easier to liquefy than the carbonic acid. The mixture of oxygen and ozone contains an explosive gas, and should always be compressed slowly and cooled; for if one does not meet these conditions the ozone decomposes with the escape of heat and light, and there is a loud report accompanied by a flash of yellowish light. Berthelot has found that the transformation of oxygen into ozone absorbs 14 cal. equivalent to 8 (0°=24 gr.); the ozone may then be placed beside explosive gases; our experiments show that, like them, this body is susceptible to sudden decomposition.

One may also observe a portion of these new facts by compressing the oxygen which has gone slowly through the electric apparatus at ordinary temperature; for if we compress this gas rapidly in a capillary tube, placed in some water at 25°, we often destroy the ozone with explosion, but if this same gas is cooled at -23°, the ozone which it contains may be brought to a tension of 10 atmospheres, and may be preserved for hours in these conditions of temperature and pressure, if the gas is separated from the mercury by a column of sulphuric acid.

It is ascertained then almost equally clearly as in the preceding experiment that the ozone is a gas of a beautiful azure blue, for its color is so intense when we increase its density tenfold, that we have been able to see it in a tube of 0.001 m. internal diameter, when we were operating in a very poorly lighted room in the laboratory of l'Ecole Normale. It is then established that ozone under a strong tension is a colored gas—but may the same be said of ozone at the tension of some millimeters? The blue color characterizes the ozone as surely as its smell, for at all the tensions we find it on examining the gas to a sufficient depth. It suffices in order to make this manifest to interpose between the eye and a white surface a tube of 1 m. in length, traversed by the current which has passed through the electric apparatus of Berthelot. The color which the gas then possesses reminds one of the blue color of the sky. This blue is more or less deep, in proportion to the time the oxygen has remained in the apparatus, and it is consequently more or less rich in ozone. As soon as we stop the electric tube the coloration disappears, the ozonized oxygen being replaced by pure oxygen.

The following numbers, from a paper by the same authors in *Comp. Rend.*, are of interest, as showing the exact influence of temperature and pressure on the ozonizing of oxygen. Diminution of pressure does not tend to increase the amount of ozone produced, but decreased temperature exerts a marked action in increasing the amount of oxygen transformed into ozone:

Tension of oxygen.	Tension of ozone.				Proportion of ozone by weight.			
	-23°.	0°.	20°.	100°.	-23°.	0°.	20°.	100°.
760	108.70	82.64	53.96	—	0.214	0.149	0.106	—
380	51.68	38.76	31.54	1.48	0.204	0.152	0.125	0.0117
300	40.20	30.60	22.20	—	0.201	0.1525	0.112	—
225	24.80	22.95	15.52	0.088	0.191	0.153	0.104	0.0118
180	22.30	16.58	10.52	—	0.181	0.137	0.089	—

THE OXIDATION OF ALCOHOL.

WHEN a dilute or alcoholic fluid is exposed to the action of free oxygen, the latter gradually combines with the former, and according to the generally received opinion, acetic acid is the product of the change. The formation of acetic acid may be expressed in chemical notation as follows:

This oxidation is brought about through the influence of a certain ferment called *Aceti mycoderma*, but other organisms are also capable of inducing the same change. Acetic acid, however, can be formed from alcohol without the agency of organized ferments. Doberiner's celebrated experiment proved that the vapor of alcohol was converted into acetic acid by the agency of porous platinum, which possesses the remarkable property of condensing gases upon its surface, and thus the alcohol and oxygen are brought into sufficiently close contact to combine. Other porous materials besides spongy platinum possess this power, although, perhaps, not to the same degree; but it is right to mention that, according to Pasteur, the oxidation of alcohol into acetic acid is due to the intervention of one of the oxidizing ferments. The cells of these organisms probably act much in the same manner as the spongy platinum—namely, by condensing the oxygen within them, and thus, by rendering it more concentrated, making its energy greater. In a dilute alcoholic fluid like beer, we have all the conditions favorable to the production of acetic acid, and the change is undoubtedly hastened by the porous character of the wooden vessels in which beer is usually stored. Even if the pores of the wood do not possess the power of converting alcohol into acetic acid, which Liebig and many other chemists claimed for them, they at least serve as efficient carriers of the minute organisms which, according to Pasteur, are the principal agents in the phenomena of acetification. In drawing attention to the changes in the oxidation of alcohol, which have so much interest for brewers on account of their disastrous effects on beer, it is necessary to point out that there is an intermediate product. Midway between alcohol and acetic acid there is a substance called aldehyde; as its name implies, it is alcohol deprived of hydrogen, and its formation and subsequent conversion into acetic acid is explained by the following chemical equation:

It will be seen that the change is really an oxidation of the alcohol, although the oxygen acts by removing a portion of the hydrogen. The subsequent conversion of aldehyde into acetic acid by means of another equivalent of oxygen is expressed by the following equation:

When the vapor of alcohol is oxidized by means of spongy platinum or other powerful oxidizing agents, aldehyde is formed in large quantities, and it is more than probable that the cells of the oxidizing ferments act in a similar manner. In this way aldehyde may often be formed in beer, and as aldehyde possesses a peculiar fruity odor and is very volatile, this body may play an important part in developing the characteristic flavors and aromas of different kinds of beer. The detection of aldehyde in beer is attended with considerable difficulty, but several investigators have identified it as