a great disadvantage, because none of the hospitals issues a printed annual report, and I did not succeed in obtaining a printed copy of the schedule of lectures and clinics of the medical college. The total absence of such printed publications is only another evidence of the present poverty-stricken condition of the seaside republic of South America.

SANTIAGO DE CHILE, August 25.

Clinical Notes

PNEUMATIC SCROTAL COMPRESSOR FOR USE IN EPIDIDYMITIS.

EDGAR BALLENGER, M.D.

Lecturer on Genito-Urinary Diseases, Atlanta School of Medicine.
ATLANTA, GA.

Generally speaking, the chief factors desired in the routine treatment of epididymitis are rest, cold or heat, compression, and suspension. The value of compression, especially in the latter part of the disease, is admitted by the best authorities and is attested by the various methods that have been advocated with this purpose in view, such as strapping with adhesive plaster, the appli-

Scrotal compressor showing attachments for either hydraulic or pneumatic pressure.

cation of thin rubber bandages and the use of a large number of suspensories.

An attempt to utilize any one of these plans of treatment will readily demonstrate their serious disadvantages. Strapping with adhesive plaster works well for the first twenty-four hours until the swelling has yielded to the compression, then no further benefit can be obtained without reapplying it. The manipulation and handling required for this procedure usually causes sufficient irritation to counteract the good effect of the compression. The plaster also frequently irritates the skin. So much skill and judgment is necessary in order to bandage the swollen part with thin rubber that the average practitioner finds it very unsatisfactory. It is either applied so loosely that it will not hold and no benefit is obtained, or it is too tight and has to be removed to relieve the pain. The suspensories on the market are still more unsatisfactory as they do not furnish uniform pressure.

I have devised a pneumatic compressor which is sim-

ple, easily applied and effective in maintaining uniform compression of any degree, and I think it is the most satisfactory method so far advocated for the treatment of epididymitis. It is composed of two layers of rubber, the outer one is inelastic while the inner is elastic. The space between them is to be inflated with air after the compress has been applied and laced around the swollen part.

The pressure can be accurately regulated by gradually forcing the air in or allowing it to escape. As the swelling subsides more air should be injected to maintain the desired pressure. A commendable feature is that this can be done without any manipulation to aggravate the condition. A light gauze stocking should be placed under the compress to absorb the perspiration. By using this method of treatment patients can be gotten out of bed more promptly and with little danger of a relapse.

Local medication in the form of guaiacol in from 10 to 30 per cent. solution in glycerin or tincture of iodine painted on the swollen part once or twice daily may be used in conjunction with the compression. These I have found to be the most useful local remedies.

The larger size compressors are supplied with an exit valve for drainage, and instead of using an ice bag in the beginning of the disease a fountain syringe containing ice water (or hot water if preferred) may be connected to furnish hydraulic pressure in the place of air. By regulating the height of the bag and the rate of outflow both compression and cold can be adjusted to suit the comfort of the patient and the lessened swelling. It is well known that the benefit of an ice bag can be augmented by pressure to force some of the blood out of the tissues and prevent the too rapid dissipation of the cold.

The compresses are manufactured in several sizes, and the smallest one that can be used with comfort should be selected and snugly laced before being inflated. They are attached by straps and held in place as an ordinary suspensory to prevent any tension on the cord. Their simplicity, with the admitted value of uniform pressure which can be maintained and easily regulated or combined with cold or heat, make unnecessary any further exposition of the advantages to be derived from this plan of treatment.

1014 Century Building.

THE HOSPITAL ARRANGEMENTS OF A MAN-OF-WAR.

ROBERT A. BACHMANN, M.D.

Passed Assistant Surgeon, U. S. N.; Senior Medical Officer, U. S. S. St. Louis.

There was a time when the only place devoted to the sick in a man-of-war was a small, dark, stuffy place situated on one of the lower decks, usually forward in the very bow. In a way these sick quarters were commensurate with the demands of the profession at that time. There was little antisepsis and no asepsis in surgical work, laboratory methods were confined to small limits and the dispensary was not called on for the variety of medication as at present.

The importance of a well-equipped medical department on board of a man-of-war, too, was less apparent and the value of hygiene in war not known. So long did these inferior conditions exist that even now the impression prevails in many minds that on board a man-of-war little or no improvement has taken place; at least I

doubt if the modern, greatly improved hospital arrangements are as well known as they deserve to be.

The U. S. S. *St. Louis*, while not an example of our largest class of war vessels—the battle ships—being rated as a protected cruiser, having less tonnage than the former, by from 2,000 to 6,000 tons, and greater speed, still is a fine type of ship for the purpose of this article. Its length is 426 feet, extreme beam 66 feet and displacement 9,700 tons.

In order to afford the greatest safety to the sick the sick quarters on men-of-war are usually located on one of the lower decks behind armor or coal bunker protection. On the *St. Louis* the sick bay is situated on the berth deck forward, but as the ship is not armored, enjoys no greater immunity to shell fire than any other part of the ship. In close connection on the same deck are the operating room and the dispensary. Farther forward is the isolation ward.

The sick bay (Fig. 1) is a large and well-lighted room containing twenty-four folding

Fig. 1.—Sick-bay looking into operating room.

bunks arranged in pairs one above the other. The air contents are sufficient to allow about 220 cubic feet per bed. Four large supply louvres (air shafts) furnish a steady influx of fresh air, and are constructed so that the current may be directed to any part of the ward.

Electric fans, port holes, etc., furnish additional aids to the ventilation. The deck of the ward is covered with thick linoleum, which is shellacked at regular intervals, making it a smooth waterproof surface easily kept disinfected, and comparatively noiseless to the tread. A bath room with mosaic tiled floor, porcelain tub, shower and modern sanitary fittings adjoins the sick bay. In all the washstands of the ward, toilet rooms, operating room and dispensary nothing but distilled water flows through the pipes. Part

of the space has been utilized as a dressing station, where minor surgery is done and the various dressings attended to.

The operating room (Fig. 2) is directly across from

Fig. 2.—A view of a portion of the operating room.

the sick bay, a small passage extending between them, and the two take up the entire width of the ship—about 50 feet at this point. It is equipped with an enamel operating table and electric reflector for night operations, instrument cabinet, autoclave, small steam sterilizer for instruments, basin stands, dressing rack, etc.

The supply of instruments is amply sufficient for all the usual major and most of the special regional operations. The sides of the room are of steel covered with white enamel paint and the floor is mosaic tiling.

The dispensary (Fig. 3) is located just forward of the operating room, to which it is nearly equal in size, and part of it is arranged for laboratory work. A microscope with mechanical stage, the most frequently used stains, sterile typhoid cultures, a Zeiss hemacytometer, centrifuge, Talquist's hemoglobinometer, etc., and a chemical case containing ureometer, albuminometer, Erlenmeyer flasks, burettes, pipettes, tubing, fermentation tubes, Petri dishes, graduated cylinders, etc.; in short, all apparatus and chemicals the medical officer might require for a bacteriologic or chemical diagnosis are supplied.

The isolation ward, still farther forward, is a small compartment, entirely closed in and containing four folding bunks. The deck is of mosaic tiling and the sides enameled steel. It is devoid of all furniture save

Fig. 3.—Dispensary and laboratory.

the metal bunks, and can be rapidly disinfected by either fumigation or solutions. The air supply here is entirely from leakage and one port hole, facilitated by an exhaust blower which evacuates the air and conducts it to the exterior at the stern of the ship.

On the superstructure of the ship a tent (Fig. 4) can be pitched for additional cases in case of an epidemic and at other times used for phthisis and pneumonia cases.

There are aboard about 650 men including the

Fig. 4.—Tent for the treatment of tuberculosis and pneumonia, located on the after bridge.

officers. For this complement there are one hospital steward, two hospital apprentices, first class, and one hospital apprentice (Fig. 5).

Hospital stewards are usually graduates of a college of pharmacy or have equivalent experience; hospital apprentices, first class, are graduates of a naval training school in most cases and have passed an examination for the grade. The hospital apprentices are recruited from civil life and need merely possess a general education. Besides having a knowledge of their profession,

Fig. 5.—Trained hospital corps.

these men are drilled in the special requirements of a military organization.

In this connection it might be well to add that no little part of the medical officer's time and attention are given to working out preparations for conditions in the result of actual warfare. First aid instruction is given to every man on the ship. To some showing adaptability more advanced training is allowed for the purpose of

using them in time of battle as assistants to the regular hospital force.

Drills are held aboard ship simulating actual battle, and at these times the medical corps assumes positions at relief and dressing stations, established with a view to prompt and systematic aid to the wounded of the crew in mind, and makes all the preparations that would be necessitated by a real engagement.

It has taken some years and much labor in the face of difficulties to bring the medical department of a ship to its present state. Confronted by the problems of crowding a great deal into a small space, arranging rooms to conform best with the architecture of a ship, allowing for a wide range of climatic conditions, beset with questions of ventilation, unusual ship motion and distance from a supply source, the results have been very satisfactory and the benefit to the naval personnel marked.

AN IMPROVISED REFLECTOR FOR THROAT EXAMINATION.

J. HARVIE DEW, M.D.
NEW YORK CITY.

I have used the following simple device for a number of years in making examinations of the throat:

Showing method of using an improvised reflector in throat examinations.

It will prove of exceeding value whenever it is difficult to secure satisfactory light.

The idea was not original with myself nor was it obtained from any professional sources.

A candle, a tablespoon and another spoon suitable for depressing the tongue are the several essentials needed. The accompanying cut illustrates the procedure.

Hold the candle between the middle and ring fingers and the fore and little fingers of the left hand; then place the tablespoon between the thumb and fore and little fingers of the same hand. The concave bowl of the spoon should be placed immediately in rear of the flame. In this way a bright light can be thrown on any desired point.

Eczema of Extremities.—N. E. Aronstam, in *Medical Fortnightly*, says that, in acute erythematous eczema of the extremities, in which the subjective symptoms are very pronounced and annoying, the application of a dilute solution of the adrenal principle to the lesions will produce a rapid blanching of the parts and ameliorate the intolerable itching. After the acute features of this dermatosis have subsided an ordinary Lassar's paste with or without ichthyol wherein a small amount of adrenalin solution has been incorporated, will hasten the restitution of the involved areas to normal.