tain the opinion of the surgeon, so it would be advisable to consult the dermatologist before the *x*-ray is applied for an affection of the skin.

Regarding my own experience with it, I can report the following: Two cases of epithelioma originating from senile warts, both cured; five cases of ringworm of the scalp in which from six to twelve exposures produced total alopecia in the areas treated and in which the slight irritation soon subsided, and subsequent repeated attempts at cultures from the hair follicles proved negative; and one case of cutaneous tuberculosis which yielded splendidly and is healed over. Cases of acne rosacea and psoriasis showed improvement.

DR. PUSEY—I have not seen rheumatism that could be apparently attributed to the treatment with *x*-rays. Several gentlemen have spoken in this discussion about some poison being liberated in the system from neoplasms degenerating under *x*-ray treatment. I believe that there are no good grounds for such assumptions. It is theoretically possible that such a liberation of poisonous material might occur, but I have seen nothing to indicate that it does. I have seen large masses of carcinoma tissue and of gland tissue in Hodgkin's disappear, and I have never seen any symptoms to indicate that there was a liberation of any substance in the destruction of these tumors that in any way affected the patient's condition.

DR. C. E. SKINNER, New Haven, Conn.—In deeply-seated abdominal cases my experience so far has been that we do secure improvement usually and that we ought not to consider the rays inefficient in this situation as yet. It is only a question of how far the improvement can be carried. Up to the present time the improvement which I have noted has consisted of increase in weight, relief of pain, increase of appetite, and lessening of discharge where any was present. In one case of cancer of the uterus involving the bladder and rectum, improvement along these lines was marked and in addition the growth itself had diminished in size as a whole. All of a sudden, however, she developed uremic symptoms and died in a week. In view of the fact that she had been improving markedly up to this time, the sudden development of renal trouble might be construed as indicating that a toxin of some sort had been developed or liberated. I think we ought to be careful to recognize and weigh all the elements that appear in these cases.

DR. MONTGOMERY—Dr. Allen has very properly called attention to the danger of radiotherapeutics in the hands of the unskilled and inexperienced. We shall be fortunate, indeed, if, for a time at least, more harm than good is not done with this comparatively new therapeutic agent as the result of its indiscriminate use by those who lack the requisite skill for its successful and safe administration. Papers like the excellent one just read by Dr. Schmidt, giving full details of technic, will be of great value to those desiring to take up this work; but even with full instructions the beginner can not be too cautious in acquiring the experience necessary to a safe use of this somewhat uncertain therapeutic agent. There is a tendency to use the rays for all sorts of cutaneous and other disorders for which we have no other specific treatment, and there is no doubt that we are all using them now in a tentative and experimental way in cases for which the older and better understood methods will prove to be preferable. In other words, it is probable that as our experience enlarges we shall find that radiotherapy and phototherapy, while exceedingly valuable in certain disorders, will have a narrower field of usefulness than that in which we are now trying to apply to them. We have had good results in a few cases of acne and rosacea. Dr. Pusey refers to the use of the rays in a case of blastomycosis. In two of our recent cases, areas which persisted after a long course of potassium iodid disappeared quite promptly after a few exposures to the *x*-ray.

The matter of personal idiosyncrasy is, I believe, an exceedingly important one, and I am glad that the other men have spoken so strongly on this subject. Practically the only guide we now have in the treatment is the reaction in each individual case.

It has been suggested by two speakers that the Finsen light was for all practical purposes identical with the *x*-rays. The immediate results of the treatment are quite different. With

the Finsen light an exposure of ten or fifteen minutes will result in a dermatitis often vesicular or even bullous. Reaction occurs within twelve hours and disappears in a few days. No deep burn is ever produced. On normal skin pigmentation follows such a reaction and serves to protect the tissue from the light, so that a second treatment produces, as a rule, little or no reaction. In the diseased areas this protective pigmentation does not take place, and the reactions to successive exposures are like the first. With the *x*-rays, on the other hand, not only is the reaction slower in appearing, more persistent, and deeper seated, but reactions to successive treatments on both normal and diseased areas come, as a rule, more promptly and are more severe than those obtained from the first application. As to ultimate differences, we have found cases of lupus erythematosus and at least one case of lupus vulgaris in which the Finsen light has proven more effective than the *x*-rays. In at least one case of verrucous tuberculosis and in some other cutaneous affections the *x*-rays have given excellent results where none was obtained with the Finsen light. Our experience is as yet too limited to let us determine accurately the factors which cause one method to be more effective than the other in a given case.

DR. PFAHLER—I want to correct the statement of Dr. Pusey that I said no case of cancer had been cured. My paper states that cases have been cured and others are still improving under treatment. Another point that was brought out in the discussion was the protection of the hands of the *x*-ray worker. I use a mitten made of chamois skin covered on the back with lead foil and a piece of adhesive plaster over the foil. In regard to the point of heating the platinum anode, I always have my anode red hot, often white hot. I fully agree with Dr. Corlett and Dr. Allen that this is not only a very valuable therapeutic agent, but also a dangerous one.

DR. SCHMIDT—I think a large number take no precaution whatever to protect the hands of the operator. In fact, I think there is nothing to be applied or that can be applied to prevent the *x*-rays doing any damage, if an idiosyncrasy of the individual exists. The patients that come to us with *x*-ray dermatitis are those who do a great deal of photographic work. They handle chemicals and then possibly with the action of the *x*-rays a dermatitis is readily set up.

In regard to what Dr. Skinner said about heating the anode red hot or white hot, I am pretty positive that when we take a large number of workers in this field they will agree, if such a condition occurs, the tube is of a very low vacuum, and in addition to that there is an excessive amount of electricity passing through and some of the electric energy is converted into heat rays instead of *x*-rays. If any such condition exists for a long time, a puncture of the anode may readily occur.

TREATMENT OF FRACTURE OF THE NECK OF THE FEMUR.*

CHAS. E. THOMSON, M.D.
SCRANTON, PA.

For fifty years or more we find a sentiment pervading medical literature on the subject of fracture of the neck of the femur; that it is an irreparable injury; that a large percentage of these cases, being old, die of the trauma and shock following the accident, or the complications incident to it, viz.: Hypostatic congestion of the lungs from recumbent position or sepsis from bed sores; that the majority of those cases who live remain permanently disabled, getting about only on crutches or canes, on a badly deformed or functionally weak limb, the so-called "fibrous union." In support of this sentiment allow me to quote from the days of Sir Astley Cooper down to the latest literature obtainable, as taught in text-books on surgery, medical journal articles and medical schools.

* Read at the Fifty-third Annual Meeting of the American Medical Association, in the Section on Surgery and Anatomy, and approved for publication by the Executive Committee: Drs. H. O. Walker, A. J. Ochsner and DeForest Willard.

THE OLD TREATMENT.

COOPER.—In reference to the first-named author, that great surgeon admits that he had never seen bony union in a case of intercapsular fracture of the neck of the femur. His treatment consisted in placing the limb on a long pillow for ten days or two weeks, afterward allowing the patient to be up and about on crutches.

SAMUEL D. GROSS, in his admirable work, Vol. I, p. 1027, commenting on the treatment of Sir Astley Cooper twenty years ago, considered it too pessimistic and added that for many years he had been in the habit of treating fractures of the femoral neck (intercapsular) as any other fracture of the femur, viz., extension and counter extension, not with the hope of obtaining bony consolidation, but with the view of keeping the ends of the fragments in more accurate contact and thus affording them an opportunity of becoming united by fibro-ligamentous tissue.

AGNEW.—Writing one year later, Vol. I, p. 931, the great Agnew also criticised Cooper's treatment and advised treating fracture of the femoral neck on the same general principle as fractures elsewhere in the bone, viz., by extension on a straight or an inclined plane, adding that we have seen enough of fractures that were "believed" to be intercapsular recover with bony union to justify such procedure.

ECKHART MINES, MD., *North Carolina Medical Journal*, January, 1885, Vol. xv, p. 309. Youth of 18, crushed by fall of coal. Two inches shortening, inversion, no crepitus. Diagnosis: "The readiness with which the leg is brought down, absence of crepitus, coupled with the fact that after five weeks no attempt at union, nor discoverable callus thrown out, warrants me in believing that the fracture is within the capsule. Treatment: leg supported by pillows three weeks, Porter's modification of Smith's ant. splint, two weeks, further treatment useless. Allowed on crutches. Condition at writing: Shortening apparent enough, wears a high heel and walks with a peculiar rolling gait."

DR. SWING, Coatesville, Pa.—Lost 16 cases out of 18. All died in from six to nine weeks. Ages 70 to 80. Treatment: extension, counter extension and weights to limb and sand bags. Cases developed bed sores.

In subsequent case allowed patient to move about on mattress, got well with shortened everted limb, but could walk about her room last year of her life. Two years later second case could walk with crutches after two months, limb shortened and everted, lived one year. The doctor considered "the only rational" treatment to be to allow patient to move about on mattress of hair. He finds they will get well if spared the necessity of useless apparatus and will recover with as good a limb as they are likely to have if they could survive the torture of the apparatus. This treatment saves life and suffering. All treatment should have these two ends in view, rather than the glorification of the surgeon in displaying his skill in adjusting apparatus and setting limb. He feels sure that some of his cases would have gotten well if they had been "let alone" instead of receiving the orthodox treatment. He concludes by saying "the surgeon is not a mechanic to adjust appliances, but one whose judgment and experience should determine their necessity."

EVERHART, Scranton, Pa.—I am indebted to Dr. Everhart for report of the following case: Mrs. H., aged 71, in 1896 fell down stairs, was picked up and put to bed, about $\frac{3}{4}$ -inch shortening, suffered considerable pain, but did not have much deformity for four days, when she suddenly sneezed and at the same time felt something give way in the hip. Upon measuring the hip $1\frac{1}{2}$ inches shortening was found. The limb was markedly everted.

June 7, 1902: patient is still in bed, where she has remained for six years; 2 inches shortening, marked external rotation. Patient suffers great pain in hip, which is aggravated on slightest movement.

CORSER, Scranton, Pa.—I am indebted to Dr. John Corser for report of the following case: Mrs. C., aged 65, in 1887 fell, fracturing her hip; patient was put to bed, where she remained

for 15 years, when she died of exhaustion. During all these years was unable to walk or help herself in any way.

W. PUGIN, Thornton, April 2, 1894.—Patient, aged 87, was knocked down by horse, chloroformed, crepitus found, long splint, four nurses for 3 weeks. Water bed for one month. Got bed sore while on water bed. May 2 splint taken off and sand bags used. Four and a half months in bed, now goes about in bath chair; is able to walk a short distance with crutches. I do not expect he will improve much in this respect on account of his age.

OSCAR H. ALLIS.—*Medical Times*, August, 1883: "From first to last I make the patient my first care and regard the fracture as of secondary importance." He recommends getting patient up in rocking chair at end of first week, and from the first daily shiftings in bed. He thinks that getting patient up daily, carefully, does not produce separation of fragments. "In conclusion, I ask what has been, is and must ever be the outcome of all this? Will the surgeon risk his little all for the good of his patient or must he steer his course for that true but selfish standing, 'Self preservation is the first law of nature.'?"

HENRY LEHENAN.—Read March 19, 1894, before the Philadelphia County Medical Society: Mrs. —, aged 60, fell Jan. 23, 1881, while attending household duties; was not dizzy at the time. Saw her first Jan. 24, 1881. She was then lying on a couch and was unable to stand or move about. On examination, from the preternatural mobility, intracapsular fracture was diagnosed. My attendance was continued at intervals during the months of February, March, April and May. She was not able to help herself in any way. By help she could be gotten on a Charleston chair, and this was the only treatment attempted. The most of her pain was in the adductors; there was shortening and eversion of toe. She continued in the same room and on the same couch and chair until I was called to see her, Jan. 6, 1884, when I found her dying from exhaustion. She died January 8. Permission was granted to examine the hip. Specimen here presented was obtained. The acetabular cavity was normal in appearance. The head of femur lay in cavity free. The socket and head of bone were covered over with a membrane firm but somewhat incomplete. Upon this surface the upper extremity of the femur had formed an artificial joint, the neck having been absorbed. In the surrounding tissue on the inner side was a fragment of bone $1\frac{1}{2}$ inches long.

SCUDDER.—"In case of fracture of the neck of the thigh bone occurring in elderly individuals, treat the patient and let the fracture be of almost secondary importance." Again: "The impacted cases will unite; the unimpacted may unite." And again: "In only two cases out of sixteen could it be said that the leg was functionally useful."

PARK.—"Treatment must be so ordered as to save life even at the expense of an imperfect leg."

BERGER.—Gould's Year Book, 1897, p. 357: "In articular fractures those only in which operation is justifiable are about the elbow and knee, because an imperfect result here means a most grave functional loss."

AMERICAN TEXT-BOOK OF SURGERY.—Page 310, 1892, recommends extension and fixation with sand bags and plaster of paris, and in ununited fracture reports that several cases have been cut down on but fail to give results of such operations.

CURTIS.—"In certain cases, of over 60 years of age, give up the idea of bony union."—From lectures on surgery, University of New York.

From the above it is apparent that at least a goodly number of our authors have been rather skeptical about obtaining bony union in those cases, and in discussing the subject, both in medical societies and in private conversations, I find the average practitioner—under whose care those cases are usually treated—is satisfied if his patients are able to hobble about on crutches for the remainder of their lives, nor has this pessimistic standing been confined to cases treated by ordinary methods and general practitioners.

THE OBTAINING OF BONY UNION.

BRYANT.—Having reported 6 cases from König, Cheyne, Myre, Gillette and Curtis, all of whom obtained firm bony union and strong useful limbs, Dr. J. D. Bryant, in his valuable work on operative surgery, vol. ii, p. 1279, 1901, concludes as follows:

"There is yet a reasonable doubt regarding the advisability of this operation except in special cases. The difficulty of securing and maintaining proper adjustment of the fragments because of the lack of command of the inner portion, its porous character and low vitality, present obstacles to success that can not be gainsaid. The degree of shortening that follows the successful results suggests an initial failure of reduction of the deformity or the maintenance in proper place, and bespeaks in any event a considerable amount of absorption at the seat of the fracture. A more extended experience is needed and a careful comparison of the favorable results by different methods of treatment is required before a final judgment can be recorded."

BERGER.—Gould's Year Book on Surgery, 1897, p. 357: "In articular fractures those only in which operation is justifiable are about the elbow and knee, because an imperfect result here means a most grave functional loss."

Is not the hip joint an important one and quite as worthy as the elbow and knee of our special consideration? I am pleased to add Cases 5 and 6 to the successful operative list, and hope that others may be added in sufficient numbers to put the operation out of the experimental class beyond all doubt, even in the minds of the most skeptical.

Among the authors that have written more hopefully on this subject are Hamilton, Wyeth, Dennis and Ridlon.

RIDLON.—The last-named surgeon, in his excellent article on treatment of fracture of the femoral neck, *Annals of Surgery*, July, 1901, has so fully expressed my views that I take the liberty of quoting from it as follows:

"Let me assure you, gentlemen, that in all these cases a strong and useful limb, in good position, may be as confidently anticipated as in any other fracture of this bone or any other bone. In no other fracture is the surgeon's task easier; in none is a good result more certain." Again: "I believe we can only treat the patient properly, by properly treating the fracture from which she is suffering and likely to die." And again: "The average surgeon treats fracture of the neck of the femur precisely as he treats a solution of bony continuity where he aims to get a false joint; and, verily, he usually has his reward. My plea is simply this: Show the fracture at the neck of the femur the same consideration that you show all other fractures. It is useless to quibble about fractures within the capsule and fractures outside the capsule; fractures impacted and fractures unimpacted; about the thin, weak bone of fat old women; these cases invariably result in strong and useful union if they are given a fair chance."

I cannot possibly improve on the above quotations, nor write anything that more aptly expresses my views. I could never understand how any thinking man can pretend to benefit his patient in any way by leaving a fracture unreduced and unimmobilized. I suppose men who have never reduced nor immobilized a fractured femoral neck cannot be expected to have noticed the great relief given to a patient by such a procedure.

METHOD OF TREATMENT.

I regret that I have to differ from my friend, Dr. Ridlon, in his method of treatment, yet I believe he has the most complete and efficient apparatus ever devised for the treatment of fracture of the femoral neck, but such apparatus is made well only by skilled mechanics, whose work is expensive and hard to obtain when most needed. Then, again, each case requiring a specially made apparatus, a number of circumstances may arise causing long delay and loss of valuable time,

while the patient is suffering from unreduced fracture and the friends are impatient because the limb has not been "set." All this care may be obviated by the surgeon who carries in his satchel sufficient material, and in his brain sufficient intelligence, tact and skill to make a plaster-of-paris spica. That a surgeon should stop before he really commences to treat a difficult case and wait for the whims and efforts of a mechanic is surely a reflection and a humiliation on modern surgery and should not prevail. Then, again, I think my friend has rather overestimated the task of putting up a fractured femoral neck in plaster-of-paris spica. Beside the man who is giving the anesthetic, all that is required is a McBurney hip rest and one assistant with sense enough to do what he is told, and strength enough to make the necessary extension. Then, again, they speak of disturbing the fracture by changing the plaster. I have never found it necessary to change a plaster cast on one of these cases, and have frequently left them on three months, or long after the patient was up and going about. It is not necessary to get these casts soiled, and as to excoriations and bed sores, they simply never occur, if the plaster is properly applied. On the contrary, I frequently put on a plaster-of-paris spica to cure a bed sore; by so doing you can distribute the pressure and support the circulation, two very important factors in the treatment of bed sores.

What cases should be operated on? No recent case. Why operate when we get such excellent results without operation? The two cases I have operated upon, of three and eighteen months' standing, respectively, were not even diagnosed before I saw them. I think a fair rule would be that any case that remained ununited for three months and had received judicious care, would be a case for operation.

Dr. Bryant has said, *Operative Surgery*, vol. ii, page 1287: "In operating on cases of ununited fracture the refreshing of the ends of the fragments is a necessary step in the technic." I made no attempt to freshen the ends of bones in my cases and yet I got prompt and firm union.

The method of treatment I have used and intend to recommend, simply carries out the principle of treatment of fracture in any other bone in the body, viz.: Reduction of fracture and retention of fragments in corrected position, for sufficient length of time to allow Nature to make the necessary bony repair. This I believe to be best obtained by anesthetizing the patient, so as to relax the muscles and enable the surgeon to ascertain the exact nature of the injury and more completely replace the parts. Impacted fractures should be very carefully handled. In fracture of the femoral neck this is best accomplished by having an assistant make forcible extension on the limb (while a second assistant, the anesthetizer, makes counter extension from shoulders) until it is drawn down to equal length with its fellow, while the operator manipulates the upper end of the bone, reducing the fracture as well as possible; and holds it while the assistant carries the limb into abduction of about 10 degrees. This position of abduction has a tendency to hold the fractured ends together, and facilitates in keeping the patient clean afterwards. The operator now applies shaker flannel bandages to the entire limb and body, as high as the arm pits, amounting to about four layers. Over this is applied plaster-of-paris dressing (reinforced with soft steel plates at hip and knee) sufficiently strong to maintain the form. During the application of the spica the limb is constantly held by the assistant in extension and ab-

duction as noted above. Care must be taken for the first twenty-four hours that undue pressure or strain is not brought to bear on the plaster before it is sufficiently dry to resist. If at any time the patient complains of pain over prominent parts, as the knee or malleoli, a portion of the plaster may be removed to relieve the pressure. After the plaster has become sufficiently dry the patient may be turned from side to side to relieve the hypostatic congestion of lungs, or for the purpose of bathing. If the spica is continued sufficiently high on the thorax it will not materially interfere with respiration. It is only when it ends at the free border of the ribs that patient complains in this respect. By this method of treatment the patient is enabled to materially assist in the use of the bed pan without the danger of disturbing the fragments.

SUCCESSFUL CASES.

The following is a list of my own cases, together with a number of successful cases from other authors, that I take the liberty to present.

SHEEHAN.—*British Medical Journal*, 1895: January, 1893, J. H., aged 91, knocked down by cab; two inches shortening, no crepitus. Treatment: Extension, weight and pulley, sand bags. Allowed up in a few days, on crutches, got about until his death from nephritis, April, 1894. Specimen showed bony repair in good position. Amount of shortening not given.

JOYCE.—Page 907: Dec. 28, 1895, patient fell while intoxicated, three-quarter inch shortening. Treatment: weight and pulley. Jan. 7, 1896, showed signs of pneumonia, propped up in bed, turned over; improvement followed. January 20, extension had to be removed as the skin about ankle and foot gave way. Next five weeks no treatment, but kept in bed. In three weeks walked with crutches. May 12, walked with two sticks, one inch shortening, complete bony union. Was since able to follow his trade of peddling.

ALEXANDER HATTEN, Portsmouth, Pa.—*New York Medical Journal*, Nov. 13, 1897: On Sunday, March 8, 1896, was called to see Mrs. C., aged 73; much emaciated, being about 5 feet 3 inches in height and only weighing 95 pounds. She had fallen while entering a church door and complained of severe pain at the left hip joint and lost motion in the left thigh. She was placed on a stretcher and carried to her residence, where on examination I found an intracapsular fracture of the neck and left femur with no impaction. She was placed on her back in a fracture bed and her leg put in good position. Buck's extension was applied and two sand bags were used, one on inner and one on outer side of leg, the latter extending up to waist. She remained in this position for six weeks, at the end of which time Buck's extension was removed, as it was found the fracture had entirely united. In three weeks more she was able to walk with the assistance of crutches. She now walks well without assistance and only has about one-half inch shortening.

AUTHOR'S CASE 1.—Mrs. C., aged 62, on August 19, 1901, was thrown from her carriage, striking violently on her hip. Seen by the writer three days later, limb everted, 2½ inches shortening, very painful. Patient chloroformed, fracture reduced, plaster-of-paris cast applied. Two weeks later was transported six miles over country road and 125 miles on railroad train to her home. Plaster left on eight weeks, when she was allowed on crutches.

Present Condition.—June, 1902, nine months after accident, limb in good position. Can walk without crutches or cane but uses cane on street. At present time is visiting friends and enjoying herself.

AUTHOR'S CASE 2.—Mrs. S., aged 66, 1897, sustained fracture of hip, 2 inches shortening, chloroform administered, fracture reduced and limb encased in plaster-of-paris spica, in extension and abduction. Cast left on for 8 weeks, when she was allowed on crutches.

Present Condition.—Has done all her house work for three years; walks with slight limp. Patient absolutely refuses to have limb measured.

AUTHOR'S CASE 3.—Mrs. F., aged 76, on Oct. 18, 1900, fell off porch 2 feet, striking on hip; 2 inches shortening, limb everted, great pain. Under complete chloroform anesthesia, limb placed in best possible position, plaster-of-paris cast applied and left on for nine weeks, during which time patient was kept in bed, and then allowed up on crutches.

Present Condition.—Shortening 1¼ inches, limb straight, 45 degrees free and painless motion. Walks without cane or crutch, no pain in hip, goes up and down stairs unaided although 78 years old.

AUTHOR'S CASE 4.—Mrs. P., aged 59, in April, 1901, fell down steps 4 or 5 feet, striking on hip; seen by writer one week later: patient in bed suffering a great deal of pain which was intensified on slightest attempt at motion; 1 inch shortening. Temporary plaster-of-paris spica applied. Patient transferred 4 miles over country road, and 20 miles on railroad train to hospital. Next day, under complete anesthesia, plaster-of-paris spica was removed, limb carefully examined and previous diagnosis of impacted fracture confirmed. Permanent plaster-of-paris carefully applied, patient put to bed for nine weeks, spica removed, allowed to walk on crutches.

Present Condition.—June 3, 1902, union firm, position and motion of limb normal, 1 inch shortening; can bear entire weight of body on limb without pain, and walks well without cane or crutch. Has done all her housework for 3 months.

SUMMARY OF AUTHORITIES ON OPERATIVE TREATMENT.

KOENIG.—The operative treatment in these cases was first suggested by Langenback, and successfully carried out by Koenig. The latter operated in a case of recent fracture, making a small incision over the trochanter major, drilled a hole through it with a metal drill in the direction of the head of the bone, applied extension to the limb to the extent necessary to overcome the deformity, and then drove a long steel nail through the canal in the trochanter into the head of the bone and left it there. The limb was then immobilized and extended for six weeks. There is no record of the ultimate shortening, but good union and free motion of the joint was obtained.

CHEYNE.—In a case of recent fracture, Cheyne exposed the fragments through a longitudinal incision made over the anterior aspect of the joint, exposed the fracture, made extension and internal rotation of the limb, and with the fingers in the wound manipulated the fragments into place; then a small longitudinal incision was made over the outer side of the trochanter major, and two canals drilled through the fragments at a distance of half an inch apart. Ivory pegs were then driven through the holes made by the drill, and the limb immobilized. Good union and motion were obtained, but there is no record of measurements of the limb.

MEYER.—In a case of ununited fracture with three inches shortening, Meyer made the Langenback incision for excision of the hip joint (Bryant's Operative Surgery, Fig. 393), exposed the seat of fracture, scraped the ends of the fragments until they bled, reduced the deformity by extension of the limb and fastened them together by driving two nails through the trochanter major. A useful joint was obtained, with an inch and a half shortening of the limb.

PARKHILL.—This method is commended for fracture of the neck of the femur (Fig. 1550 in Bryant's Operative Surgery).

GILLETTE.—Three cases of ununited fracture of the neck of the femur are reported operated on in the following manner: A horseshoe incision with its convexity downward was made, beginning an inch below and an inch posterior to the anterior superior spine of the ilium, carrying it down two inches below the trochanter major and bringing it up the buttocks to about the center of the gluteus maximus muscle. The skin and the two layers of fascia were dissected up en masse. A chain saw was then passed between the posterior border of the tensor vaginae femoris and the gluteus medius, hugging the neck of the femur and the base of the trochanter major; it was brought out through the posterior surface of the gluteus medius and the anterior surface of the gluteus maximus; the trochanter major and its muscular attachments were sawed off, turned back, thus exposing the capsule of the joint. Then, by making a longitudinal incision into the capsule the fracture

could be easily seen. The surfaces of the fractured ends were denuded and a bone peg driven through the neck of the femur, thus holding the fractured ends together. The capsule was

Fig. 1.—Showing fracture before operation and atheromatous femoral vessels.

stitched with catgut, the trochanter major restored and nailed in place with a small bone peg, the skin closed and the limb immobilized. There was union and good motion obtained in

Fig. 2.—Showing hip after operation.

all cases, with shortening of from an inch to an inch and a half.

CURTIS.—In a case of ununited fracture of neck of the femur of three months' standing, Curtis exposed the fracture through an anterior incision, passed a drill into the callus and be-

tween the fragments to cause irritation, applied extension to the limb and reduced the deformity, after which a drill was passed through the trochanter major from the outer side, transfixing the fragments; the handle of the drill was then removed and the drill itself left in situ. The anterior wound was closed and the limb immobilized. The extension was maintained for six weeks, at the end of which time the drill could be easily removed. Good union and a useful limb were obtained with three-quarters of an inch shortening.

LISTER.—*British Medical Journal*, Aug. 26, 1871: Sir J. Lister has recorded the case of an ununited extracapsular fracture of the femur in a man aged 45, where, eighteen months after the injury, he cut down on the fragments with antiseptic precautions and gouged them, the fracture being then finally put up. Recovery was complete, the man walking well.

AUTHOR'S CASE 5.—M. M., aged 60, miner by occupation, on Feb. 7, 1900, fell on ice, striking on hip, was carried home and put to bed for three days, afterwards allowed to sit in chair, where he spent 3 months; could not put weight on limb for 18 months, had constant pain in hip, walked with both hands on cane (bent almost double) on which he carried the greater part of his weight, dragging his right limb; one and one-half

Figure 3.

inches shortening, marked eversion of limb, on manipulation distinct movement could be made out between head and trochanter (Fig. 1). On examination patient appeared much beyond 60 years, and showed signs of senile degeneration, viz., general muscular atrophy, arcus senilis, and atheromatous arteries (Fig. 2).

On Aug. 28, 1901, under complete chloroform anesthesia, a horseshoe-shaped incision was made around the great trochanter down through the muscles to the periosteum. In this case no bone flap was made, the capsular ligament was opened, and the free motility of the head ascertained. We deemed this step necessary because the skiagraph showed what we feared was ankylosis of head in the acetabulum. I then drove two solid silver nails, 3/16 inch in diameter, and 2 and 3 inches long, respectively, through the great trochanter, neck and into head of bone. Only the long nail reached the head, as was afterward shown by x-ray (Fig. 2). It was not necessary to use drill before driving nails, capsular ligament was closed with catgut sutures, external wounds with silkworm gut without drainage, aseptic dressing applied and entire limb and body encased in plaster-of-paris spica. Patient did not suffer from operation, was allowed to turn over in bed, no elevation

of temperature. Wound was dressed through a fenestra in spica at end of third week, when sutches were removed, primary union complete. Was removed from hospital to home at end of six weeks, and spica taken off at end of nine weeks, when union appeared firm, 45 degrees of free motion in the hip. Three months after the operation he could bear over half the weight of body on the limb, could walk upright; 1 inch shortening.

Present Condition.—Nine months after operation the limb is straight, motion normal, shortening 1 inch, can bear entire weight of body on limb without pain, walks with cane but can walk very well without it. No pain except on direct pressure over seat of operation.

AUTHOR'S CASE 6.—This case was reported at the St. Paul meeting and published in *THE JOURNAL* of Dec. 21, 1901. Sustained fracture of hip by a fall of rock while working in mines. Was seen by me and operated on 3 months later; practically the same operation as in Case 5, except in this case a bone flap was made before driving nail and but one nail was used besides the small nail for bone flap. Plaster-of-paris dressing the same; wound healed by first intention. Patient walked in three months with one inch shortening. Improvement continued until at the time of his death he could walk without cane or crutch; had 90 degrees of free and painless motion. Death 18 months after operation from intercurrent disease. Specimen obtained. Bony union firm in good position (Fig. 3).

SUMMARY.

1. I believe that the teachings and writings of surgical authors have been rather discouraging to the general practitioner on this subject.

2. That in all cases of fractured femoral neck, firm bony union and useful limbs may be anticipated.

3. That age is not a counter indication to treatment nor to obtaining bony union.

4. That the patient is best treated by reducing the fracture and immobilizing it.

5. That this is best accomplished under anesthesia and by the use of the plaster-of-paris spica.

6. That the immobilization should be continued for a long time, and three months should elapse before allowing weight on the limb.

7. That Buck's extension with weight and pulley is not sufficient immobilization to obtain bony union.

8. That the use of apparatus in these cases is expensive, hard to obtain when needed, not so efficient in immobilization, and less convenient for the patient and nurse than plaster-of-paris spica.

9. That the operative treatment in old and neglected cases has succeeded beyond all expectations and deserves a place in surgery among the radical cures for troublesome conditions.

10. That the patients who usually suffer from the accident being old and enfeebled is a good reason why the physician should make their declining years as peaceful and pleasant as possible.

Intravenous Injection of Chloral in Treatment of Varicose Ulcers.—An article in the *Gazzetta Med. di Roma* for June states that Scalzi has treated 200 patients with varicose veins or ulcers by the intravenous injection of a Pravaz syringe of equal parts chloral hydrate and distilled water. No inconveniences were noticed in any instance. He applies a constricting bandage above. The coagulum forms so rapidly that the constriction can be removed ten minutes after the injections. If the lesion is merely a varicose vein, the chloral mixture is injected at different points in the vein; if an ulcer, the injections are made in the most swollen portions of the vein, close to the edge of the ulcer. The beneficial effects are evident in a few days and the ulcer gradually heals. Enrico, the author of the article, lauds this as the simplest and most effective of all methods of treating varicose veins and ulcers.

THE TREATMENT OF TRANSVERSE FRACTURE OF THE PATELLA BY SUBCUTANEOUS PURSE-STRING SUTURES.*

JOHN B. ROBERTS, M.D.

PHILADELPHIA.

In April, 1896, I reported¹ a case of patellar fracture satisfactorily treated with a catgut ligature carried around the fragments, in the coronal plane, making a sort of purse-string suture. This method seemed to me preferable to Barker's method, in which the ligature is introduced in the sagittal plane and traverses the knee joint and prepatellar bursa. I do not know that the method was new, but it is certainly deserving of extended use. Further experience has increased my confidence in it and leads me to advocate its adoption. It requires no special apparatus, such as Malgaigne's hooks, does not invade the knee joint or the synovial sac in front of the patella, and demands no general anesthetic. It is practically free from septic risks, allows the patient to get out of bed soon, and results in such a close union of the fragments that, even if true bony union is not obtained, the function of the limb is perfect.

Clinical experience and laboratory investigation seem to justify me in suggesting some modifications of the original procedure, which will, I believe, add to its surgical value.

The only instrument used is a stout needle five or six inches long and a strong ligature of silk, soft wire, kangaroo tendon or chromicized catgut. About a week is permitted to elapse after the receipt of the injury, in order to give time for absorption of blood and synovial fluid. During this period I keep the patient in bed on his back, with the knee extended and the limb elevated in order to relax the rectus muscle, which, by reason of its origin above the hip joint, may tend to draw the upper fragment upwards. An ice bag is applied to the knee to hasten the reduction of inflammatory swelling. If the intra-articular effusion were very great, I would not hesitate to tap the joint with a small aseptic trocar. This is probably seldom necessary.

When the purse-string suture is to be applied, the skin is sterilized and the aseptic needle thrust transversely through skin, muscle and fascia, just below the lower fragment of the patella. It is carried deeply enough to insure pressure against the bone when the purse-string is drawn tight and tied; but it is not expected to penetrate the joint cavity. The aseptic suture is drawn through the tissues by means of the needle. The needle is then inserted at the puncture of exit and thrust upwards, through the tissues, alongside of the broken patella, until its point emerges at a point a very little higher up the thigh than the upper edge of the upper fragment. The suture is drawn through this needle track. During these manipulations the two fragments are held together by the surgeon or an assistant. The needle is then introduced at the second puncture of exit and carried through the muscles just above the upper margin of the upper fragment. Its transit carries the ligature across the thigh superficially to the synovial sac of the joint. The needle is then inserted at the fourth puncture and is thrust downwards alongside of the two fragments until it emerges at the first puncture. It is pulled through this open-

* Read at the Fifty-third Annual Meeting of the American Medical Association, in the Section on Surgery and Anatomy, and approved for publication by the Executive Committee: Drs. H. O. Walker, A. J. Ochsner and DeForest Willard.

1. *Annals of Surgery*, vol. xxiv (1896), p. 229.