NOTES SUPPLEMENTARY TO TABLE.

Note 1.—Case 16. In this case pia is also thickened, but over different regions and asymmetrically.

Note 2.—Case 5. Exaggeration of deep reflexes on left side.

Note 3.—Case 9. Infantile cerebral palsy at six months, i. e., right hemiplegia. Examination showed athetoid movements of left (or right?) side, athetoid movements of right hand.

Note 4.—Case 10. Spastic ataxic gait. "Exaggerated reflexes" on both sides. Temporary paraplegia at age of 21. Syphilitic infection at age of 25 (or earlier?).

Note 5.—Case 14. Spastic gait. No history notes accounting for it.

Note 6.—Case 16. Slight right hemiplegia; defective, uncertain, slow speech. "Exaggerated reflexes." Paralysis came on at age of 3 or 3½ years, during an illness accompanied by high fever and convulsions. Attributed to overdose of opium (?).

Note 7.—Case 17. In this case of hydrocephalus there was divergent strabismus, present since admission and for some hours before death a tonic spasm of right arm and right side of face. Otherwise no reliable date pointing to organic brain lesion.

Note 8.—Case 18. At age of 2½ stroke of right hemiplegia; at 6½ years stroke of left hemiplegia. Both recovered (?).

Epilepsy set in after the second stroke. Clinical notes say only: "Exaggerated reflexes."

Note 9.—Case 19. "Left knee jerk exaggerated."

Note 10.—Case 22. Weakness of left arm, chiefly after attacks. Epileptic seizures of Jacksonian type, with preservation of consciousness and involvement almost exclusively of left arm, left side of face and left side of neck. Chancre two years before death. A few days before death he developed thick speech, partial aphasia and paraphasia, disturbance of deglutition and apparent involvement of vocal cords.

Note 11.—Also Case 22. The valvular lesion was here evidently an acute one, possibly a syphilitic one.

discussion. In view of their relative rarity as compared with the thalamic changes, they can not be given as much attention as the latter, but yet should not be allowed to pass without mention.

Claus and Van der Stricht,⁹ who discuss the pathology of epilepsy quite extensively, make the following statements on cerebellar changes in epilepsy.

The sclerotic indurations of the cerebellum are very frequent. They are often found at the autopsies of the epileptics. They have no special significance, they indicate likewise (*egalement*) a more generalized state of sclerosis. If localized exclusively in the cerebellum they can, just like the sclerosis of the cornea ammonis, of the olivary bodies, or of other parts situated outside of the motor centers, have an etiologic significance in the production of epilepsy.

The authors then go on to quote Duquet as having reported three cases of sclerosis of the cerebellum. They further mention Claus as giving the history of a female epileptic whose gait at the onset simply unsteady (*chancelante*) became more disturbed in the further course of the disease. The autopsy revealed a sclerous atrophy of the cerebellum. The cells of Purkinje and the nerve cells of the external layer were also diminished and atrophied.

Finally, Claus and Van der Stricht mention Pierret as quoting without comments a case of peripheral atrophy of the cerebellum with accompanying atrophy of the two olivary bodies and integrity of the rhomboid bodies. They conclude by saying: "We find it useless to multiply these instances."

If I have dwelt at some length on the changes of the cerebellum in epileptics it is because some authors have attributed to the cerebellum a relation to emotions. Close contact with epileptics shows, on the other hand, how frequently in these patients the emotions are at fault. (See table, emotional state.) However, I am fully aware that the small material at hand and the speculative character of the emotion theory of the cerebellum does not permit of any definite conclusions.

DISCUSSION.

DR. DAVID I. WOLFSTEIN, Cincinnati, declared that this paper is valuable, as it certainly supports in a startling way

9. Claus and Van der Stricht: *Essai sur la pathogénie et le traitement de l'épilepsie*, Bruxelles, 1895, p. 165.

the recent contribution to the study of these cases made by Dr. Starr, in which he made an effort to pick out a class and place them under the head of organic. It seems that in this group there is confirmatory evidence that organic lesions would be found.

DR. WILLIAM G. SPILLER, Philadelphia, stated that he has many brains of epileptics from the Pennsylvania Training School for Feeble-Minded Children, and has observed changes similar to those which have been noted by Dr. Onuf. He does not think it wise to remove the pia, though by so doing better photographs can be made, but one can not then study the condition of the pia. Dr. Onuf's method of hardening the brain has been employed in a somewhat modified form by others. A physician, for the purpose of studying the retina, injected formalin into the eyeball immediately after the death of the patient, and succeeded very well in preserving the nerve cells. Dr. Spiller agreed with Dr. Onuf that it would be well to devote more time to the clinical study of the ocular and aural symptoms.

DR. N. B. ONUF said that in principle Dr. Spiller's objection to removal of the pia is entirely justified. He explained, however, that the removal is not so objectionable as it seems on the surface, since the removed pieces were marked in the relation to the region from which they were taken. The pia was, for instance, removed in a strip over the second frontal convolution, and a note was made which end was the anterior one and which end adjoined the anterior central convolution. In such a manner the pia strips were all marked and removed for future study. Dr. Onuf declared that it is but fair to state that most of these cases were cases of advanced epilepsy with marked mental changes, usually marked dementia, but this does not exclude the possibility of finding such changes also in earlier stages of the disease.

CEREBROSPINAL FEVER.

EPIDEMIC CEREBROSPINAL MENINGITIS.

J. C. WILSON, M.D.

PHILADELPHIA.

HISTORICAL.

Two diseases have especially excited the interest and baffled the researches of students of epidemiology and clinical medicine. One is influenza, the other epidemic cerebrospinal meningitis. Influenza has occurred in world pandemics which have been succeeded for varying periods extending over years by recurrent epidemics and sporadic cases and finally by periods of apparent absence of the disease until the sudden recurrence of another great pandemic. The opportunities afforded by the outbreak which began in 1889 have resulted in the discovery of the pathogenic agent of influenza—the bacillus of Pfeiffer—in the recognition of the transmissible nature of the infection and of the existence of sporadic cases by which the disease has been maintained for an indefinite time. But the regions in which the organism has continued to exist and the efficient mechanism by which, after long periods of latency, it suddenly gives rise to its pandemic effects, have thus far eluded research.

Epidemic meningitis—cerebrospinal fever—for about a century has been recognized as a substantive disease. Its occurrence in epidemics more or less extensive and prevailing under conditions wholly at variance with those which obtain with other transmissible or contagious diseases, its terrible symptom-complex, high mortality and the desperate state of many of those who survive the attack, together with the fact that the disease remains in sporadic form as a continuing menace to the health of communities in which it has once oc-

curred in the epidemic form, have given it a singular and gloomy position among the pests of mankind.

The preliminary work of Weichselbaum (1887) and Jäger (1895) and the definite report of Councilman, Mallory and Wright (1897) have demonstrated the exciting cause of this specific form of cerebrospinal meningitis to the *Diplococcus intracellularis meningitidis*. A few days ago Councilman¹ published a most important contribution to the bacteriology and pathology of the subject, in which a brilliant side-light is thrown on the obscurities of the disease in certain of its epidemiologic and clinical aspects.

Quincke's lumbar puncture as a familiar procedure in clinical work has proved of inestimable advantage in the differential diagnosis between cases of pneumococcus and streptococcus meningitis and those due to the *Diplococcus intracellularis meningitidis*. Much has thus been done to clear up the obscurities regarding the origin and nature of cerebrospinal fever, but much remains to be done, especially with reference to the cases that are not well characterized and in the direction of treatment. The subject acquires a peculiar interest at this time on account of the epidemic which has prevailed in New York City since February, 1904, resulting in more than 1,000 deaths, the number recently reaching 130 deaths per week, and the evidences of its tendency to spread in Philadelphia since the localized outbreak of 23 cases which occurred at the U. S. Navy Yard at League Island in the spring of 1904. At the present time there are 3 cases in the medical service of the Pennsylvania Hospital. It is also reported that the disease has assumed an epidemic form in some parts of Europe, especially in Silesia.

Cerebrospinal fever may be defined to be an acute, infectious, epidemic disease caused by the *Diplococcus intracellularis meningitidis*, characterized clinically by sudden onset, with headache, vomiting and painful contraction of the muscles of the back of the neck, irregular fever, profound nervous symptoms, rapid course and high death rate; anatomically by inflammation of the meninges of the brain and cord.

This disease was first recognized as a distinct affection about the beginning of the nineteenth century. It is probable, however, that it has existed from a remote period. Vieusseux described an outbreak which appeared in Geneva in the winter and spring of 1805. The following year an outbreak in Medford, Mass., was described by Danielson and Mann. From that time the disease was extensive in the New England states, prevailing in circumscribed epidemics until 1816. At the same time and subsequently, as shown by Hirsch, the disease prevailed in various parts of Europe and in the United States, chiefly in isolated epidemics. Stillé fully described the disease as it prevailed in Philadelphia and in other parts of Pennsylvania in 1867. In the last three decades scattered cases occurring singly or in small groups have led to the belief that the disease has become naturalized in regions in which it previously prevailed in epidemic form.

Climate appears to have little influence as a predisposing factor. Epidemics have occurred in Sweden and on the Mediterranean coast, and in this country from the Gulf coast to Canada. Outbreaks are more common and extensive in the winter and spring than in the warm seasons of the year. Densely populated cities and sparsely settled agricultural regions are alike subject to its prevalence. Observers concur in the statement that

damp, overcrowded and unclean habitations favor its spread and that persons living on the ground floor are especially apt to suffer. In adult life the proportion of males attacked is greater than that of females. Among children the number of males and females is about equal. Cerebrospinal fever is especially a disease of children and young adults. After 40 it is uncommon, though the diagnosis has been verified postmortem in individuals over 70. Individuals of all occupations and professions are liable to this disease. Military life involves a special liability.

COMMUNICABILITY.

Cerebrospinal fever does not appear to be contagious in the sense in which we are in the habit of using the term in speaking of smallpox, scarlet fever and typhus. Councilman, in his first report, states that the evidence, on the whole, is not conclusive that the disease is incapable of being transmitted from one individual to another. The definite micro-organism which causes it is, in the majority of instances, confined to the meninges of the brain and cord, with little or no opportunity of transmission to other individuals. In cases in which there are lesions in the lungs, ears and nose, however, infection of neighboring objects or persons may readily take place. It is important at this point to note the following statement of Councilman in the recent contribution to the subject above referred to:

The presence of (these) sporadic cases is of importance in the occurrence of epidemics. The *diplococcus intracellularis* is an organism of feeble vitality; it dies out easily on exposure to drying and light and is incapable of a saprophytic existence. In the absence of intervening infections, it would be impossible for the period of epidemics to be bridged over. Not only this, but there is evidence that this organism can produce other infections and may even live as an inhabitant on the normal mucous membrane.

The older literature contains numerous accounts which suggest the possibility of the specific exciting cause of this disease being communicable from one person to another, or at least portable on the person or among the effects of those who have been exposed to it. Many of these relate to the outbreaks among French troops and are highly suggestive. The two following cases are cited by Hirsch as occurring in the epidemic of West Prussia in 1865:

On February 8, K., aged 20, fell ill in the township of Sezakau. He was nursed by the girl W., who had hastened to him out of the village of Sullenczyn. After the death of K. his nurse returned home and there died, on February 26, of epidemic meningitis. This was, with the exception of one on January 15, the first fatal case of meningitis in Sullenczyn. To the burial of this maid came the family of the farm-steward K., to Sullenczyn from the township of Podgass, accompanied by the servant D. and the 4-year-old daughter O. of the teacher R. in Podgass. After their return from the funeral a little child of K.'s fell ill and died in twenty-four hours, then the servant D., who died on March 4, and finally the girl R. on March 7.

At another village, two children of one family, aged, respectively, 1½ and 3½ years, died of the epidemic, one on January 27 and the other on February 7. The clothes of the deceased were taken to a neighboring village, and came into the possession of a girl, aged 5 years. She soon sickened of the epidemic and died on February 14.

Stokes reports the following case:

A child was seized with epidemic cerebrospinal meningitis and died. A second child of the same family was attacked with the malady a few days later. The day following the

1. The Journal A. M. A., April 1, 1905.

attack of this child the mother, who slept in the same bed with it, sickened of the disease.

The next case was observed by J. Lewis Smith:

A boy, 12 years of age, died of cerebrospinal fever and was buried on Saturday or Sunday. On the following Monday the mother washed the linen of the boy, which had accumulated, and within two days was herself affected with the disease. She and her infant, who was also seized with it, died.

The following personal observation is not without interest:

I saw at 8 o'clock in the evening of March 13, 1905, in consultation with a medical colleague, a physician, 33 years of age, suffering from fulminant cerebrospinal fever, the first symptoms of which had appeared on the previous evening and which terminated fatally during the course of the night in a little more than twenty-four hours from the onset. This man had been in constant attendance on a patient who had died of malignant cerebrospinal fever of two days' duration about twelve hours before his own sickness began.

In view of the foregoing facts it can not be affirmed that cerebrospinal fever is not under certain circumstances a transmissible disease.

THE SPECIFIC ORGANISM.

Second attacks are exceedingly rare. The Massachusetts report refers to five instances collected from the literature. It is probable that in most cases persistent immunity follows the attack. The pathogenic organism is found in the fluid obtained by lumbar puncture and in the meningeal exudate. The cerebrospinal fluid is usually more or less turbid, sometimes very turbid, especially early in the course of the attack. In a chronic case under the care of my colleague, Dr. Lewis, in the Pennsylvania Hospital, the fluid was turbid when first withdrawn, but perfectly clear on a second puncture made 43 days later, although the symptoms persisted. In one of my own cases in which the duration of the illness could not be determined and the patient prior to his admission to the hospital was supposed to be suffering from uremia, the fluid was perfectly limpid, but showed pure culture of the *Diplococcus intracellularis meningitidis*.

While turbidity of the spinal fluid is of diagnostic importance, its perfect limpidity does not constitute a negative sign, and in either case cultures are necessary to the diagnosis.

The organism grown in pure culture appears in diplococcus form with an unstained interval. It reacts to ordinary stains and is decolorized by the Gram method. It grows best on Loeffler's blood serum mixture. In the tissues the diplococcus is almost constantly confined to the interior of the polynuclear leucocytes. The diplococci are found only in connection with the lesions of the disease. Mixed infections are not uncommon. In the Boston cases the pneumococcus was found seven times, once in connection with Friedlander's bacillus. Terminal infections with staphylococci and streptococci were occasionally found. Lumbar puncture was performed in fifty-five cases and in some of these several punctures were made. Diplococci were found either on microscopic examination or in cultures in thirty-eight of these cases. The average period from the onset of the attack before the puncture was made was seven days in the positive cases and seventeen days in the negative cases. The longest time after the onset in which the fluid drawn by puncture gave positive results was twenty-nine days.

SYMPTOMS OF THE ORDINARY FORMS.

Cerebrospinal fever presents a great diversity of

symptoms in different cases. No other acute disease appears in such various guises. Stillé has well called it a "chameleon-like disorder." The period of incubation is unknown. Prodromes are rare. When present they consist of headache, dragging muscular pains, vertigo and a sense of fatigue. The onset of the attack is usually abrupt. It is marked by a chill, agonizing headache, nausea and vomiting. In some cases headache is not a conspicuous symptom. The attack begins with vertigo and the patient acts like a drunken man. This was the mode of onset in a case which I recently had the opportunity of seeing. Presently dragging pains in the neck come on, which spread along the spine and into the extremities and are soon followed by motor symptoms which progressively develop. These consist of titanic stiffness of the spinal muscles, great pain on attempting to bend the head forward or to turn it from side to side and awkwardness and difficulty in movements of the extremities. Strabismus, inequality of the pupils and palsies of the facial muscles are common. In the course of a little time opisthotonos develops, the head is drawn back, the spine curved, the forearms flexed on the arms and the legs on the thighs. Muscular cramps and spasmodic twitchings come and go. In young children general convulsions may occur.

Hemiplegia has been frequently observed. Paraplegia may also occur. Paralysis may develop during the course of the attack and disappear shortly or it may persist for some time.

The sensory symptoms consist of headache, which may be sharp, lancinating or boring and commonly referred to the back of the head; sometimes it is felt as a constricting band; pain in the back of the neck and in the lumbar and epigastric regions and general hyperesthesia, most marked in the face and neck. Various disturbances of the special senses occur. There is a photophobia, intolerance of sounds, ringing in the ears and vertigo.

The physical disturbances are striking. The patient is restless and distressed. His face is seldom flushed, usually pale and slightly cyanotic. In children there is great irritability. Delirium occurs early and may be active, even maniacal or of a busy wandering character. After a time it passes into somnolence or stupor, which may be still attended by more or less restlessness and continual movement on the bed. In the worst cases stupor deepens to coma.

The symptoms referable to the alimentary tract are not characteristic. The tongue is at first slightly covered with a whitish fur. In conditions of great depression it becomes dry and brown and sordes collect. Taste is lost and the patient refuses food; nevertheless the vomiting persists. Constipation is commonly present throughout the sickness. Toward the end of the attack, however, diarrhea and involuntary discharges may take place. The abdomen, as a rule, is neither tender nor tympanitic. Slight, exceptionally marked enlargement of the spleen may be made out.

The fever is generally moderate, very irregular and does not observe a typical course. There are frequent remissions. In some instances fever is slight or absent altogether. On the other hand, the temperature may reach 105° or 106° F. In some cases it abruptly rises before death. The fever, even when intense, may be of short duration. There is no constant relation between the intensity of the febrile movement and the severity of the other symptoms. Defervescence may take place without improvement in other respects, so that there

are cases in which severe nervous symptoms persist for weeks after the temperature has fallen to normal or even to subnormal ranges. Many of the cases show a temperature range of irregularly remittent type. In the milder cases the temperature is sometimes distinctly intermittent. The subfebrile temperatures are sometimes broken by rapid and transient elevations.

The pulse is also irregular. There is no constant correspondence between the pulse and temperature. It may be soft and weak, even slower than in health, and is often intermittent and arrhythmic. Abrupt changes in the force and frequency of the pulse are common. Change in the frequency from 80 to 100 has been observed in the course of a minute. The rhythm of the respiration may be disturbed and Cheyne-Stokes breathing may occur in the graver cases.

Leucocytosis is present throughout the disease, diminishing toward the end of the attack in cases which recover. In 33 cases in the Boston epidemic in which blood counts were made the maximum number of leucocytes was 31,000, the minimum 9,350. In a case under my observation at the present time there is a leucocytosis of 40,000. Differential counts show that the increase is chiefly due to the polynuclear leucocytes.

Lesions of the skin are common. To their prominence is due the old name of spotted fever. They vary greatly in different epidemics. In many cases they are absent altogether. They have been more frequent in the American and Irish epidemics than on the Continent. They are polymorphous and several different forms may occur in the same case. Herpes is far more common than any other eruption. It usually appears on the lips and nose, but may involve other parts of the face or body and may vary from a crop of a few fine vesicles to an abundant eruption of large vesicles. A petechial rash is frequently observed, and in some instances extensive hemorrhagic areas develop in the skin. The petechiæ often resemble flea-bites. They are distributed in varying numbers over the whole surface, but particularly about the knees and elbows. In some cases the rash is abundant and develops with great rapidity. Patches of erythema, dusky mottlings and rose spots disappearing on pressure, like the rash of enteric fever, have been observed. Among the rarer cutaneous manifestations are urticaria, erythema nodosum, pemphigus and gangrene.

The urine is, as a rule, increased. It may be much increased even with high temperature. The reaction is usually acid. A moderate amount of albumin is frequently present. Recent personal observations lead me to believe that there is a special form of cerebrospinal fever characterized by symptoms of an acute nephritis and corresponding to the renal form of enteric fever—the nephrotypus of the French. I have seen two such cases in which the urine from the onset was densely albuminous, of high specific gravity and contained epithelial and blood casts and free blood. One of these cases terminated fatally; the other is still under observation. Glycosuria occasionally occurs, and in malignant cases hematuria has been observed. Retention of the urine is common in the graver cases. Polyuria is frequent in children, and in some cases has persisted for years after convalescence.

The eye lesions are referable to three causes. First, neuritis, due to the involvement of the nerve in the exudate at the base without extension of the inflammatory process to either the orbit or the eye. This condition may affect the oculomotor and the optic nerve.

Second, inflammation from the meninges may extend directly into the eye along the pia-arachnoid of the optic nerve, causing purulent choroido-iritis and in very rare instances suppuration in the orbit. Keratitis may arise in consequence of an extension of the inflammation from the iris and ciliary region. The third cause is neuritis of the fifth nerve, with destruction of the gasserian ganglion and loss of sensation. To this most of the cases of keratitis and purulent conjunctivitis are due.

Symptoms relating to the auditory apparatus are very common. The auditory nerve is generally swollen and surrounded by the exudate. Extensive degeneration of the nerve fibers is frequently found, the degeneration being most marked in the chronic cases.

The subject has been especially studied by Moos, who found that of 64 cases of meningitis which recovered 59 per cent. were deaf, in 7 per cent. the hearing was impaired and in 31 per cent. it remained normal. Of 43 cases closely investigated, deafness appeared in 2 on the tenth day, 6 on the second, 3 on the third, 17 between the fourth and tenth days and 15 between the fourteenth day and the fourth month. This observer thinks that it is possible that the abortive form of epidemic cerebrospinal meningitis is the cause of many cases of early acquired deafness. Deafness is frequently due to disease of the labyrinth. Otitis media and mastoid disease occur. The diplococci are found in the pus cells.

Coryza has been frequently observed in the course of the attack. Weigert first advanced the opinion that in meningitis the nose forms the portal of entry for the infectious organisms. It may be, however, that their presence is due to an extension from the brain and not to primary invasion. Epistaxis occurs also. In 15 of the Boston cases 10 showed the presence of the diplococci in the nasal secretion, and in 18 cases examined by Scherrer the diplococci were found in the pus cells.

The wasting in severe cases is rapid and extreme. An early, sudden and great loss of strength is a frequent and prominent condition in this disease.

The symptoms may be divided into those due to the inflammatory lesions of the cerebrospinal organs and those due to a general infection. In the malignant cases both these groups of symptoms are of overwhelming severity. In the mild cases the nervous symptoms are predominant. The foregoing symptoms indicate the nature and severity of the disease in its ordinary form.

SYMPTOMS OF THE ANOMALOUS FORMS.

A. *Malignant (Meningitis Cerebrospinalis Epidemica Siderans).*—The patient is struck down without warning and speedily falls into a state of collapse. A violent chill is followed by cyanosis, coldness of the surface, profuse perspiration, intense headache which alternates with drowsiness and brief delirium followed by unconsciousness. There may be contraction of the neck. Respiration is slow and labored; the pulse rapid and feeble; the urine scanty and loaded with albumin. Purpuric blotches appear on the surface. Cases of this kind have occurred in many epidemics and with greatest frequency at the beginning of the outbreak. They may occur sporadically. Death may ensue in the course of a few hours.

B. *Abortive (Meningitis Cerebrospinalis Epidemica Abortiva).*—The onset of the attack is severe. In the course of a few days the symptoms subside and convalescence is rapid. Under this head must be included

the mildest ambulant cases. Patients complain of headache, stiffness in the neck and spine and malaise. Vomiting occurs. Fever is, as a rule, absent. Stillé relates the case of a girl who was fully convalescent on the fifth day, and states that he observed many slight but distinctly marked cases that were convalescent within a week. J. Lewis Smith has reported similar cases in childhood. Cases of this kind can only be recognized in the light of a prevailing epidemic.

c. *Intermittent (Meningitis Cerebrospinalis Epidemica Intermittens.)*—Intermittent and remittent cases are common. Not only the fever, but other symptoms of the disease show extraordinary exacerbations and remissions, which may be repeated at intervals of twenty-four or forty-eight hours. These cases may be due to successive involvement of areas of the meninges or to fresh growths of the organisms. They rarely, however, present the well-marked periodicity of the malarious diseases.

d. *Chronic (Meningitis Cerebrospinalis Epidemica Chronica.)*—Cases of this form occur in all epidemics. The disease lasts, with numerous complications, remissions and exacerbations, for several weeks or in some instances for five or six months. Emaciation is extreme. The symptoms may be due to the persistence of conditions left by the acute attack, such as chronic hydrocephalus or abscess of the brain, or general neuritis.

COMPLICATIONS AND SEQUELS.

Among the complications and sequels are pleurisy, endocarditis and pericarditis. Bronchial catarrh and deglutition pneumonia are very common. Croupous pneumonia has been very common in some of the epidemics. This complication occurs more frequently at the close than at the beginning of an epidemic. This was especially the case in an epidemic observed by Immerman and Heller at Erlangen. It is uncertain whether in some instances the cases of pneumonia reported in connection with epidemic meningitis have been cases of true croupous pneumonia or cases of diplococcus pneumonia. In the Boston epidemic the latter condition was found in 8 cases, in one of which the amount of lung involved was so extensive as to resemble both clinically and anatomically ordinary croupous pneumonia.

Arthritis, commonly slight, but in rare instances suppurative, has been noted. The wrist joints are most commonly involved. I have observed transient, painful swelling, usually involving the wrists or knees, in a number of cases. In the League Island epidemic, among 23 cases there were 8 in which purulent arthritis occurred. Swelling of the parotid glands is an occasional accident of the disease. It may be slight or may run on to suppuration—parotid bubo.

Intestinal catarrh may occur as a complication. Malarial and enteric fever and measles, scarlet fever and cholera have been encountered as intercurrent affections.

The convalescence is irregular and uncertain. After severe cases it is apt to be tardy. Relapses are not uncommon and are often fatal.

Among the more important sequels are prolonged debility and emaciation, palsies and various forms of paralysis, impairment of intelligence in consequence of chronic meningitis and chronic hydrocephalus, especially in children, and more or less complete deafness and loss of vision. General motor weakness and paralysis of individual cranial nerves or of the lower extremi-

ties may persist for a long time. They depend on lesions of the brain or spinal cord or pressure exerted by extensive organized inflammatory exudate, or on peripheral neuritis.

The symptoms of chronic hydrocephalus are paroxysmal. They consist of severe headache, intolerance of light and sound, vertigo, pains in the neck and limbs, vomiting, involuntary discharges, convulsions and loss of consciousness. The attacks occur either at long and irregular intervals or in rapid succession. The mental and bodily condition of the patient during the intervals is sometimes such as to lead to delusive hopes of his recovery.

DIAGNOSIS.

The direct diagnosis of cerebrospinal fever by ordinary clinical methods is a matter of difficulty in sporadic cases and at the beginning of outbreaks. The diagnosis of any form of meningitis is occasionally obscure. Sudden onset, chill, fever, vomiting, delirium, tremor and painful rigidity of the back of the neck may occur in pneumonia, the malignant form of variola, typhus and especially in the cerebrospinal form of enteric fever. The phenomenon described by Kernig in 1884 and known as Kernig's sign is found to be present in 80 to 90 per cent. of the cases of meningitis and only exceptionally present in other cases. This test is often attended by evident pain on the part of the patient. Kernig's sign is not available in cases of rheumatic or other forms of arthritis of the knee or hip, myositis, contractions from nervous disease and sciatica.

If meningitis be present there is usually no great difficulty in recognizing cerebrospinal fever during an epidemic. The ordinary and anomalous forms alike show a symptom-complex that in the course of a little time is distinctive. In doubtful cases and, indeed, in all cases when practicable lumbar puncture should be performed. The method is easy and has been shown by experience to be devoid of danger. If carried out early in the attack, at the time when the diagnosis is often as important as it is difficult, the result is commonly conclusive.

If the patient has meningitis the fluid withdrawn is, as a rule, but not invariably, more or less cloudy; if cerebrospinal fever, the *Diplococcus intracellularis meningitidis* (meningococcus) will be found on direct microscopic examination or in cultures. Positive conclusions can only be drawn from positive results. When the result is negative the operation must be repeated.

DIFFERENTIAL DIAGNOSIS.

The differential diagnosis between cerebrospinal fever and the following diseases demands consideration:

1. *Pneumococcus Meningitis.*—This may occur alone or in connection with croupous pneumonia. The pulmonary lesion may be latent. In fact until very recently this organism was very generally regarded as the cause of epidemic cerebrospinal fever. Symptoms indicating extensive infection of the meninges, of the cord and spinal roots and extension of the infective process along the cranial nerves are less marked or absent altogether. Contraction of the muscles of the neck may be absent, delirium and coma are present and occur early, and this form of meningitis is commonly fatal, while cases of the epidemic form may recover.

2. *Streptococcus Meningitis.*—This form is secondary to infection elsewhere. Fracture of the skull, especially fracture of the base, local abscess formation, acute endocarditis, erysipelas of the face and scalp, otitis media

with extension to the mastoid or the meninges, are forms of primary infection. Opisthotonos is neither common nor well developed. The symptoms develop slowly and are often for a time obscure. The association of painful rigidity, intense headache and vomiting is not conspicuous. Eye symptoms are common.

3. *Tuberculous Meningitis*.—This, perhaps the most familiar form of meningitis, is to be distinguished from cerebrospinal fever by a protracted period of prodromes, more gradual onset, slower course, slow and irregular pulse, great irregularity of the respiration and the absence of eruption. Antecedent tuberculous disease, failure of health following measles or influenza, tuberculous glands or a hereditary predisposition to tuberculous infection are found in the history of the patient. In children or during the prevalence of an epidemic of cerebrospinal fever and in cases in which the tuberculous process involves the spinal meninges (Hirsch) the diagnosis is far from easy.

4. *The Cerebrospinal Form of Enteric Fever*.—The onset is sudden and the symptoms of the period of invasion are those of cerebrospinal fever. About the end of the first week the cerebrospinal phenomena usually subside more or less abruptly and the subsequent course of the illness is that of an ordinary case of enteric fever.

5. *Scarlet Fever*.—In some instances the sudden onset, high febrile movement, vomiting, convulsions and stupor suggest cerebrospinal fever as it occurs in children. The presence of the peculiar redness of the palatine half arches rapidly followed by general erythematous angina are important. In the course of twenty-four or thirty-six hours the efflorescence will clear up any uncertainty.

6. *Typhus Fever*.—At one time cerebrospinal fever was confounded with typhus or regarded as a variety of that disease. To Stillé is due the credit of having finally settled every question of doubt concerning the identity of these two diseases in this country. They are in strong contrast in respect of their causes, symptoms, course, lesions and sequels.

7. *Hysteria*.—Cases of cerebrospinal fever occurring in nervous females at the close of epidemics or sporadically have presented a delirium so peculiar and an array of symptoms so little characteristic that they have been looked on as manifestations of hysteria. This error in diagnosis is no longer possible.

PROGNOSIS.

In individual cases the prognosis can never be made with certainty. The abortive and fulminant cases run the most rapid course. Hirsch has emphasized the fact that certain cases which at the onset present the symptoms of cerebrospinal fever recover after an illness of a few hours, which terminates in free sweating. The malignant cases, on the other hand, prove fatal in a few hours or two or three days. Moderately severe cases may last one or two weeks or several months. The first week is the time of greatest danger. Symptoms rendering the prognosis unfavorable are intense excitement, early depression, persistent vomiting, irregular respiration and convulsions alternating with coma. The average mortality is about 40 per cent. It varies in different epidemics from 20 to 75 per cent. The mortality in the Boston outbreak was 68 per cent.

Relapses are not infrequent and are often fatal.

Bad as the prognosis in cerebrospinal fever is it is still worse in pneumococcus, streptococcus and tuberculous meningitis, these conditions almost invariably terminating in death.

PROPHYLAXIS.

In times of epidemics cleanliness of dwellings and the avoidance of overcrowding should be instituted in the hope of diminishing the severity and mortality of the disease. The successive attack of different members of a household demands the abandonment of the dwelling and thorough disinfection. Isolation of the patients and sterilization of their clothing should be carried out.

TREATMENT.

The mildest cases require no treatment; the malignant cases react to none. Medicine with all its resources is neither capable of combating the attack nor responsible for its results. Of this group of cases Stillé has said "the first symptoms of the disease are the first phenomena of death."

The room occupied by the patient should be darkened and quiet. The diet should consist of milk, eggs, meat juice and broths administered at regular intervals. When food can not be taken by the mouth, nutritious enemata should be given. Water should be freely administered. Alcohol is to be given when the symptoms of depression appear, whether at the onset of the attack or later. A guide as to its amount is the effect on the character of the pulse and the first sound of the heart. Venesection, generally practiced in the early history of the disease, is in no case admissible. On robust individuals wet cups may be applied to the nape of the neck and along the spine or leeches to the temples and in the neighborhood of the mastoid process. The continuous direct application of cold to the head and spine by means of ice-bags is of value. The pain is relieved and in some instances quietude and sleep ensue. If the temperature be high systematic cold bathing may be practiced. During the application of cold to the head and spine and after cold bathing the patient should be covered with warm blankets and artificial heat applied to his sides, thighs and extremities. Warm baths, 104° F., of ten minutes' duration, have been recommended. Blisters applied to the occiput and nape of the neck early in the attack have appeared to be of service. They should be carefully watched and immediately removed on the development of signs of vesication, which is liable to be followed by troublesome sloughing. The thermo-cautery may be used.

Laxative doses of calomel should be given early in the attack. Quinin has been extensively employed. There is no evidence that it exerts a favorable influence, much less that it is capable of aborting the disease. Opium may be regarded as one of the indispensable remedies in the treatment of epidemic meningitis. The indications for its use are to be found in the headache and spinal pains, restlessness, spasm, hyperesthesia and sleeplessness. It should be given early and in full doses. Experience has demonstrated a remarkable tolerance for this drug in cases of cerebrospinal fever. Strong gave in one case 60 drops of laudanum every half-hour until half a fluid ounce was taken. Some of the early American physicians gave enormous doses, half an ounce of the tincture of opium or from 30 to 60 grains of the crude drug in the course of twelve hours. These large doses were frequently followed by recovery. Stillé gave one grain every hour in grave cases and every two hours in moderately severe cases without even an approach to narcotism. He adds that "under the influence of the medicine the pain and spasms subsided, the skin grew warmer and the pulse fuller, and the entire condition of the patient more hopeful." Morphia may be administered hypodermically

in doses of from 1/6 to 1/2 grain. With great restlessness and jactitation hyoscin hydrobromate is indicated. In the case of children the dose must be cautiously regulated.

Ergot, belladonna, cannabis indica, gelsemium, chloral, zinc oxid and the bromids are of inferior value. They are frequently, however, given in connection with opium or morphia. Inhalations of ether or chloroform may be cautiously employed to secure temporary relief from torturing pain, jactitation and spasm. Veratrum viride and aconite are mentioned only to condemn their use in this disease. The modern analgesics may be employed alone or in conjunction with opium and its derivatives in the management of the milder cases of the disease. Large doses are to be avoided.

R. Phenacetini 3|
Chart No. viii.
Div. S. One powder q. tertia vel quarta hora.

R. Pectophenini 2|
Sacch. alb. 1|
Chart No. vi.
Div. S. One powder q. t. v. q. h.

R. Antipyrini 3|
Sacch. alb. 1|
Chart No. viii.
Div. S. One powder q. t. v. q. h.

If the headaches and pains increase cerebrospinal fluid may be withdrawn by lumbar puncture in order to diminish the intracranial pressure. Performed early in the course of the disease and repeated at intervals, it has been frequently followed by beneficial results.

Manges has reported 3 successful cases treated by the injection of lysol in 1 per cent. solution into the spinal canal after the removal of the exudate. This method was originally suggested by Carlos Franca at Lisbon in 1903. Wolff, of Hartford, Conn., basing the procedure on an antagonism which he found to exist between the *Diplococcus intracellularis* and the Klebs-Loeffler bacillus and the fact that the former were precipitated from active bouillon cultures on admixture with diphtheria antitoxin serum, was led to use the antitoxin in the treatment of this disease with satisfactory results, and Waitzfelder² treated 17 cases in the same manner, employing from 4,000 to 10,000 units, according to the age of the patient and the severity of the symptoms, at intervals of two or three days. Of his 17 cases, 5 recovered completely, 3 died and 9 were still under observation at the time of the publication of the report, 5 of the last "giving promise of speedy recovery." These observations are now being extensively repeated. In two cases of which I have knowledge the antitoxin treatment was employed without result, but these two cases were of fulminant type, one dying at the end of forty-eight hours and the other in a little more than twenty-four hours.

During a period of depression stimulants are to be freely given. Preparations of alcohol, ammonium carbonate and the spirit of chloroform are recommended. Cold effusion may be occasionally practiced.

During convalescence the bitter tonics and iron may be given. Cod liver oil and iodid of potassium are also employed. Delirium, spasm and irritability of the stomach too often in the severe cases render the administration of medicine and food impracticable. As recovery takes place the appetite is often excellent and an abundant, easily digested and nutritious diet of mixed solid food should be given.

THE ACTION OF THE INTRACELLULAR POISON OF THE COLON BACILLUS.*

VICTOR C. VAUGHAN, JR., M.D.

ANN ARBOR, MICH.

It is now generally conceded that the pathogenic bacteria bring about those changes in the body cells which form the basis of what we call disease by virtue of certain poisons which they produce. These poisons are supposed to be specific for each kind of bacteria and to them the name of toxin has been given. Certain bacteria possess a toxin which is not an integral part of the cell itself, and hence is not essential to its existence. As examples of such organisms may be mentioned the bacillus of tetanus and that of diphtheria.

With these bacteria the toxin is found free in the culture medium and there are differences in opinion concerning its origin, some regarding it as a cleavage product formed by the action of the bacillus either directly or through a chemical ferment on the constituents of the culture medium, while others think that the soluble toxin represents a group of the bacterial cell which easily drops off. On the other hand, the filtrates of cultures from such bacteria as the typhoid and the colon bacillus are non-toxic, or, at the most, possessed of but feeble toxicity. In the later instance the poison exists as an integral portion of the bacillus itself and can be set free only on the death of the latter. We therefore have two distinct classes of so-called specific bacterial poisons, the one being easily cast off from the living cell and hence called an extracellular toxin to distinguish it from the toxin which can not be split off from the bacterial cell without destroying the latter, the intracellular toxin.

That the colon bacillus, when introduced into the animal body, is one of the most deadly of all bacteria is a fact which is too well known to require further proof. There is no other germ which causes more anxiety to the abdominal surgeon, and the fatal results following typhoid perforation, strangulated hernia, and appendicitis when left to pursue their course uninterruptedly are in almost all instances due to its ravages.

The colon bacillus is a normal inhabitant of the intestinal canal, and so long as it remains within the canal it is practically outside of the body and can do no harm. When it obtains an entrance into the peritoneal cavity, however, it multiplies and at the same time is disintegrated by the protective agencies of the body. As it is broken up, its intracellular poison is set free and then, and not until then, is it capable of acting on the body of the host. It is not, therefore, from the growth of the bacterial cell in the peritoneal cavity that death directly results, but rather from its destruction. Primarily, of course, it is essential that the growth of the organism should have advanced to such a degree that enough bacterial cellular substance shall be present to furnish a sufficient quantity of the poison to cause a fatal issue in the body on its liberation. The necessity of employing a large amount of the living culture in order to obtain a fatal dose of poison from it has long been urged as one of the principal arguments against the intracellular poisons being true toxins. As compared with diphtheria and tetanus, this is, indeed, true, but the comparison is hardly a fair one. Thus, in the case of the diphtheria bacillus, the same germ may go on producing toxin without loss of life, for all one knows, indefinitely, whereas, in the case