The Journal of the American Medical Association

Published under the Auspices of the Board of Trustees

VOL. XL.

CHICAGO, ILLINOIS, MARCH 21, 1903.

No. 12.

Original Articles.

OBSERVATIONS ON THE MECHANICS OF DIGESTION.*

WALTER BRADFORD CANNON, M.D.
INSTRUCTOR IN PHYSIOLOGY, HARVARD MEDICAL SCHOOL.
BOSTON.

The results which are to be presented in outline in this paper have been obtained in researches carried on in 1897-98 and in 1901 in the Laboratory of Physiology of the Harvard Medical School, and they are part of an extensive investigation into the mechanical factors of digestion to be carried on during the next few years. The method thus far pursued has been that of mixing with the food subnitrate of bismuth, a harmless, almost inert, powder, opaque to the *x*-rays, and then with the fluorescent screen observing the changes in the shadows of the alimentary contents as they are worked on by the muscles of the digestive organs. The clearness with which masses of food can be seen is indicated in Figure 1, a radiograph of a cat. Observations have been made chiefly on cats, but many of these observations have been confirmed on dogs, rats and guinea-pigs, as well as on the human subject. The results of the work on the movement of food in the esophagus,¹ on the movements of the stomach² and on the movements of the intestines³ have already appeared. In this paper I shall review briefly some of the results already published on the activities of the stomach and intestines and report certain new observations.

The mechanical factors in digestion perform three functions: the propulsion of the food through the alimentary canal, the mixing of the food with the digestive juices and the exposure of the digested food to the absorbing mucous membrane. These functions are performed in different ways in different parts of the alimentary tract. For the sake of clearness, the methods and results of the activities of the stomach will first be considered, and later those of the large and small intestines.

When the stomach is observed according to the method above mentioned, it is seen to consist of two physiologically distinct parts, divided about half way between the cardia and the beginning of the antrum (Fig. 2). To the left of this division lies the cardiac portion or fundus of the stomach; to the right is the pyloric portion. The distinction between these two portions arises

from a difference in muscular activity. Over the food in the pyloric portion waves of constriction are repeatedly passing. These waves start at the beginning of this division of the stomach and pass as deepening contraction rings to the pylorus. In the cat six of these waves pass a given point in a minute, in the white rat, also, about six waves per minute, in the dog between four and five, and in the guinea-pig seven. The fundus is characterized by a total absence of peristalsis; as the food is pressed out of the pyloric portion, however, by the repeated constrictions, the muscles of the fundus slowly contract and squeeze the contents into the more active division. Since the stomach is an organ more devoted to digestion than to absorption, the mechanical processes of mixing the food with the digestive juices and of propelling the food into the intestines are here of first importance.

The efficiency of the gastric peristalsis as a means of mixing the food with the gastric secretions depends on the contraction of the pyloric sphincter. Careful observation reveals that the pylorus does not open for every wave of constriction which approaches it, but only at irregular intervals. So long as the sphincter holds tightly, each ring of constriction as it courses from the middle to the end of the stomach presses the food into a blind pouch; the food, unable to escape through the pyloric opening, has as its only outlet the opening in the advancing ring. Thus each peristaltic constriction in moving toward the pylorus has all the contents of the pyloric portion streaming back through the contracted ring. Beaumont, in his famous researches on Alexis St. Martin, declared that the gastric contents circulated first along the greater curvature from cardia to pylorus, then back along the lesser curvature. He gave, however, very meager evidence to support this statement. Brinton, for theoretical reasons, believed that the food must move forward along the sides of the stomach and backward through the middle. The value of the circulation of the food, as described by these men, lay in the supposition that the food was thus brought near to the secreting gastric wall and that the gastric secretion could thereby more readily exert its action. Observations made with the *x*-rays do not support these theories of mixing currents running throughout the stomach, but nevertheless they show that the pyloric portion is an admirable device for bringing all the food under the influence of the glandular secretions of that region.

Little paste pellets of bismuth subnitrate with starch enough to keep the form may be distinguished from the remainder of the stomach contents by the darker shadows which they cast. Several times I have been fortunate in seeing two of these pellets about a centimeter apart in the axis of the stomach. As a constriction wave ap-

* Read at the Fifty-third Annual Meeting of the American Medical Association, in the Section on Pathology and Physiology, and approved for publication by the Executive Committee: Drs. Alfred Stengel, L. Hektoen and Winfield S. Hall.

1. Cannon and Moser: *American Journal of Physiology*, 1898, 1, p. 435.

2. Cannon: *Ibid.*, 1898, 1, p. 359.

3. Cannon: *Ibid.*, 1901, vi, p. 251.

proached them, both moved forward, but not so rapidly as the wave. When the constriction overtook the first ball, the ball moved backward through the constricted ring in the direction of least resistance. The wave then overtook the second ball and this also passed backward to join its fellow. At the approach of the next wave they were both pushed forward once more, only to be again forced backward, one at a time, through the narrow orifice. As the waves recurred in their persistent rhythm the balls were seen to be making progress, an oscillating progress, toward the pylorus; for they went forward each time a little farther than they returned. Sometimes twelve minutes elapsed before the balls had advanced to the pylorus from the place where they were first

Fig. 1.—Radiograph of a cat's abdomen, taken 5½ hours after eating. Spinal column in middle line with pelvis below. On right, above, the pyloric end of stomach; on left, dimly outlined by gas, the ascending colon, and the cecum containing small amount of food. Loops of small intestine containing food in various parts of the abdomen.

affected by the waves; since the waves were running at the rate of six per minute, these little pieces of food were moved back and forth on their way to the pylorus more than seventy times.

The significance of the motions of these food particles is made clear when the conditions attending the formation and passing of a peristaltic wave are considered. As a constriction occurs, the secreting surface enclosed by the narrowed muscular ring is brought close around the food lying within the ring. As this constriction passes on fresh areas of glandular tissue are continuously pressed inward about the narrow orifice; and, furthermore, as this constriction passes on, a thin stream of food is continually forced back through the orifice and

thus past the mouths of the glands—a food stream sweeping in one direction past an ever-refreshed mucous surface moving in the opposite direction. The result of this ingenious mechanism is that every part of the secreting surface of the pyloric portion is brought near to every bit of food before the latter leaves the stomach, it may be seventy times or more, as shown by the moving balls.

The propelling function in the stomach is subserved by these same waves of constriction whenever the pylorus relaxes. Relaxation of the pylorus occurs at irregular intervals and is due to factors as yet little understood. Curiously, it allows carbohydrate food, on which the stomach does not act, to pass onward into the intestine much sooner than it releases a proteid food which the stomach can digest. Hard particles, when brought by the peristaltic waves to the pyloric sphincter, act as a mechanical stimulus and greatly prolong the periods of its contraction. Under these circumstances over and

Fig. 2.—Outline tracing from radiograph of cat's stomach. *x-y*, Line of division between cardiac and pyloric portions; cardiac portion, with even contour, pyloric portion with waves of constriction passing to the pylorus (*p*); *o*, position of the esophageal opening.

over again the hard particle is swept up to the pylorus, only to shoot back through the advancing ring of constriction. By this repeated rubbing the peristaltic waves of the stomach serve to break up the more solid parts of the food, a function which the teeth perform with manifestly greater efficiency. It may be that the prevalence of pathologic conditions in the pyloric part of the stomach rather than in the fundus is due to injuries which the greater activity of the pyloric part must bring on itself.

The fundus of the stomach, as stated above, is characterized by a slow tonic constriction of its musculature, which forces its contents into the more active pyloric portion. Pellets of subnitrate of bismuth in this part of the stomach contents retain their relative positions for an hour or more after the food is ingested. In 1898 I presented evidence that the absence of mixing currents in the fundus results in the food there retaining its original reaction for a considerable period, so that salivary digestion may readily take place in that part of the stomach without interference by the acid gastric secre-

tion. These results have recently been confirmed by Heyde⁴ working in Grützner's laboratory in Tübingen.

The value of the stomach as a place for holding the bulk of the food and serving it out a little at a time so that the intestines may not become congested during their digestive and absorptive processes has been pointed out by Moritz. All the advantages secured thus to the intestines may be claimed also for the stomach itself, for, as has been shown, the active portion of the stomach is the pyloric portion, and the fundus serves as its reservoir—an active reservoir pressing out its contents a

Fig. 3.—Radiograph showing region of persistent rings of contraction (descending colon), and region of antiperistalsis (cecum, ascending and transverse colon). The slight indentation above the cecum marks the position of the ileocolic valve.

little at a time as the pyloric peristaltic mechanism is ready to receive them.

I have described in detail the effects of peristaltic waves in bringing the food into intimate contact with the mucous surface of the stomach, because the action is similar to that in the cecum and in the ascending and transverse colon, where the normal activity is an antiperistalsis. Constriction waves start in the transverse colon and pass backward toward the cecum. These waves, unlike those of the stomach, do not run in continuous rhythm; they occur usually in periods lasting

about five minutes and recur every fifteen or twenty minutes.

The antiperistalsis in this part of the large intestine seems indeed to give a reason for the presence of a valvular structure at the entrance of the small intestine into the large. Hundreds of antiperistaltic waves have been seen coursing toward the cecum, and only twice has food been observed to be pressed back into the ileum through the ileocolic valve. Inasmuch as the valve is competent for the food which has gone from the small into the large intestine, the antiperistaltic waves have the same effect here as the peristaltic waves have in the stomach when the pylorus remains closed. For, when a constriction occurs, some of the mucous surface of the colon becomes enclosed by the narrowed muscular ring; and, as this constriction passes on, fresh areas of this surface are continually pressed inward around the narrow orifice, while a thin stream of food is passing by it in the opposite direction. As waves recur about five times a minute, twenty-five waves or more affect every particle of food in the cecum and in the ascending colon in this churning manner during each normal period of antiperistalsis. The result must be again a thorough mixing of the contents with the digestive secretions brought from the small intestine and an exposure of the digested food to the absorbing wall.

Immediately after food passes from the ileum into the large intestine a strong tonic contraction of the cecum and the ascending colon is commonly observed, which serves to press onward toward the rectum the contents of these parts. Antiperistaltic waves follow at once the general contraction, so that much of the food which has been pressed onward is returned into the cecum. With the repetition of this process, however, more and more material appears at the end of the transverse colon, and on its appearance there a persistent ring of contraction cuts it off from the region of antiperistalsis; as still more food appears in the large intestine this ring moves slowly onward toward the rectum, pressing a mass before it, and is followed by other similar rings carrying onward similar masses by very slow peristalsis (Fig. 3).

Thus, in the large intestine the function of mixing the food with the digestive secretions and of exposing this food to the absorbing walls is performed by an antiperistalsis at the beginning of the large intestine. It is here that the last valuable constituents of the food are worked over and taken into the body. The remnant of unused material is propelled onward by occasional strong contractions of the whole region of antiperistalsis. These contractions squeeze the material toward the end of the transverse colon where slowly-moving peristaltic waves force it to the rectum.

The manner in which the food is mixed with the digestive secretions, exposed to the absorbing wall, and carried forward in the small intestine is considerably different from the processes observed in the large intestine and in the stomach. To the admirable process by which these functions are performed in the small intestine I have given the name "rhythmic segmentation." This activity of the muscular wall is first seen in the duodenum when a mass of food has accumulated there after repeated relaxations of the pylorus. The mass of food is observed at first to be wholly quiet (Fig. 4). Suddenly irregular movements take place about it, and then at regular intervals along its length constrictions of the circular musculature separate the mass into a number of segments of equal size. Hardly have the constrictions thus taken place when similar constrictions begin to appear

4. Grützner: Deutsche Medizin-Zeitung, 1902, No. 28.

about the middle of each segment. As these new constrictions deepen, the first constrictions begin to relax. Finally, when the new constrictions have completely divided the segments, the first constrictions have entirely relaxed and the neighboring halves of the divided segments unite to form new segments in the region of the first constriction. Now these new segments are again divided by circular constrictions about their middle, and neighboring halves of these divided segments unite to make a third series, and so on. This process of rhythmic segmentation of the food may go on for several minutes in the duodenum (at the rate of thirty divisions per minute in the cat) so that the food, coming from the stomach, becomes thoroughly mixed with the outpouring secretions of the liver and the pancreas. It is usual for the process to be brought to an abrupt end in the duodenum by active peristalsis. A peristaltic wave forms behind the mass of food and sweeps it swiftly forward through several coils of the intestine.

When an animal has received a meal of bread and milk or salmon the characteristic appearance of the alimentary organs is as follows: over the pyloric end of the stomach constriction waves are continually passing.

Fig. 4.—Diagram representing the process of rhythmic segmentation of the contents of the small intestine. Constrictions cut the mass into a series of uniform ovoid segments; a moment later each segment is divided into two parts, and immediately after the division neighboring parts, as *a* and *b*, line 2, rush together and merge to form new segments, as *c*. The next moment these new segments are divided to form a third series, and so on for a thousand times or more.

Along the cecum and the ascending colon an occasional series of antiperistaltic waves is to be seen, and in the small intestine a mass of food here and there is undergoing rhythmic kneading. The quiescence of the food in the intestines is in marked contrast to the continuous churning which it is receiving in the stomach. The food lies in scattered, long, slender masses in the ileum; only a few of these masses show any signs of muscular activity affecting them. When segmentation begins on one of these masses it follows the same course as that described for the duodenum. The only variations from this type are divisions of the segments into thirds instead of halves when the masses are large, or a division into two parts and the subdivision of each part before the mass is united again to undergo a repetition of the process. In the cat, as already stated, segmentation of the food occurs about thirty times in a minute. These constrictions may continue for half or three-quarters of an hour without moving the food perceptibly along the canal. The food is thus divided and redivided over and over again a thousand times or more with no advance. This same operation has been observed in the dog at the rate of from 18 to 22 segmentations per minute and in the white rat at the rate of about 44 to 48 segmentations

per minute—a most remarkable activity for smooth muscle.

The result of the process of rhythmic segmentation must be a thorough mixing of the food with the digestive secretions brought from the stomach and the duodenum and with the secretions of the intestine itself, and a thorough exposure of the digested food to the absorbing walls—since, with each constriction, the absorbing mucous membrane is plunged directly into the midst of the little masses of food. Not only are the processes of digestion and absorption favored by these movements, for Mall has shown that contraction of the intestinal wall has the effect of forcing the venous blood from the submucous plexus into the radicles of the superior mesenteric vein, and has noted that it must have a similar effect on the flow of the lymph. Thus by this single mechanism of segmentation the food and digestive fluids are churned together, the absorbing mucosa is brought in most intimate contact with the digested food, and the blood and lymph, carrying the absorbed material, are sent away through veins and lacteals.

A review of the activities of the small intestine shows that the functions of mixing the food with the digestive juices and exposing the food to the absorbing structures is performed by the process of rhythmic segmentation. The function of propelling the food along the canal may also be performed to a very slight extent by this same process; it is performed typically, however, by the well-known peristaltic wave which has so long been regarded as the chief activity of the intestinal muscles.

There are two matters of practical importance which remain to be mentioned: the effect of antiperistalsis of the colon on rectal injections, and the effect of emotional states on the activities of the alimentary canal. About eight years ago Grützner⁵ reported observations which indicated that materials injected into the rectum might be carried back into the small intestine even as far as the stomach. Since that time some observers have confirmed Grützner's results; others, failing to confirm them, have affirmed that such results were due to carelessly allowing the test materials to become mixed with the food which the animal took normally by mouth. The presence of antiperistalsis in the colon led me to believe that there exists here an activity which would fully account for Grützner's results. Accordingly, I injected into the large intestine enemata of different amounts and of different consistencies. The experiments showed that, when a small amount of the nutrient material was introduced, it lay first in the descending colon. In every instance antiperistaltic waves were set going by the injected mass and the material was thereby carried in considerable amount to the cecum. Never in my experience have these small enemata been carried back through the ileocolic valve. When large injections are given, i. e., amounts nearly filling the colon and cecum, the antiperistaltic waves are effective in forcing the food from the large intestine into the ileum. The ileocolic valve,

5. Grützner: Deutsche medicinische Wochenschrift, 1894, xx, p. 897.

which is competent for food coming normally from the small intestine into the large, is curiously incompetent for this injected substance, even when the substance is of the consistency of thick cream. Radiographs taken at intervals after the injection show that, as time passes, the antiperistaltic waves almost completely empty the transverse and ascending portions of the colon and that the small intestine gradually receives more and more food. After a short time the typical segmenting movements can be seen in the loops, busily separating the food into small masses and again and again dividing and re-dividing them. It is evident that as far as the mechanical activities of the small intestine are concerned this portion of the alimentary canal is bipolar. Food material may enter it at either end; injected food and the nutriment which has come through the stomach in the normal manner are exposed to the same mixing and absorbing activities.

Observations on the stomach of the cat, made in 1898, showed that the peristalsis is stopped whenever the animal manifests signs of anxiety, rage or distress. The extrinsic innervation of a large part of the intestinal tract is the same as that of the stomach, and it is of interest to note that emotional states have the same effect on the movements of the intestines as they have on the movements of the stomach. Any signs of emotion, such as violent rage or even the restlessness and continued whining which indicate anxiety and distress, are accompanied by a total cessation of the segmenting movements in the small intestine and the antiperistalsis of the colon, as well as the total quiescence of the gastric mechanism.

DISCUSSION.

DR. A. C. CROFTAN, Chicago—To what extent did the presence of bismuth in large quantities influence these movements? I believe that some investigators deny the existence of antiperistaltic movements. I would like to ask if bismuth, which would act as a foreign body in the intestine, would modify the intestinal movements?

DR. J. E. O'BRIEN, Scranton, Pa.—I have thought for a long time that many cases of appendicitis were due to loading of the cecum, and this brilliant illustration of the antiperistaltic movements of the ascending colon should shed some light on the subject. Some cases of appendicitis, I believe, begin as congestion of the cecum and extend to the appendix.

DR. W. B. CANNON, Boston—As regards the antiperistalsis of the colon, the animals on which these experiments were performed were kept without food for a day, and I have no means of knowing what takes place except in the food containing bismuth. When food enters the colon, antiperistaltic waves may be seen coursing along the food. This may continue for about five minutes, then there will be an interval of fifteen minutes—the antiperistalsis has been seen starting from the first of the constrictions which you see in one of these photographs of the large intestine. These waves start from the end of the food as it is gradually accumulating in the cecum or the first of the tonic constrictions which occur after the food has come to the further end of the transverse colon. Nothnagel has denied antiperistalsis, but his investigations have only referred to the small intestines, and I agree with him to this extent. In the large intestine, however, there is no doubt that it takes place, for I have seen it in animals whose abdomens have been opened under salt solution. I have also seen it in the dog, and think it probably bears a relation to animals having large ceca. The bismuth does not seem to have any effect on these waves, but this is a critical matter, on which I am working at the present time. I intend adding to the food other substances which are not readily soluble, and which have a sufficient molecular weight to cast a shadow on fluoroscopic examinations.

THE CLINICAL MANIFESTATIONS OF HYDROPHOBIA.*

D. J. MCCARTHY, M.D., AND M. P. RAVENEL, M.D.

From the William Pepper Clinical Laboratory (Phoebe A. Hearst Foundation), and the Laboratory of the Pennsylvania Live Stock Sanitary Board.
PHILADELPHIA.

The usual text-book descriptions of hydrophobia give a clinical course of the disease divided into three stages. It is, however, difficult in practice to separate the clinical manifestations by any fixed rule.

The incubation period is usually about six weeks. In both the cases reported in this paper it was about that length. The onset of the disease is marked by several days of malaise, with slight fever, anxiety and, in a large number of cases, irritation at the seat of the infecting wound. The temperature at this stage is usually elevated one or two degrees, although it may remain at normal. This stage of irritation soon passes over to the convulsive stage. In some cases there is no convulsion or spasm, and the premonitory stage is followed by the paralytic symptoms. The symptoms of the second stage begin with more or less difficulty in the act of swallowing. Anything that excites the swallowing reflex mechanism causes an overflow spasm to the glottis, the muscles of the neck and sometimes to the jaw muscles. As the disease progresses the local spasm is replaced by a general convulsion. The mental state is one of increasing anxiety, both as to the outcome and as to other conditions, until a maniacal condition supervenes, or much more frequently the mental state of a hallucinatory delirium.

The patient may die in the convulsive stage from exhaustion and heart failure or a paralytic group of symptoms supervenes. This so-called paralytic stage may occur in cases where there is very little spasm of the muscles. In these cases it is spoken of as dumb or paralytic rabies. The paralysis is usually one beginning in the lower extremities and later affecting the other extremities. In the delirium there may be passing of the urine and stools in the bed. Some authors include among the paralytic group of symptoms the evidence of cardiac and respiratory failure.

ILLUSTRATIVE CASES.

The following two cases give a picture of the disease as it is met in private and hospital practice.

CASE 1.—W. M., male, aged 5, had been perfectly well up to May 13, 1901. At that time he was bitten on the cheek by a stray dog. The wound healed and he appeared to be in good health until Saturday, June 15. During that night he cried during his sleep. The following day he was up and about, but did not appear to be well. He ate very little during the day. Citrate of potassium was administered by the mother; free purgation resulted. Sunday evening he did not appear to be able to drink, as he asked for water and then told them to take it away after attempting to drink it. Monday he was admitted to St. Joseph's Hospital, with a temperature of 101. He was very much frightened, according to notes of the case, evidently the irritative stage of the disease. He was tested as to his ability to drink, and every time he attempted to swallow there resulted a spasm of the glottis and the muscles of the neck and face. He refused to repeat the first attempt. After reassuring him, he again attempted to swallow a mouthful of water, but it came out through the nose by the spasm of the neck, throat and face muscles as in the first attempt. Later the nurse succeeded in getting him to swallow a portion of an apple. In attempt-

* Read at the Fifty-third Annual Meeting of the American Medical Association, in the Section on Pathology and Physiology, and approved for publication by the Executive Committee: Drs. A. Stengel, Winfield S. Hall and Frank B. Wynn.