should be instructed to distinguish and guard against outward rotation of one eye with diplopia, etc.—a quickly harmful procedure. *f.* So fast as adduction-power grows, the strength of the prisms should be increased. This will usually be by 3 degrees or 5 degrees at a time, and every two or three days, until the limit is reached, esophoria induced, or the symptoms disappear. *f* have sometimes carried the adduction-power as high as 80 or 100 degrees before the desired result appeared. *g.* With a wished-for increase of adduction-power there is likely to be a change of ocular pressures and corneal curvatures so that a retesting of the ametropia becomes necessary.

It seems useless to take up your time with a brave array of case-histories. Hundreds, perhaps thousands, could be collated. The lessons derivable from them is all that seems necessary. The chief ones I gather from mine are as follows:

The infinite patience and clinging to hope on the part of the physiologic and neurologic ocular mechanism is a constant source of wonder and delight, and teaches the daily law to help Nature along the lines of her desires by common-sense and physiologic methods. Her motto is surely *nil desperandum*, and it should be ours. We shall rarely fail when we try to aid Nature's method of spontaneous therapeutics. We should beware of our impertinent surgical short-cuts. Surgery is the despair of medicine, and we should never adopt surgical methods while there is a glimmer of hope by natural means and in the directions indicated by the subtle and inherent strivings for normality. To show but one case illustrative of what Nature and the doctor may do when they work together let me cite:

CASE 2870.—A physician's little girl of 7 came to me with 28 degrees of exophoria, i. e., her exophoria was really exotropia, the divergent strabismus being habitual except under the stimulus of the violent will to converge, when binocular vision was possible for only a few seconds. With a 3-degree prism, base out, this temporary power was impossible. I would not consent to operation. It took about four years of treatment, but to-day the child has slight esophoria, constant binocular vision, perfect acuity, and absence of all eye strain.

Even in cases of positive strabismus natural or non-surgical methods may prove effective if the patient is taken in hand sufficiently early in life, before amblyopia has become too pronounced, and before morbid habits have been too long continued.

THE OPERATIVE TREATMENT OF HETEROPHORIA.*

G. C. SAVAGE, M.D.

PROFESSOR OF OPHTHALMOLOGY, VANDERBILT UNIVERSITY.
NASHVILLE, TENN.

On what cases of heterophoria should we operate? On what muscle should the operation be done? What kind of operation should be done? These are all questions that should be answered by the operator before undertaking this class of work. Certainly, an innervational heterophoria should never be operated upon. A pseudo-esophoria is always curable by convex glasses. A pseudo-exophoria is likewise curable by concave glasses. There are cases that present themselves now and then in whom there is a want of convergence innervation, i. e., there seems to be a fault in the third conjugate innervation.

* Read at the Fifty-second Annual Meeting of the American Medical Association, in the Section on Ophthalmology, and approved for publication by the Executive Committee of the Section: Drs. J. A. Lippincott, Casey A. Wood and H. V. Würdemann.

This condition, as a rule, should be treated by exercise, though occasionally an operation must be done.

No operation should be done in cases of heterophoria, intrinsic in character, that can be relieved by gymnastic exercise, even though this treatment should have to be resorted to for a long while. Nor should those cases be operated upon whose muscle error is small, and for whom comfort can be obtained by the wearing of weak prisms in positions of rest.

The heterophorias not curable by correction of errors of refraction, by prisms in positions of rest, or by rhythmic exercise, should be subjected to operative procedure. Such cases are not infrequent, and the relief from operations skilfully done is by no means uncertain. There is no department of surgery that requires more care in the making of the diagnosis. The condition of every extrinsic ocular muscle must be determined before any one muscle is to be operated upon. There are but two objects in view in muscle operations, the one is altering the tension of the muscle, the other is changing its plane of action.

In order that I may be the better understood when I speak of the operation involving the changing of the muscle plane, it would be well to define it. The rotation plane of any muscle cuts the center of origin of the muscle, the center of rotation of the eye, and the center of the muscle insertion. The axis of rotation is always at right angles to this plane. The plane of an internus or an externus muscle may coincide with the horizontal plane of the eye, when, of course, the vertical axis of the eye becomes the axis of rotation. In such a condition the contraction of an internus or externus will effect only one motion, viz., the rotation of the eye directly in, by the internus, or the rotation of the eye directly out, by the externus. When the center of attachment of the internus is above the horizontal plane, no longer can the muscle plane coincide with the horizontal plane, and a contraction of the internus results in a complicated movement of the eyeball. An internus thus attached not only turns the eye in, but also turns it up and torts it in. The center of the insertion of an externus above the horizontal plane results always in a complicated movement, turning the eye out, turning it up and torting it out. When the centers of attachment of these muscles are below the horizontal plane, the secondary results of their action are the reverse of those mentioned, i. e., a too low internus turns the eye down as well as in, and torts it out; a too low externus turns the eye down as well as out, and also torts it in.

The tension of a muscle is to be altered either by a central partial tenotomy, as when operating on the too strong muscle; by shortening the muscle in the line of its original plane or by advancing it straight forward, as when operating on the too weak muscle. In making either one of these operations the existence of a cyclophoria must be first excluded. When there is cyclophoria complicating any one of the other heterophorias, the operation on the rectus muscle should alter the tension of the muscle and at the same time change the plane of its action. In such a case a partial tenotomy should not be central only, but should include those peripheral fibers, a division of which would be corrective of the cyclophoria. A shortening should be done in such a way as either to raise or depress the plane of action of the muscle as might be indicated by the complicating cyclophoria. In making advancements, the new attachment should be

carried either higher or lower than the original attachment, as the character of the cyclophoria might determine.

The operation to simply alter the tension of a rectus muscle with the view of lessening its power is a central partial tenotomy. This, as in all operations, should be done under strict aseptic precautions, and while the eye is under the influence of cocain. If the operation is to be bloodless, a drop or two of adrenalin chlorid solution (1 to 1000) should be instilled at the time the cocain is used. The lids having been separated sufficiently by a stop-speculum, the patient should be asked to look as far as possible in the direct opposite direction from the muscle to be operated upon, so as to expose well the field of operation. With forceps the conjunctiva alone, over the insertion of the muscle, may be raised, when a small opening should be made with a Stevens scissors. Through this opening the capsule of Tenon may be grasped and then divided. Thus the tendon of the muscle is exposed, the central fibers of which should be lifted by the forceps and then divided by means of the Stevens scissors, thus buttonholing it. While still holding the tendon up with the forceps the buttonhole may be enlarged with the scissors by cutting first in one direction and then in the other, close to the sclera, being very careful not to encroach too far toward the peripheral fibers. The forceps should now be laid aside and the Stevens hook should be inserted through the buttonhole beneath the uncut fibers, first on the one side and then on the other, by means of which the amount of resistance is determined. If this is still too great other fibers must be cut with the scissors, between the hook and the sclera, always being careful to leave a sufficient number of peripheral fibers to act as stay-cords to prevent the cut muscle from retracting too much. The strength of the uncut fibers in both directions should be equal, so that the plane of the muscle may not be changed. Both judgment and skill must be exercised, else too much or too little of the tendon may be cut. It is better to aim at leaving some of the old error uncorrected than to transform it into the opposite condition. The conjunctiva may, but the capsule of Tenon must, be divided co-extensively with the division of the tendon, to obtain the effect desired. In no kind of heterophoria should a complete tenotomy ever be done; but, if by accident it should happen, the tendon should be stitched to the sclera directly behind the original insertion, and at that distance behind determined by a correct understanding of the exact character of the error for which the operation has been undertaken.

A skilled operator will find it easy to grasp, with the forceps, conjunctiva, capsule of Tenon and tendon of muscle all at the same time, and with one snip of the scissors buttonhole all these structures and then proceed with the operation as set forth above.

If a complicating cyclophoria is to be corrected by a partial tenotomy of a rectus, not only must the tension of the muscle be altered for the correction of the main error, but its plane must be changed so as to correct the complicating cyclophoria. The kind of cyclophoria having been determined—and it is nearly always a plus cyclophoria—one is not left in doubt as to how the operation should be done. To cure a hyperphoria and a plus cyclophoria the nasal and central fibers of the superior rectus of the hyperphoric eye should be divided, while the temporal fibers should be left uncut and sufficiently strong to prevent an over-correction of the hyperphoria; or the temporal and central fibers of the inferior rectus of the cataphoric eye should be

divided, leaving the nasal fibers uncut and of sufficient strength to prevent an over-correction of the error.

In a partial tenotomy for sthenic esophoria complicated with a plus cyclophoria, there being no hyperphoria, the lower and some of the central fibers of both interni should be divided, leaving the upper fibers uncut. In this way the tension of the muscle is altered, curing wholly, or in part, the esophoria, and the plane of each muscle is elevated so as to correct the plus cyclophoria. The plane of both interni having been equally elevated, there is developed, of necessity, a slight double hyperphoria, which, however, will give no trouble, being easily overcome by a pose of the head.

In operating on a case of sthenic esophoria complicated by a right hyperphoria and a plus cyclophoria, the first operation must be done on the internus of the cataphoric eye, and should consist of a complete division of the lower and central fibers, leaving uncut the upper fibers of the tendon. The threefold effect of this procedure is a correction in part, or wholly, of the esophoria, a correction of the cataphoria, and a cure of the plus cyclophoria. Whatever part of the esophoria may remain, after this operation, should be corrected by a central partial tenotomy of the right internus. Should some of the right hyperphoria remain, but no cyclophoria, a central partial tenotomy of the right superior rectus should be done; but if there should remain some uncorrected plus cyclophoria, as well as right hyperphoria, the inner and enough of the central fibers of the right superior rectus should be cut to cure these conditions. These three operations—sometimes even one or two of them—will cure a complicated case of this character.

If the esophoria is asthenic and uncomplicated, the operation of shortening should be done on one or both externi, and the plane of these muscles should be the same after, as before, the operation, otherwise a cyclophoria would be created.

If the asthenic esophoria is complicated with a right hyperphoria only, the externi should be shortened as though no complication existed, and later the hyperphoria could be treated by exercise, by a prism in position of rest for the hyperphoric eye, or by a central tenotomy of the superior rectus of the hyperphoric eye. In such a condition no muscle plane should ever be changed. The plane of a muscle should never be changed unless there is a cyclophoria to be corrected.

If the asthenic esophoria should be complicated with a plus cyclophoria alone, not only should both externi be shortened but the plane of each should be depressed so as to cure both esophoria and the plus cyclophoria. This double effect is accomplished by having the loop of the suture through the muscle near its upper margin, while the two ends are passed through the tendon near its lower margin. Tying the knot on the silver suture-plate pulls the muscle lower on the globe, thus changing its plane of action. The double cataphoria resulting will not be harmful.

If asthenic esophoria is complicated by a right hyperphoria and a plus cyclophoria, the first operation should be on the externus of the hyperphoric eye, and should be a shortening so done as, at the same time, to depress the plane of its action. The triple effect will be to cure more or less completely the esophoria, the right hyperphoria and the plus cyclophoria. If the left externus must be shortened for a remaining esophoria, whether complicated or not by a cataphoria and plus cyclophoria, one or both, for obvious reasons the plane

of this muscle should not be altered. If the plane were elevated, it would lessen the cataphoria, but would increase the plus cyclophoria; if the plane were lowered, it would decrease the plus cyclophoria, but would increase the cataphoria. After the straight-forward shortening of the externus of the cataphoric eye, whatever hyperphoria alone may exist should be treated, if sufficiently great in quantity, by a central partial tenotomy of the superior rectus of the hyperphoric eye; but if the remaining hyperphoria should be complicated with a remaining plus cyclophoria, the inner, and as much as necessary of the central, fibers of the superior rectus of the hyperphoric eye should be cut.

The operation for sthenic exophoria uncomplicated, is a central partial tenotomy of one or both externi, preferably both. The object in view being only the alteration of tension, care must be exercised that the plane of rotation shall not be changed. In operating on an externus it is far more easy to grasp, with fixation forceps, the conjunctiva, capsule of Tenon and muscle tendon, and buttonhole them with the scissors simultaneously, than it is with other muscles.

When sthenic exophoria is complicated by a hyperphoria the operation on the externi must not be done with a view of affecting the hyperphoria, hence a central partial tenotomy is indicated. Later the hyperphoria must be relieved by a central partial tenotomy of the superior rectus of the hyperphoric eye, and, if necessary, a central partial tenotomy of the inferior rectus of the cataphoric eye. The tension of these muscles should be altered without a change of plane.

In sthenic exophoria complicated by a right hyperphoria and a plus cyclophoria, the first operation should be done on the externus of the hyperphoric eye, and it should consist of a division of the upper and central fibers, leaving uncut the lower fibers. The threefold result of this operation will be: 1, relaxing the tension of the externus, lessening, if not curing, the exophoria; 2, a turning of the eye down, thus counteracting the hyperphoria; 3, torting the eye in, curing the cyclophoria. If some of the exophoria remain, whether still complicated or not by right hyperphoria and plus cyclophoria, the operation on the left externus must be a central partial tenotomy. The reason is clear: a division of the upper and central fibers would so change the muscle plane as to increase the cataphoria, although diminishing the cyclophoria; while a cutting of the lower and central fibers would so change the plane as to lessen the cataphoria but increase the plus cyclophoria. The only safe course between this Scylla and Charybdis is a central partial tenotomy of the left externus. These two operations having been done, any remaining right hyperphoria, without a plus cyclophoria, should be relieved by a central partial tenotomy of the right superior rectus; but if the remaining hyperphoria should be complicated with plus cyclophoria, the nasal and central fibers of this tendon should be cut, with the double purpose of altering the tension for the hyperphoria and changing the plane for the cyclophoria.

When sthenic exophoria is complicated by a plus cyclophoria only, the upper and central fibers of both externi (not one alone) should be cut. The triple effect of these operations is: 1, alteration of tension for the exophoria; 2, lowering the plane of both externi for the plus cyclophoria; 3, the development of a double cataphoria which, in itself, is not bad.

In asthenic exophoria uncomplicated, the tension of the externi must be increased by shortenings or advance-

ments so done as not to change the plane of rotation. The same is true of asthenic exophoria complicated by a hyperphoria alone. Later the hyperphoria may be relieved by a central partial tenotomy of the superior rectus of the hyperphoric eye.

When an asthenic exophoria is complicated with a right hyperphoria and a plus cyclophoria, the internus of the cataphoric eye should be so shortened or advanced as to alter its tension for the exophoria, and elevate the plane for counteracting the cataphoria and curing the plus cyclophoria. If the internus of the once hyperphoric eye must be shortened or advanced to still further correct the exophoria, not any longer complicated, it must be so done as not to change its plane; and the same is true if the only remaining complication is a hyperphoria, for in counteracting the hyperphoria, a plus cyclophoria would be developed. It is also true that only the tension of the internus should be altered by a shortening or advancement when the remaining exophoria is complicated by hyperphoria and plus cyclophoria, for the reason that lowering the plane would increase the cyclophoria although lessening the hyperphoria, while elevating the plane would increase the hyperphoria although diminishing the cyclophoria. Later the hyperphoria or hypercyclophoria should be remedied by the correct operation on the superior rectus.

An asthenic exophoria complicated by a plus cyclophoria must be treated by such shortening and advancement of both interni as to alter the tension for the exophoria and elevate both planes for the plus cyclophoria. The double hyperphoria resulting would be counteracted by a pose of the head.

Cyclophoria may exist alone and may be so high in degree as to demand relief by operation. This can be accomplished by operating on both superior or both inferior recti. A plus cyclophoria uncomplicated can be relieved by dividing a few of the nasal fibers or advancing a few of the temporal fibers of both superior recti. In doing the former a double cataphoria is developed while the cyclophoria is cured; in doing the latter the cyclophoria is cured but a double hyperphoria results. Since a double cataphoria is preferable to a double hyperphoria, a division of the nasal fibers of the superior recti should always be chosen.

The plus cyclophoria can be cured by a division of the temporal fibers or an advancement of the nasal fibers of both inferior recti. The former would give a double hyperphoria while the latter would give a double cataphoria, hence, of the two, the latter should be chosen.

As to final results a division of the nasal fibers of the superior recti and an advancement of the nasal fibers of the inferior recti are precisely alike; but the former should be preferred, for it is more easily done and gives the patient much less inconvenience.

DISCUSSION ON PAPERS OF DRs. GOULD AND SAVAGE.

DR. S. D. RISLEY, Philadelphia—While I regret very much that Dr. Gould finds it necessary, under the qualifying conditions stated in his paper, to regard tenotomy of certain ocular muscles as malpractice, I find some consolation in the fact that Dr. Savage did not insist that it was malpractice to omit tenotomy. I am willing to admit that after the series of papers by Dr. Stevens, published several years ago, that the ophthalmological world ran tenotomy-mad. I heard one surgeon say that he had done more than 200 tenotomies in a single year. In one of these papers I believe we have an illustration of how the pendulum may swing too far in the opposite direction.

In 1894 in a paper I had the honor to present at Edinburgh to the International Congress, I pointed out some of the diffi-

culties that lie in the way. Attention was called to the frequent distortions to the human skull as shown in any cursory study of a hatter's forms taken as patterns from the heads of his patrons; many of which are so ill-shaped as to appear grotesque when compared with the diameters of a model skull. It is highly probable that these anomalous forms of the skull modify the form and dimensions of the bony orbit by changing the line of direction of the orbital planes. It is reasonable to suppose that during development in an anomalously shaped orbit that not only certain modifications would occur in the form of the eyeballs and in the length and line of direction of the optic nerves, but also in the origin and point of attachment, the length and direction of the ocular muscles; conditions, any one of which would lead to more or less marked disturbance of motility, and in many complex ways modify the conditions of binocular vision. In view of such considerations it is plainly obvious that any effort to reduce all the possible abnormalities of binocular vision to the simplicity of a law governing their occurrence and correction will probably result in failure. I believe, therefore, that we can not afford either in the study or management of these conditions to be dogmatic. It is my opinion that no chapter in ophthalmology is filled with more difficulties; that no subject should be approached with more critical caution than the abnormalities of binocular vision. That much harm has been done by unwisely performed tenotomies I am ready to believe; but, on the other hand, I am also sure that much suffering has been relieved and many careers rendered more hopeful by the wise aid of the skilful and cautious surgeon.

DR. M. F. WEYMANN, St. Joseph, Mo.—When a physician practices very largely in any one line, he acquires a certain amount of perfection not obtained by the average man. Thus, Dr. Savage may be able to get results above the understanding and reach of the rest. However, a measure not assimilable by the professional corps at large is a dangerous practice to advocate for general purposes, on account of the failures and actual damage in some instances that may result. Surgical interference with the muscles of the vertical plane seems to me a bad practice, although in tenotomies of the muscles of the horizontal plane I have had many nice results. Yet, even here, the complex nature of the muscular mechanism upsets often all seemingly correct and logical reasoning. A few weeks ago I operated for internal strabismus on a little girl. I decided to cut the upper fibers of the internal rectus tendon, but through some unaccountable mistake I clipped the lower portion. I was surprised to see the un-intended section produce the intended result. Likewise, I have failed when faithfully carrying out partial section according to established notions. Dr. Risley's reference to the great variability of orbital conditions also tends to show the impracticability of hard and fast rules.

DR. DUDLEY S. REYNOLDS, Louisville—There are a great many other questions to settle before proceeding to partial, or complete tenotomy in such cases. Conditions of malnutrition, state of the nervous system, habits of the individual, mode of living, age and hereditary conditions, all have to be considered. It is very important to remember what Dr. Risley has just said about physical conformation of the face and head in connection with abnormal shape of the orbit, and consequent anomalies of the orbital contents. Surely no one would expect to change the shape of the orbit by dividing the tendons of ocular muscles. Many a case of heterophoria due to anxiety and overwork, with consequent malnutrition and loss of rest has been ruined by frequent tenotomies. One recent case in my observation was operated 13 times, and the surgeon wanted to operate again, in a person with a small central scotoma in one eye. A multitude of cases of heterophoria may be cured by constitutional treatment with suitable regulation of diet, habits and mode of living. Many are cured by constitutional medication and correction of errors of refraction, without either prisms or tenotomies. Reflexes from the genito-urinary system are sometimes causes of heterophoria. I have known more than one case cured by successful operation for the cure of hemorrhoids. Many of them are due to auto-intoxication from con-

stipation of the bowels. All these, and many more matters should be duly considered before subjecting the patient to any sort of tenotomy.

DR. J. E. COLBURN, Chicago—I must plead guilty to having performed a good many of these operations, and while I have had many occasions to regret operations I have had a great many occasions to congratulate myself and the patients upon the good results obtained. I believe with Dr. Risley that the conformation of the orbit has very much to do with the position of the eyes, their relative activity and with the production of heterophoria. I believe that errors of refraction are the cause of a great many cases of pseudo-heterophoria and also that general debility is the cause of some instances; such patients can not bear the discomforts of a muscle error with the same degree of patience they have been able to do when in a better systemic condition. I believe it is possible in a certain class of cases to accomplish many of the things Dr. Gould has referred to in his paper, but I feel that his rules must be exceedingly limited in their application. That a tenotomy, graduated or partial, is a serious operation, I am well aware, but I believe it is a justifiable procedure when done under the proper conditions. I believe in giving the patient the benefit of correcting the refraction, gymnastic exercises, etc., but that there are a certain class of cases where we must consider the patient and his inability to give us the months of time for exercises necessary to bring about the conditions which Dr. Gould says will result and that we must give such patients the relief if only for a time that comes by operation.

DR. C. H. WILLIAMS, Boston—Before we do any operations on the eye muscles, or in fact before we make a diagnosis in these complicated cases, it seems to me we need to get at all the facts of the case, as to how much muscle error there is and where it is located. I wish to call your attention for a moment to a piece of apparatus which I made last summer and which I have used for the past six months and found very helpful in this direction [exhibiting new instrument]. It consists of a hollow tin case shaped like a Geneva cross, each arm being about eighteen inches long; inside this case are a series of incandescent lights. A slot is cut in the front of each arm about half an inch wide and the length of the arm, behind which is fastened a strip of red glass. In front of this case is a wheel of wire pivoted at the intersection of the arms of the cross; this wheel carries sixteen small incandescent lights arranged in the form of a cross, to correspond to the arms of the case below. These lights are covered by having a small cylinder of tin about each lamp, and on the front of each cylinder is a number in green glass; these lights are placed on the wheel at such distances apart as would correspond to the deviation produced by a prism, when viewing the lights at a distance of twenty feet, and the number before each light gives the number of the prism which would at that distance cause a displacement of that amount. The person to be tested wears a pair of spectacles, of which one eye is filled with a red glass, and the other eye with a green glass. When testing a patient, the electric lights of the instrument are controlled by a switch so that the vertical red line and the horizontal numbers are lighted at one time, or the horizontal red line with the vertical numbers, or the vertical red line and the vertical numbers. Starting with the first combination, the vertical red line and the horizontal green numbers, the patient is asked to state where the red line seems to stand; if it comes in its true position between the 1 and 2 we know that there is no deviation of the axes of the eyes to the right or left, but if the red line seems to stand to the right (the red glass in the spectacles being before the right eye) it shows there is a convergence of the axes, and if the red line is over No. 4 it shows this deviation is equal to that produced by a prism of four degrees. If on the other hand the red line appears over No. 5 to the left, it shows there is a divergence of the axes equal to a prism of five degrees. The even numbers are to the right of the center, and the odd numbers to the left. Again, when testing a patient with the horizontal red line and the vertical green numbers showing, if he sees the red line in the center between the 1 and 2 it shows he has no

displacement of the axes up or down, but if the red line stands near the number 5 below the center it shows that the axis of that eye stands above the other equal to the displacement caused by a prism of five degrees. To show the amount of torsion, if any, the vertical red line and the vertical green numbers are lighted. If they appear superimposed, or standing parallel to each other, there is no torsion, but if the red lines and the green figures appear crossed the vertical meridians of the two eyes are not parallel, and the number of degrees which it is necessary to turn the wheel with the green figures in order to bring the red line and the green figures parallel, measures the amount of torsion. Through these spectacles the patient can only see the red lines with his right or red eye, and can only see the green figures with the left or green eye, the glasses cutting off the other rays. In this way each eye is unconsciously fixing a separate object, and if the axes of the eyes are in their proper position the red lines and green figures will seem to be in their proper place, but if there is a deviation its character and amount can be quickly and accurately measured. It is desirable to have a second pair of spectacles with the red glass in the left eye, and the green glass in the right eye; and it is essential to select the color of the spectacle glasses with care, so that the red glass will not transmit any of the green light, nor the green glass any of the red light from the lamps.

DR. J. L. THOMPSON, Indianapolis—In the beginning of my thirty years' practice I paid more attention to insufficiencies of the internal muscle than to esophoria, but since I have been examining for esophoria I find that 98 out of 100 of my patients have from 1 to 3 degrees and that it seems as natural as the flexion of the hands and arms. In fact, if the patient does not have esophoria I look upon the case as an insufficiency of the internal recti muscles. I started in my practice with a great fear of operations upon the internal muscles except in cases of strabismus, where I operated freely, and it astonishes me now after twenty years to see that I have not had much over-correction in these cases. I have never operated on a case of strabismus until it has been under observation for a year, and I think in the future it will be five years before I operate on these muscle cases.

DR. G. C. SAVAGE, Nashville, Tenn.—Dr. Risley mentioned malformations of the skull as causative of heterophoria. There is nothing clearer than that a malformed orbit must contain an eye out of its normal relationship. The result will be one kind or another of heterophoria. In the treatment of such heterophoria, however, it is impossible to alter the shape of the orbit; therefore, if the error is to be corrected, it must be done by treatment of the muscles involved. Whatever then may be the cause of a heterophoria which is intrinsic in character, the principle of treatment remains the same. I wish to say that the instrument that Dr. Williams has very ingeniously invented will do everything that he claims for it. I do not want anybody to go from this meeting believing that I am an extremist on the muscle question, for I am not. My paper was of necessity confined to the operative treatment of heterophoria, and I am always delighted when I have a case of intrinsic heterophoria which can be cured by exercise, or by weak prisms in positions of rest. There are, however, a great many cases that can not be thus cured, and on these I always operate.

DR. FRANK ALLPORT, Chicago—In considering the question of graduated or other tenotomies for the relief of so-called muscular asthenopia, I like not only to be honest with my patients and refrain from burdening them with unnecessary expenses, but I like also to be honest with myself. I must, therefore, ask myself the question: Do I find it necessary to perform tenotomies for asthenopia in order to secure satisfactory practical results, both to my patients and myself? A negative reply is inevitably the result of such an interrogatory. I have the average number of asthenopic patients in private and dispensary practice, and I presume I secure about the average results. At all events, in the course of a year I do not see many asthenopic patients who express much dissatisfaction from their treatment. I do not claim better re-

sults than others, and I can honestly say that I very rarely perform a tenotomy for asthenopia. Sometimes a year or so may elapse without the performance of such an operation. I therefore find it difficult to understand why some men perform this operation so frequently. I find that most of my cases do very well without tenotomies and without prisms, by a thorough correction of refractive errors and observance of the laws of health and hygiene. It may be with the unusual knowledge and skill possessed by such men as Drs. Stevens and Savage, the frequent performance of such operations may be not only useful but wise, but I am free to confess that from my own standpoint I do not feel myself capable of freely interfering with the functions of the ocular muscles. I recall the case of a gentleman operated on in Chicago who had only one eye. Through an accident his other eye was completely blinded. Some years after this accident, while suffering from asthenopia, he sought the advice of a professional muscle clipper, who, through motives which I do not attempt to analyze, performed several partial tenotomies on the good eye.

DR. C. F. CLARK, Columbus, Ohio—My experience has been very like that of the last speaker; yet I have been inclined to think of late that I have been too conservative in the matter of operations. I am willing to run the risk with being charged with radicalism when I say that within a few months a case that I had treated with muscle exercises faithfully, and for a time successfully, came to me and I found that I had kept that patient for several years with only a partial use of her eyes when she had the need for the full use of them; a very slight advancement gave me a brilliant result. By muscle exercise following the operation the abduction was carried from 18 degrees to 58 within three weeks and she had no discomfort. It convinced me that in many cases I had been guilty, as perhaps Dr. Gould has been, of being too conservative.

DR. HIRAM WOODS, Baltimore—I can fully agree with Dr. Risley that if you once admit Dr. Savage's premise and then work out his conclusions you are compelled to admit that his reasoning is logical. The question with me is the premise, as given by Dr. Savage: "Heterophoria not curable by correction of refraction errors, prisms in position, or strengthening exercise." I think he should add, "provided the heterophoria is shown to be the cause of trouble still persisting." Experience has taught me that with more exact and careful methods of diagnosis the number of cases that are left for tenotomy are very small. I formerly did from ten to twenty a year, while now I do not believe I do more than one or two, and then usually for esophoria. One of the most important steps in the examination is to try to determine what the heterophoria is due to and in this connection we must not only look after the refraction and the health but into the school and home hygienic conditions. It takes a considerable amount of time to get down to these facts and possibly Dr. Thompson is right in putting it at five years. I can not go so far as Dr. Gould in saying that operation is never justifiable, for there is a class of cases in which enormous benefit results, namely, in esophoria. I do not find the high degrees of hyperphoria that some do. I have never used the instrument that Dr. Savage exhibits, but I have looked for cyclophoria persistently with the Maddox prism, as Dr. Savage taught, and found it only three or four times. In esophoria, however, with low abduction, and with the parallax displacement which Duane described, a tenotomy is often followed by the best results. I know that in my own experience a tenotomy done by Dr. Harlan of Baltimore benefited me enormously. I want to ask Dr. Gould in regard to his method of muscle exercise whether he considers the rapid development of adduction an increase in muscular strength or a development of positive relative convergence. Landolt dwells upon defective relative convergence as a cause of asthenopia. Why is it claimed that orthophoria is as bad a condition as some degrees of heterophoria? While the normal condition probably is a degree or two of esophoria for distance and three or four degrees of exophoria at the near point, as Dr. Theobald pointed out some years ago, the statement that orthophoria causes asthenopia is not in accord with my observations.

DR. L. CONNOR, Detroit—It is perfectly evident that this is a very complex problem and that some gentlemen see one part of it with greater distinctness than others. So far as my own observation goes we may seek for the cause of these muscular disturbances along perhaps three or four lines: one is a reflex occurring with certain conditions of the nose or allied organs; another is malnutrition of the eye whose muscles wobble as the muscles elsewhere do in similar conditions; a third is a physiological lack of power, and lastly there are unquestionably anatomical defects. I think it is well established that when we have settled upon an anatomical error as the cause of the trouble an operation is absolutely necessary. In other conditions the removal of the cause of irritation, etc., has certainly resulted well. It is entirely unsafe to say that anatomical conditions alone are the cause of heterophoria and that all cases should be operated upon. A friend of mine was operated on thirty-eight times within eight months and has more muscular unbalance to-day than ever before. You all know there are cases of disease of various sorts cured by operation *per se* and I am certain that many cases of heterophoria are cured simply by the cutting, and a cutting on the ear would do just as much good. I have never seen any good resulting from a partial tenotomy; I might just as well have cut the conjunctiva somewhere else. I do not mean to say the men who operate do not benefit their patients, but the good should be ascribed to its proper cause and, in some instances at least, to hypnotic influence.

DR. D. B. WYLIE, Milwaukee—The gentleman who has just had the floor touched upon a point which I think is not looked after to the extent it should be, which, in my practice at least, has appeared to be of great importance, that is, external irritation. In neither of the papers, nor in the discussion have I heard anything about correcting diseased conditions or irritated conditions of the conjunctiva. You may call these pseudo-heterophorias if you will, but they produce the same asthenopia. I have time and again seen small degrees of esophoria or exophoria disappear entirely after the cure of a very mild degree of conjunctivitis. I was somewhat surprised at one of the gentlemen stating that surgery is the despair of medicine. It seems to me that is beyond the power of any man to defend.

DR. D. M. CAMPBELL, Detroit, Mich.—The differences of opinion seem to be on the causes of heterophoria; in the treatment of the condition we are probably not so far asunder as might appear from the discussion. Most of us correct the refraction and look after the general health of the patient and cure a large number of cases, and yet in a large practice there will be a certain residual number uncured by whatever we do. In a report I made to the Michigan State Medical Society upon a study of 500 cases of asthenopia I found less than 2 per cent. that had to undergo operation.

DR. H. HARLAN, Baltimore—I envy very much the men who can have such positive convictions as to permit dogmatic statements such as we have heard. I want to ask Dr. Gould whether his statement that no case should be operated on is the swinging backward of the pendulum as the result of unsatisfactory experience in operating, or whether that has always been his opinion.

DR. GEORGE M. GOULD, Philadelphia—It is my belief that we are emphasizing entirely too much this question of muscle trouble. It may be highly scientific to do this work, but I have found many patients coming to me wearing prisms that seemed to my mind utterly unnecessary and never could have been called for. I feel it a higher duty to emphasize the importance to correct refraction, of applying the proper glasses in relation to these muscle troubles. Several misunderstandings have taken place which I wish to correct. I never said or implied that no case should be operated on. I made certain reservations and further said that this paper was written from my own experience limited to a private practice in a good class of intelligent people; I have been able to cure their symptoms or relieve them so perfectly that I have kept them as satisfied patients. I do not wish to be dogmatic because I thoroughly believe that in the matter of the practice of medi-

cine individual genius has its own law and one man will accomplish by his method what another man could not accomplish with it.

I wish to correct a misinterpretation about "surgery being the despair of medicine." What I meant was that none of us believe in proceeding to surgical measures if we can cure our patients by medical and physiological means. We certainly despair of curing our patients naturally before we proceed to surgery. I do believe that local irritation of the conjunctiva may produce a temporary heterophoria, but on the other hand I believe these conditions are usually corrected by proper glasses which remove the heterophoria at the same time. The question has been asked why orthophoria should be considered a disease. My feeling is that in orthophoria with the Maddox rod at 20 feet there is almost always an insufficient adduction power for close work and that the trouble thus produced justifies its being considered a diseased condition.

A TABLE OF OCULAR EXTRINSIC PARALYSES.*

HORACE M. STARKEY, M.D.
CHICAGO.

In the "International Clinics," Vol. iii, Fourth Series, 1894, appeared an article by Prof. F. C. Hotz, on "The Diagnosis of Paralysis of the Ocular Muscles by the Double Image Test," which was a distinct advance in simplifying the methods of detecting the affected muscle in these perplexing cases. In studying that article, it occurred to the writer that the subject might be further simplified by exhibiting these paralyzes with the position of the false image in graphic or tabular form. Soon afterward he prepared the table, which, with some modification of terms, has been used by him in teaching and in practical work since. As the table has stood the test of time and has commended itself to many practitioners and students, it is now presented for the consideration of this Section, in the hope that it may prove of value to others.

Let us consider briefly the more recent nomenclature of the movements of the eyes, mainly following that of Maddox; the muscles producing those movements, and the action of each muscle in the primary position of the eyes as well as in secondary positions.

PRIMARY POSITION.

The primary position may be stated to be that in which with the head perfectly erect, the eyes are directed toward a point upon the horizon midway between the eyes.¹

In testing the eye muscles the primary position should be first assumed and the relations of the images formed by the two eyes studied in this position and later the relation of the images in departures from this position should be considered.

MOVEMENTS OF THE EYES.

In passing from the primary position, the eyes are capable of eight simple co-ordinate movements and, by combinations of these, all possible positions may be attained. Two of these movements, adduction and abduction, are employed in convergence and divergence; four, dextroduction, levoduction, superduction, and subduction, are employed in parallel movements; and two, extortion and intortion, are employed in rotating the eyes around their antero-posterior axis "so as to make the cornea revolve like a wheel." Because of this

* Read at the Fifty-second Annual Meeting of the American Medical Association, in the Section on Ophthalmology, and approved for publication by the Executive Committee of the Section: Drs. J. A. Lippincott, Casey A. Wood and H. V. Würdemann.

1. While this definition is not strictly accurate for all movements of the eyes, it is sufficiently accurate for all practical purposes in making the diplopia tests.