edema and albuminous urine. At the time of the interview he had a normal pulse and respiration, no cyanosis, edema or albumin, and nearly every day went to walk with his mother, sometimes climbing a hill of three-quarters of a mile of considerable grade in the course of the walk.

OBSERVATIONS ON MALFORMATIONS OF THE RECTUM AND IMPERFORATE ANUS.*

SAMUEL W. KELLEY, M.D.

Professor of Diseases of Children, Cleveland College of Physicians and Surgeons; Surgeon-in-Chief Holy Cross Home for Crippled Children; Pediatricist to the City and to Cleveland General Hospital; Pediatricist and Orthopedist to St. Clair Hospital.
CLEVELAND, OHIO.

I shall not present a monograph in twenty minutes; but shall pass by history and classifications and also the simple cases in which a membranous septum closes the anus and which are comparatively easily managed. Your attention will be directed for a moment, however, to one rather simple case, so far as treatment was concerned, in which the configuration was peculiar (Fig. 1).

This case was referred to me by Dr. I. W. Bard. A depression marked the proper site for the anus, as is not uncommon (though in imperforate anus, even with a raphe, the perineum is usually smooth); but across this depression extended forward a tapering tail-like process from the coccygeal region which continued into the raphe of the scrotum, dividing the little cuplike cavity of the proctodeum into two lateral halves.

This is a fairly marked example of a comparatively rare malformation. A few others of its kind have been reported in which the anal depression reached the rectum, but left a central bridge across the opening. They seem to contradict the usually accepted view that the ectodermal layer simply dimples in until it meets the terminus of the enteron. It rather confirms the description of Tourneux, that there is a definite anal membrane or specialized cellular mass thicker in the middle line than at its sides, forming the anterior wall of an internal cloaca, or rectal ampulla; and that this cellular body is absorbed away, the greatest absorption taking place at each side of the central line, where the thicker portion remains last to be absorbed and, in such malformation as this, fails to be absorbed.

In this case there was restlessness, distress and moderate straining and refusal to take the breast, but no vomiting. Patient was operated on at St. John's Hospital. Dissection in the median line extended upward a distance of about three-fourths of an inch; encountered and opened the rectal ampulla, and discharged the meconium, to the great relief of the infant, which seized the mother's breast at the first opportunity. There was no appearance of any fibrous cord extending from the anal site to the rectal pouch, as is often described in the books; and in a number of cases that have come under my observation I have never seen such a cord and think it must be unusual.

I was particularly impressed in this case with the advantage of operating without an anesthetic. In the absence of expulsive efforts the exploring finger gains no sense of proximity of the distended pouch. But during straining the increased tension is at once communicated, even through a considerable thickness of tissue. Pressure of the hand over the hypogastrium will simulate straining, but, the diaphragm not being fixed, it is not so effectual. This case, after one simple incision and

evacuation, followed by use of bougies daily for ten days, resulted in a functional cure as complete as one could wish, with perfect control. Thus my fears, taught by previous experience and reading, that simple incision of a septum of that thickness would be followed by cicatricial contraction, failed to be realized. Usually if the impervious portion necessarily cut through be more than a membranous septum, or if it be only a septum but situated at some distance from the surface, it is necessary to bring down the pouch and stitch it to the freshened edges of the skin at the anal margin, thus securing a mucous lining to the canal, to prevent recontraction.

Since writing the above I have again visited this patient, eight months after the operation, and found a fine, well-grown baby said to have normal, healthy stools. On examination, I found the anus rather small, admitting only the end of my fourth finger, and rather rigid. In case of constipation there might be trouble. But what especially struck me was the fact that the rectal cavity was encountered at a depth of a half inch or less, as though a process of development since birth had brought it nearer to the surface. There are one or two cases on record in which, after inguinal colostomy, the blind end of the gut went on developing downward so that later it could be easily reached by the perineal route.

But I wish now to relate two cases of severe malformation of the rectum.

The first case was seen March 24, 1903, in consultation with Dr. Etienne P. Crowe, accoucheur. Male babe of J. C., the father a Scotch-American and the mother a Bohemian. They had one older child, well formed and healthy, and denied all family history of deformities, and any deleterious influence during pregnancy. When seen by me the babe was forty-eight hours old and was said to have urinated, but had no bowel movement, and Dr. Crowe had discovered that no anus existed. In its place was a slight groove. The infant appeared of average size, fairly plump, and well-formed, excepting that there was a median cleft partially separating the scrotum into two lateral halves, with a testicle in each. At the bottom of this cleft, half an inch posterior to the penoscrotal junction, were two small orifices. One of these extended only skin deep, the other deeper. It was said that a half dram of blackish fluid had been found on the diaper and that the scrotal cleft had been stained with the same material. With a small probe and with a filiform bougie a persistent effort was made to trace the course of the deeper sinus, as I thought this the key to the situation, but could penetrate little more than an inch without danger of making a false passage. While inspecting the parts a small amount of gas escaped from one of the orifices. The sinus ran in the same direction as the urethra, but posteriorly to it. The urethra was patulous and the urine was not stained with meconium. Therefore, an opening between intestine and bladder was excluded. A probe passed into the bladder revealed no greater encroachment of abdominal viscera in any one direction. Dr. Crowe and Dr. Propper had made an attempt to find the rectum by making an incision through the perineum in the median line three-fourths of an inch long and one and one-half inches into the pelvis without finding the gut. A finger introduced into this incision could feel urethra, prostate, and bladder, but no gut, and no tension during straining. The distension of the abdomen was pitiful. It appeared ready to burst; while the entire surface of the infant was purple from interference with the venous return. I noted that this stasis seemed a little greater in the right lower extremity; but the abdomen was equally distended on both sides. There was no fecal vomiting. The pulse could hardly be counted.

The condition appeared so urgent that I decided to abandon the perineal route and open the abdomen. I operated at the house. A few breaths of chloroform were given; made an inch incision of a particularly prominent portion of the left inguinal region. A few drams of amber-colored fluid escaped.

* Read in the Section on Diseases of Children of the American Medical Association, at the Fifty-eighth Annual Session, held at Atlantic City, June, 1907.

No ampulla and no colon could be reached; but a greatly distended loop of small intestine made its appearance in the incision and was stitched into its margins, packed round with gauze and opened. Quantities of flatus and some half-digested flakes of casein escaped from the opening. The distension on the left side of the abdomen was relieved, but not that on the right. We considered the advisability of opening the right side, but the parents refused point blank to allow anything more to be done. The infant stood the operation nicely and appeared relieved. The pulse counted 160. Ordered a few drops of brandy with water every hour.

The babe passed a very uncomfortable night. Next morning the parents refused to allow even an examination. With great effort we obtained the privilege of looking at it without removing it from the bed at the mother's side. It was grunting with pain and the pressure against the diaphragm. The artificial anus had discharged some more of the bowel contents, but not a great deal. The abdomen was greatly distended, being painfully tense on the right side, especially in the cecal region, and the skin on the abdomen and right lower extremity was turgid. Dr. Crowe and I expended our combined logic and eloquence on the parents. They were absolutely obdurate against further interference. The babe died that evening, thirty-two hours after operation.

By the greatest effort imaginable, Dr. Crowe secured for us the privilege of an autopsy next morning. The weight was seven pounds and four ounces, length twenty-one and a half inches, length of upper extremities eight inches, and of lower extremities seven and three-fourths inches. From umbilicus to heel ten and one-fourth inches; circumference of head thirteen and one-fourth inches, of chest thirteen inches, of belly sixteen and one-fourth inches, showing the babe to be of average size and general development. Distension general over abdomen, with veins of skin full. Distension more tense on right side. On opening the abdomen the perineum was found congested like the skin, and its cavity contained more fluid than normal, of an amber color. Bladder extended as high as two-thirds of the distance from os pubis to umbilicus, but was empty. Intestines distended with gas, as well as meconium. The cecum, with the appendix behind it, was located half way from the umbilicus to the cartilage of the ribs, right side (Fig. 3). Thence the colon extended transversely to the left, made the splenic flexure, then turned to the right and crossed the abdomen at the level of the umbilicus to a large pouch which occupied the usual position of the cecum. It had a gangrenous patch seven-eighths of an inch in diameter, which burst open as soon as touched. This large pouch, which terminated the colon and occupied the cecal region, extended beyond the median line and was closely attached to the bladder. It was rounded in outline, but came to a funnel-shaped tapering end in proximity to the neck of the bladder, at the right and posteriorly to the urethra. The funnel point extended toward the canal opening in the scrotum, with which I believe it was continuous, although patient efforts failed to pass the smallest canaliculus probe from one opening to the other. There was about half an inch impenetrable. The inferior surface of the right lobe of the liver presented forward, the anterior border being turned upward behind the ribs. The stomach was normal in shape excepting that its pyloric end, like the border of the liver, was turned upward by the pressure of the intestines.

The second history I wish to relate is that of Baby S. S., male, of Jewish parents, their fifth child, all the others being normal. On the second day after birth the midwife, who was also the nurse, made the discovery that the babe had no anus. A neighborhood doctor was called. He made a small scratch with scissors at the site of the anus and advised the people to wait till next day and take the babe to a dispensary. But I was called on the third day. The babe had not at any time taken the breast nor any drink nor passed any stool. It was said to have urinated and that the urine was clear. No vomiting. He appeared rather under the average size and weight. His abdomen was greatly distended and tympanitic. He was feeble and apathetic. Straining was very moderate and at long intervals and only excited by manipulation of the abdo-

men. The skin was cyanotic from pressure within the abdomen, and was also jaundiced, the combination giving a very peculiar color. The epidermis was exfoliating heavily. There was no dimple or depression at the anal site. A raphe was indicated by the structure of the skin, but was not raised above the smooth surface. There appeared to be no deformity of the pelvis or close approximation of the ischial tuberosities, as is said sometimes to occur. The parents were of the opinion that God never intended that the child should live or He wouldn't have made it that way, and they were very anxious He should take it off their hands. The midwife declared no such child ought to live and she knew this one would not. I mention this to indicate the kind of care it was getting and was likely to get.

I persuaded them to take it to St. Clair Hospital, where I operated on it at once, without anesthetic, dissecting in from perineum. Having passed through the floor of the pelvis, I used a blunt dissector and then my fourth finger between bladder and sacrum, a distance of two inches, but felt no gut nor any bulging on straining. The babe seemed so debilitated that extensive dissection by the pelvic route seemed inadvisable, and I decided to at least try to give relief by opening the abdomen. I drew the attention of my class to the fact that the distension of the abdomen was larger on the right side than on the left, and remembering the previous case, I felt considerable hesitation in choosing which side to open. Under a very little chloroform, for fear of adding to the depression of the stercoræmia and the jaundice by anesthesia, I incised the skin and fascia, two or three finger breadths (of the infant) above the middle of Poupart's ligament on the left side, and separated the muscle layers. Going through the abdominal wall, I found my opening at once into the gut, there being no peritoneal cavity between the parietes and the bowel. This seemed very odd. Using finger and probe, I decided that the cavity entered was the rectal pouch near its lower end, blended with the abdominal wall. Great quantities of gas escaped, and also a dark fluid, not as dark or thick as meconium usually is. I sutured the layers together around the opening, which made a very good artificial anus. Dressed with abundant gauze. The cyanotic hue disappeared from the skin. Soon the effect of the chloroform passed and the babe at once took brandy and water and later an ounce of water with a dram of coffee. During operation hot water bags were used and all possible protection from cold and shock. The parents insisted on taking him home at once, on a wintry day, and the nurse, who staked her reputation on a prediction that he would not live, carried him off. He died in fifteen hours. No post-mortem allowed.

Reflecting on these two cases, I think they show, for one thing, the necessity of early discovery of the presence of malformation and prompt surgical interference; for there is no possible chance of life otherwise, and waiting only produces more distension and exhaustion, allows time for the meconium to become septic, stercoræmia to ensue, and for the development of jaundice, peritonitis, gangrene, rupture of the colon or other complication. An examination of the infant should be made at its birth as to the normal development and the integrity of all its parts.

It seems strange that fecal vomiting is not a more common symptom in these cases. It is said to be unusual. The rare cases which survived more than seven or eight days, sometimes several weeks, have had periodical fecal vomiting, but they are so rare as not to count in the prognosis.

These cases illustrate, too, the impossibility of ascertaining in cases of imperforate anus, before operation, the degree of development of the rectum or the location of the enteron. In both cases the abdomen was distended and the skin of the body and lower extremities was turgid. In the first case the terminal pouch was located on the right side, and in the second case, although the right side of the abdomen was more dis-

tended than the left, the pouch was located on the left. However, in the first case the venous stasis was somewhat greater in the right extremity than in the left, but as both were much congested and it was only a matter of degree, I did not take it as a guide. Perhaps it was an index I should have valued more. Greater venous stasis, regardless of greater abdominal distension, may be as valuable a clue to the location of the pouch as it would be of any tumor. But the tympanitis and distension being the same on both sides, I considered the probability of the colon's ending on the left side—and on going in there, missed it. In neither case did examination by the sound in the bladder prove of any value in determining in which direction the distended pouch was situated.

Certainly in the first case, if I could have succeeded in passing a probe through the sinus, I could have located the pouch. That being impossible, it is debatable

Fig. 1.—Imperforate anus with a tapering tail-like ridge extending forward from the region of the coccyx, dividing the cup-like cavity of the proctodeum into two lateral halves.

Fig. 2.—Malformation of rectum. Anus absent. A groove partly divides the scrotum in the median line; in the groove near the penoscrotal junction is the orifice of a sinus. Note the distension of the abdomen and discoloration of the skin.

ground whether, if the infant had been in good condition to endure extensive dissection, it might not have been well to follow the sinus as a guide and dissect in until the pouch was reached, or to pursue the perineal coccygeal or sacral route. The situation of the orifice of the sinus was not suitable for an anus, yet if once the pouch were located and a temporary canal for evacuation established, an attempt could have been made afterward to establish a permanent artificial anus in a more suitable location.

But I think if a case of imperforate anus or malformed rectum, without a pervious sinus, were to come to me to-day in a fit condition for operation, I would persist in the perineal, coccygeal, even a sacral route,

separate and drag down the pouch if possible and stitch it to the skin margin. Yet in the first case even that procedure, I fear, would have proved impossible of execution on account of the distance at which the pouch was located.

In a case like the second, it probably would have succeeded if performed early with the patient in good condition at the start. But after waiting till the third day, in the presence of exhaustion and jaundice, such an operation would have proved too severe and would have precipitated the fatal result, as was shown by the sequel; for the slighter operation of making an artificial anus, which took but a few minutes' time and with no hemorrhage, and which evacuated the canal and thus gave relief, was followed by death fifteen hours later.

How much the result would have been altered if the patient had had better surroundings and better nursing is hard to estimate.

It is stated by some writers that ano-rectal imperfora-

Fig. 3.—Portion of lower bowel showing pouch-like distension. The specimen is greatly shrunken by the preservative. The enormous pouch occupied the right hypogastrium and terminated in a minute sinus to the right of and posterior to the urethra.

tion is apt to be accompanied by other deformity, especially by contracted pelvis, extroversion of the bladder, spina bifida or nasopharyngeal obstruction; but in none of the cases that I have seen was such malformation present.

In one case of a very puny child with its anus closed by a membranous septum the infant's forehead was wedge-shaped, the frontal bones meeting at an acute angle. The occluding membrane had been opened by Dr. F. J. Morton and the rectum seemed patulous for several inches each time that I examined it.

But the babe persistently vomited whatever was given it to swallow, though it never vomited meconium, and never passed any noticeable amount of meconium from the bowels. It died, apparently, as a result of exhaustion, on the sixth day after birth. Probably there may have existed other occlusion somewhere in the intestinal tract. No operation and no postmortem were allowed.

2255 Fifty-fifth Street.

DISCUSSION.

DR. L. T. ROYSTER, Norfolk, Va., saw in consultation about eighteen months ago a child with absence of anus—no raphe even was present. Operation was performed under chloroform. One and one-half inches from the skin was found the end of the intestine, which was brought down. The child made an uneventful recovery with well-formed stools. It died a year later of some acute disease.

DR. E. H. SMALL, Pittsburg, Pa., reported a case of imperforate rectum which he saw about four years ago. A baby boy was born under his care, well developed, but with no anus whatever. When twenty-four hours old his condition was good, and with the aid of another doctor a trochar $\frac{1}{4}$ inch in diameter and $2\frac{1}{2}$ inches in length, was thrust into the perineum where the anus should have been toward the hollow of the sacrum. When it had entered about two inches the rectum was entered, the stylet withdrawn and the meconium gushed out. Then the trochar was withdrawn and the wound enlarged enough to admit the little finger into the rectum. The gut was so adherent to the surrounding tissues that it was impossible to bring it down to be united to the skin. A catheter was introduced and dilatation was kept up with the finger and then with sounds. The rectum gradually approached the outside and a good sphincter was formed. Now the boy has just as good control of his evacuations as any other child. The anus is funnel-shaped and the sphincter about one-half inch higher than normal.

DR. S. W. KELLEY, Cleveland, Ohio, said that these patients die unless operated on. Cases are on record where after operation they have grown up and become useful citizens. He believes that it is safer to dissect than to use a trochar, and that it is better to dissect out the pouch before opening and bringing it down and attach it to the margin of the incision.

CYSTOCELE.*

CHARLES P. NOBLE, M.D.

Surgeon-in-Chief, Kensington Hospital for Women, Gynecologist to the Stetson Hospital.

PHILADELPHIA.

Cystocele is a hernia of the bladder into or through the vaginal opening. Usually cystocele is described as a prolapse of the anterior vaginal wall and bladder walls. Cystocele as a separate condition is rarely seen. As a rule it is a part of a complex condition, consisting of a laceration of the sacral segment of the pelvic floor (perineum), with prolapse of the anterior and posterior vaginal walls and of the uterus. In order to understand the nature of cystocele and the operative procedures necessary for its cure a knowledge of the anatomy of the female pelvis and of the nature of the supports of the pelvic viscera is necessary. This knowledge is best summed up in the work of Hart and Barbour¹ on the nature of the pelvic floor, of Schultze² on the proper supports of the uterus and their functions, and of Foster³ on the nature of the anterior segment of the pelvic floor. So long as the gynecologists are not agreed on the fundamental facts it will be impossible for them to agree as to the best means of curing cystocele. It is my own opinion that the tonic condition of the abdominal muscles, intra-abdominal pressure, the proper anatomical supporting ligaments of the uterus and of the bladder each play their rôle in the maintenance of the position of the pelvic organs, but the greatest factor is the in-

tegrity of the sacral segment of the pelvic floor or, as it is usually called, the perineum. This is the reason why cystocele rarely exists alone, but is usually secondary to injury of the pelvic floor, and exists as a complication of hernia of the pelvic contents, usually spoken of as procidentia uteri. The exceptions are in cases of marked muscular relaxation of the pelvic tissues, of absorption of the perivaginal fat, of exaggerated intra-abdominal pressure, or a combination of two or three of these factors.

The causative injuries of the pelvic floor involve lacerations of the pelvic muscles and fascia, and later under the influence of intra-abdominal pressure and straining at stool there is overstretching of the fascia. As a result a pouch of the bladder greater or less in size is forced down, together with the anterior vaginal wall, into the vaginal outlet or entirely outside of the vaginal canal.

Fig. 1.—Resection of the anterior vaginal wall complete. Bladder detached from the vagina laterally and from the uterus anteriorly. Deep layer of sutures begun.

If our premises are correct that cystocele, as a rule, is a part of a complicated process, it is quite clear that the cure of cystocele must involve not only restoring the anterior vaginal wall to the normal, together with as much detachment of the bladder as may permit it to rise to its normal position in the pelvis, but in addition the various lesions of the pelvis must be corrected. The proper ligaments of the bladder are of service in supporting this organ, but the bladder itself, together with the uterus, ovaries and pelvic contents, is supported by the sacral segment of the pelvic floor. Another factor of importance is the position of the uterus. Normally the uterus is anteverted, and increased intra-abdominal pressure merely serves to increase the anteversion. On the other hand, if the uterus is retroverted, intra-abdom-

* Read in the Section on Obstetrics and Diseases of Women of the American Medical Association, at the Fifty-eighth Annual Session, held at Atlantic City, June, 1907.

1. Manual of Gynecology, 1905, p. 58.

2. The Pathology and Treatment of Displacements of the Uterus, 1888.

3. The Mechanical Action of Pessaries, Trans. Amer. Gyn. Soc., 1881, vi, 434.