to about thirteen cases of pneumonia; of these, no less than eight were jaw or tongue cases, two were in rectal operations, in one a foreign body was subsequently found in the bronchus, and one was a case in which cough and a slight rise in temperature were, apparently, the only signs of pneumonia. Practically, therefore, we have but one case in 5000 which is at all parallel to the case cited by Mr. Lucas as typical. Further, Prescott, in discussing this very subject in the *Boston Medical and Surgical Journal* for 1895, comes to the conclusion that in an estimated 40,000 cases of etherisation there were only three cases of acute lobar pneumonia. Judging from these sources of information, therefore, Mr. Lucas's experience seems to be very exceptional, and one is almost inclined to commiserate him; but possibly, now that attention has been directed to the matter, the records of future years will show a few more cases, so that after all there may be some justification for the view that cases of pneumonia (as distinguished from bronchial irritation) occur with undesirable frequency. At present, however, the belief in the occurrence of pneumonia after surgical operations appears to be a traditional survival of the pre-anæsthetic and pre-antiseptic days.

The bacillary origin of pneumonia.—If simple bronchial troubles as well as those of a pronounced pneumonic type are included, it must first be shown that there is a specific micro-organism which is the *causa causans* of such conditions. Whatever may be the theoretical probabilities of this, I hardly think that any bacteriologist is prepared as yet to proclaim the discovery of the bacillus of bronchitis. With regard to the bacillus of pneumonia it may be pointed out that Crookshank and Sternberg, in their respective text-books, enumerate no less than three micro-organisms which are said to produce, or are at any rate present in the sputum of, pneumonia—viz., the pneumococcus of Friedlander, an altogether different one of Sternberg, and a third of Klein; and not only so, but it seems to me that the inoculation experiments with any one of these are by no means conclusive, or even uniform. Add to this the fact that one or other of the specific forms referred to is often found in the normal buccal debris, and then I think that we can hardly come to any other conclusion than that, from the point of view of infective organisms, our knowledge of the subject is still very incomplete; it does not carry us much further than the stage of probabilities, or entitle us to say much more than that the pathological conditions and secretions of pneumonia furnish very suitable media for the growth of the organisms in question.

Masks and face pieces as a source of infection.—Assuming for the sake of argument that cases of pneumonia after operations are of tolerably frequent occurrence and that such cases are always due to infective organisms, I would submit that of all places such organisms are least likely to be found in the masks and face-pieces of the anæsthetist. Sternberg, in his "Text-book of Bacteriology," quotes some interesting experiments which tend to support this view. Prolonged exposure to nitrous oxide, though it did not absolutely destroy some organisms that were experimented upon, materially interfered with their development and restrained their growth. Anthrax spores, the most obstinate of known organisms, were not absolutely destroyed by immersion in chloroform, but the cholera bacillus, the typhoid bacillus, and the anthrax bacillus were killed in a very short time when placed in weak watery solutions of the drug (from $\frac{1}{4}$ to $\frac{1}{2}$ per cent.). Ten minutes' exposure to ether destroyed tubercle bacilli. Then, again, it is very well known that absolute alcohol (a constituent of the A.C.E. mixture) is rapidly and permanently fatal to nearly all micro-organisms. I do not know of any experiments that have been made with the organisms contained in pneumonic sputum, but there is no reason to doubt that the result would be in accord with the above. It is not a little curious, too, that the least potent of the drugs referred to, both from the anæsthetist's and from the bacteriologist's point of view,—viz., nitrous oxide—is the one which is probably used more frequently than all the others put together, and under conditions which would be most likely to convey infection. Except when a foreign body has slipped into the trachea it has never, to my knowledge, been recorded, that the use of nitrous oxide has been followed by anything approaching pneumonia. On the other hand, chloroform, the most powerful of anæsthetics and a very strong antiseptic, which is used without a mask, is the anæsthetic most frequently employed in septic jaw and tongue cases, in which the mortality from pneumonia is exceptionally high.

Although I cannot agree with Mr. Lucas in his deductions I should like very cordially to endorse his views as to the desirability, for several reasons, of scrupulous cleanliness whether in the matter of surgical instruments or inhalers. As he is not an anæsthetist it is of course excusable that he should have overlooked several improvements in apparatus, tending to facilitate cleaning, that have been introduced of late years. Among others I would mention the following—viz., as far back as 1887 Mr. Lloyd suggested a light, inexpensive, destroyable bag for use with the portable Clover's inhaler; Dr. Dudley Buxton has adopted an easily cleaned bag for the larger Clover's inhaler; and Mr. Davis has brought forward a metal inhaler of the Ormsby pattern. I venture to think, too, that the celluloid inhalers for the A.C.E. mixture, which I introduced some years ago, are steps in the same direction; and further, some two years or so ago I placed in the operating theatre of Guy's Hospital, and showed at a meeting of the Society of Anæsthetists, a form of metal inhaler, one of the advantages which I claimed for it being that it met the demand for an inhaler which could be subjected to the same aseptic treatment as the instruments employed in the operation.

4 Devonshire-street, W.

Clinical Notes:

MEDICAL, SURGICAL, OBSTETRICAL, AND THERAPEUTICAL.

A CASE OF IMPALEMENT.

BY JOHN JOSEPH STACK, L.R.C.P. EDIN., L.R.C.S. IREL.

ONE night at the end of last February I was called by a police officer to attend a man who was literally hooked on to an iron gate opening off a street in the neighbourhood of New Cross. The spike of the iron railing had entered beneath the chin, penetrated the floor of the mouth, issued through the mouth, and was jammed against the right cheek, the point reaching as far as the internal canthus of the eye. The man was making violent but unavailing efforts to disengage himself, and must have been in that position for a long time, as he seemed to be exhausted from his prolonged struggling. My first idea was to raise his head off the railings, but this was not easy, as the shoulders of the pike-shaped rail were locked in by the narrow horizontal rami of the lower jawbone, so that the head was held fast. I then turned the head in a line with the gate, aided by several constables who lifted the body from the ground. In this position the shoulders of the end of the rail were in the long axis of the "maxillary outlet," so that the man was easily released from his distressing position and was conveyed on an ambulance to the hospital. There was only very slight hæmorrhage; he was intoxicated and unable to give any account of himself. How he got on the gate, whether by deliberately thrusting himself on the spike or whether he was placed there by others, I am unable to say. I am indebted to Mr. M. K. Tebay, house surgeon of the Miller Hospital at Greenwich, for further particulars of the case. The man is now practically well; the wound in the floor of his mouth is granulating very favourably, and he will soon leave the hospital. I ought to add that all the front teeth in the upper jaw were broken off.

New Cross-road, S.E.

NOTES ON A CASE OF WOUND OF THE PALMAR ARCH NECESSITATING LIGATURE OF THE BRACHIAL ARTERY.

BY S. J. ROSS, M.B., CH.B. VICT.

IN May, 1896, I was called to see a boy who had run a spike into the palm of his right hand. I saw him almost immediately after the accident. He had lost a considerable amount of blood, and the hæmorrhage was still very free from the wound. A pad of cyanide gauze over the palmar wound was applied. The elbow joint was flexed over a roller bandage,