exercises. When examples of these diseases can be obtained the didactic teaching is replaced by the clinical, and what I have found to be of especial value in this latter method of instruction is not only the close intercourse with the individual student which it allows, but also the mutual stimulation to criticize and discuss the various features of the case with each other which it engenders, as these exercises are made entirely informal. Certain subjects are also taught from the text-book, but this part of the teaching is very limited, and is soon supplemented by what we are in the habit of calling case teaching. This is usually given at the end of the year, just before the examination, and is a discussion of the picked cases, the history of which has been given to the student (printed) in the previous week, so that he may have the opportunity of studying the case before it is discussed. The teaching of the whole class, both didactic, clinical, by recitations and by case teaching, plays, however, but a small part of what we consider the best methods of instruction in the department of pediatrics. The greater part of the teaching is given in sections of five or six, in the wards of our infants' and children's hospitals, and it is this class of teaching which we think most of at Harvard. The other methods are simply preparatory, and in order to systematize the student's knowledge of the subject and to crystallize his ideas.

Recognizing what I have told you concerning the importance of anatomy, physiology and pathology in the study and teaching of pediatrics, they should be made a part of the curriculum of all advanced medical teaching, as should a course on laboratory therapeutics. We expect that the anatomic department will give an especial course on the anatomy of children, and the physiologic department has intimated that it will also give especial physiologic instruction in connection with infancy and children, preparatory to our later clinical teaching in the hospitals. In this way only can pediatrics be thoroughly and intelligently studied, and the young physician can not acquire a proper knowledge of infants and of children if his anatomic knowledge comprises only the anatomy of a later period, as is usually the case in most of our medical schools in the past. There is just as much reason in preparing students for the clinical study of infancy and childhood, by especial anatomic and physiologic teaching, as that such preparation should be made for the clinical medicine of adults.

Bearing all these facts in mind, we are deducing from them a system for teaching in the department at Harvard certain practical measures which we expect to carry out and perfect when the buildings and hospitals of the new medical school are erected. In the first place, we expect to teach the diseases of the first two years of life, in a hospital built and arranged for this age, and to have another hospital arranged for the needs of sick children between 2 and 12 years of age, with especial arrangements for the teaching of the different diseases as they occur at that period. We have also decided that such diseases as scarlet fever, measles, diphtheria, varicella and pertussis are so essentially connected with the early periods of life that they should be taught in the department of pediatrics, and with this end in view an assistant professor who has control of the fine department for contagious diseases in connection with the Boston City Hospital has been placed in the department of pediatrics.

There is another question that is of importance in looking on the study of pediatrics in this broad way, namely, that a thorough knowledge of the subject is of

inestimable value to the practitioner of medicine, whether he be the so-called general practitioner, or whether he devotes himself entirely to children. The treatment of diseases in children is not necessarily a specialty. The physician can only decide how much of this clinical medicine he is able to practice or to teach well, by gauging his own capabilities to grasp the especial subject and his opportunities of so doing. There is surely enough to be learned and enough to be done in the clinical medicine of infancy and children to more than fill the time and tax the brain of the average practitioner. It is needless to say how important a great deal of knowledge of infants and children is to the general practitioner of medicine. When we consider how it is usually through the younger members of the household that the young physician enters into his practice, it is very evident that much more study in the practice of pediatrics is needed by the average practitioner, especially when we see how numerous are the mistakes and how bad the results of his treatment; in fact, it not infrequently occurs that the child is found to be suffering, not from the terminal processes of the disease which it has just been treated for, but from the inappropriate treatment to which it has been subjected.

In conclusion, let me express the hope that I have not dwelt at too great length on a subject which has only recently gained recognition as one worthy to be raised and placed on a level with other great departments of medicine. It is my belief that as time goes on the leading medical schools of this country will adopt a policy similar to that which I have outlined as prevails in Harvard University, and I shall merely say, in parting, God-speed to the development and teaching of pediatrics in your great city.

THE TREND OF MODERN SURGERY.*

JAMES E. MOORE, M.D.

MINNEAPOLIS.

Westerners as a class have the reputation of thinking well of themselves and of being rather boastful of their achievements. It is very evident that others would not think well of us if we did not think well of ourselves, and it is now conceded by those whose good opinion we prize most highly that western surgeons have accomplished much of which they may be proud and even a little boastful. Most of those present have been obliged to do surgery under a serious handicap. We have had no marble walled operating rooms and no rich endowments. We have been obliged to build our hospitals as we went along and our endowments are yet to come. In short, we have had opportunities only as we have made them. It is this ability to overcome obstacles and to make opportunities that characterizes the western man. Western surgeons are proud of the fact that, notwithstanding the many disadvantages they have labored under, they are not now accused of making up the rear guard in the surgical procession.

At the organization of the American Surgical Association in 1882 one of the leading surgeons of the day said, "In my opinion surgery has nearly reached its limits. I do not see how it can advance much beyond its present height of perfection," yet in twenty years that have elapsed since that statement was made surgery has made more advancement than in all time before. The profession has

* President's Address before the twelfth annual meeting of the Western Surgical and Gynecological Association, held at St. Joseph, Mo., December 29 and 30, 1902.

been swept by such a wave of optimism, aggressiveness and surgical furor that it behooves us to scan closely our present trend. What is there in the present tendencies of surgery that makes for good and what for ill? That the general trend is toward better things all will agree, but that there are some dangerous tendencies all must admit. It is well for us occasionally to see ourselves as others see us. To this end I wish to quote briefly from an editorial which appeared in the *Minneapolis Daily Tribune* on the occasion of Dr. Lorenz's too much advertised visit to Chicago.

Reaction from the Knife.—The brilliant operations in bloodless surgery which the Austrian, Dr. Lorenz, is conducting in Chicago and elsewhere, call attention to a distinct reaction from lavish use of the surgeon's knife which is appearing in the higher levels of the profession. It is time for a counter revolution in this direction. Tremendous impetus has been given to the use of the knife in surgery by the two discoveries of anesthesia and antiseptics. Surgeons began to carve more freely when they found a way to do it without causing torture to their victim. But their power to explore the human interior was limited by the perverse habit of dying after the operation. In many cases this was due to the secondary cause of blood poisoning, which was largely removed by the later discoveries of antiseptic surgery. Then came a perfect riot of cutting and slashing, attended undoubtedly with great benefit to the human race. It would not become laymen to say that this has gone too far, because intelligent surgery always weighs carefully the chances of death and recovery, with or without an operation. If it was a great gain for humanity to learn how to use the knife with less danger to the patient, it will be a greater gain to learn how to dispense with the knife in many cases.

While we can not endorse this editorial, there is some truth in it. Our only defense is that while we do go to extremes at times there is no greater tendency in that direction among surgeons than among any other class of men. Even newspaper men by trying cases "out of court" succeed in hanging some innocent men and assisting some guilty men to escape. Surgeons are accused of being over aggressive, but the aggressor is far less dangerous than the obstructionist, and he has the saving grace of learning something while the obstructionist never learns. Even extremists have their uses. We do not look to extremists for the best in anything, yet if it were not for them we could not know our limitations. The editor just quoted, like many medical men, has a wrong conception of conservatism. He seems to think that conservatism consists in not using the knife, when we all know that the knife in judicious hands is the greatest conservative agent at our command. True conservatism consists in doing that which will the most surely prolong human life and prevent suffering. Surgeons are more truly conservative now than ever before because they save more lives and prevent more suffering than their predecessors ever did. Surgeons should not be held responsible for the work of the would-be surgeons found in every community where surgery is being done. Their ranks are filled up with young men not yet out of their medical swaddling clothes, and with older men without experience and proper training who are seeking for a royal route to fame and fortune. Their work is at times appalling and none deplore it more than the surgeons. The time will come—and it can come none too soon for the welfare of suffering humanity and the honor of surgery—when the law will require of a man that he shall have had hospital experience and shall have shown special qualifications as a surgeon before he will be allowed to perform operations involving life and death. A diploma from the best of medical colleges is no criterion,

for not all physicians, not even all good physicians, can become surgeons.

Medical colleges are trying to turn out better educated medical men by raising their standard of entrance and lengthening their college term. There is no question but that the tendency is toward better education, but there is great room for improvement yet. The lengthening of the college term from two to four years unfortunately does not mean that the student spends twice as much time in studying the diagnosis and treatment of disease. So much of his time is taken up by laboratory work that he has very little more time to study medicine than he had when the course extended over but two years. We must not forget that we are trying to make doctors out of these young men and not embryologists and bacteriologists. We can not all be bacteriologists any more than we can all be surgeons. Much of our laboratory teaching at the present time is theoretical and of very little practical value. There is no crying need of half-baked medical men; why not extend the course to six years and limit the laboratory work to the first two? We certainly should do this or make the laboratory course a requirement for entrance to the medical course, leaving a full four years for the study of medicine. Our institutions have been spending so much money building and equipping laboratories that they have little or none left for hospitals and clinics.

We make this criticism of medical education in order that we may more consistently criticize the teaching of surgery. Our didactic teaching has improved and is still improving very materially. The didactic lecture, given alike to freshmen and seniors, is fortunately a thing of the past. It has given place to graded instruction and recitation. Our clinical teaching of surgery, however, is very faulty. The average surgical clinic has been very aptly likened to a continuous vaudeville performance. There is too much of major and too little of minor surgery. It is of infinitely greater value to a student to see a finger amputated or a fracture treated than it is for him to see an abdominal operation. The students see so much major surgery and so many brilliant results that they naturally think it is easy and that they can begin doing that kind of surgery as soon as they get out of college. This is certainly a bad tendency, but one that is easier to find fault with than to remedy. The remedy is for the medical colleges to furnish the clinician with abundance of material and with the same facilities for graded instruction that are furnished to the laboratory teachers, and to so compensate him financially that his only hope of reward is not through the advertisement he may get from his clinic. Eminent jurists and theologians are not obliged to charge their time spent in teaching to the advertising account. They have too much good sense to allow themselves to be placed in any such position. The position of the average clinical teacher is all wrong, and the humiliating part of it is that it is all his own fault. We as surgeons can fully appreciate the value of laboratory work, but as teachers we should insist that it shall supplement and not supplant the teaching of medicine and surgery. One of the great advances surgeons have made is the supplementing of clinical diagnosis with laboratory diagnosis.

The tendency now is to give the patient a more thorough and careful examination than we formerly did, making use of many scientific mechanical devices to aid us. Each year some new devices are added to our

armamentarium and old ones are being improved on. At the present time no one who is imbued with the prevalent professional spirit thinks of listening to a patient's tale of woe and then writing a shotgun prescription. On the contrary, a careful and systematic physical examination is made and the patient gets fewer drugs than ever before. This method of examination is the outcome of specialism.

The one thing that has had more to do with the advancement of medicine and surgery than any other is specialism. It stands to reason that one who devotes his whole time to one branch of the profession can do better work than he otherwise could. A specialist who has not graduated only yesterday, but who has developed from a general practitioner is certainly in a position to do better for the community and for himself than if he were to try to cover all branches throughout his professional career. One of the bad tendencies of the day is for young men to go directly from college into special work before they know to what line of work they are best adapted and before their knowledge is broad enough for good general work and while it is utterly inadequate for special work. When so many new things were added to medicine specialism became a necessity, and it in turn has developed the science of medicine to a degree that could not otherwise have been attained. As soon as men began to devote their whole attention to some one branch of the profession they became more expert in old methods and discovered many new ones. Twenty-five years ago the oculist and aurist was practically the only specialist known. Then came the gynecologist, but as long as he confined himself to the local application of nitrate of silver, to glycerin and tannin tampons and pessaries, he made very little progress. Very soon, however, he began to do plastic surgery, which led to pelvic surgery. From pelvic surgery he soon passed on up to the diaphragm and out through the abdominal wall to the breasts, when he began to lose his identity and to become a general surgeon. The consequence is that there are fewer gynecologists now than there were ten years ago. The gynecologist developed the science of abdominal surgery, for which we owe him an everlasting debt of gratitude. At the present time the bulk of most general surgeon's work is abdominal surgery. It is absurd to maintain that the surgeon who can do good work in the upper part of the abdomen can not do equally good work in the lower part, and vice versa.

Fifteen years ago there were very few specialists in general surgery. There were none in the West, and, in fact, it was not believed possible for a man to get enough surgery to keep him busy. About this time, however, a few pioneers began to realize the great possibilities of surgery as a specialty, and the consequence is that to-day we have specialists in surgery in every city of consequence throughout the land, and the more surgeons there are the more surgery there is for them to do. New fields have been entered each year and the end it not yet.

Hospitals are being established without number, and it will be interesting to note the results. Many of them being poorly equipped and poorly manned will doubtless close, but others will remain a blessing to their communities and to the profession. The good country hospitals will be helpful to the metropolitan hospitals because they will educate the country people to the hospital idea and teach them the possibilities of surgery. We will find that the more hospitals there are the more surgical cases there will be to fill them. The modern hospital, with its trained nurses, is a very essential part

of modern surgery, and the rule now quite generally adopted by surgeons that every patient requiring an operation must go to a hospital when it is possible, has much to do with advancing surgery. The people in the cities have already learned that a hospital is not a charnal house, but a harbor of safety. We find very little difficulty now in persuading our patrons to go from the finest homes to the hospital. The country people already know something of this, and through the country hospitals will soon learn more. We of the larger cities need not fear the competition of the country hospital, because much that will be done there would otherwise be undone.

A few years ago when our present technic was being developed many of the papers read were on that topic, but now we are so content with our methods that the technical paper is the exception. There is even a little tendency to criticise one who writes a paper on technic as though this subject were settled for all time, but this is wrong, for great as have been our improvements we have not yet attained to perfection. It has been suggested by a recent writer that a uniform standard of technic be established, but it is questionable whether this can ever be brought about, and if it could be done it is not desirable, for technic is the work of human hands and can never be perfect. An established standard would imply perfection which would prevent progress. The early history of antiseptic surgery was an illustration of our tendency to go to extremes. Our very elaborate technic at that time made us seem ridiculous in the eyes of unbelievers and the laity. There is a very wholesome trend now toward simplicity in technic. The best surgeons are those who have the least paraphernalia and who make the least fuss about an operation.

Strong chemicals are rapidly disappearing from our operating rooms and are giving way to soap, water, brush, boiled instruments and rubber gloves. The late Lawson Tait builded better than he knew, for when he thought he was denouncing antiseptic surgery he was really teaching us aseptic surgery. We have even learned that an operation can be conducted more satisfactorily and comfortably without flooding everything with water. There should be but one rule in aseptic surgery, viz., absolute surgical cleanliness, brought about in the simplest manner possible.

The tendency in the West is to abandon all unabsorbable buried sutures. One who becomes accustomed to catgut uses it more and more, until he finally uses it for everything but the intestines and the integument. One accustomed to it finds no objection to it.

One of the most hopeful signs of the times is the great interest surgeons take in each other's work. A spirit of kindly criticism prevails that is helpful to us all. So much good surgery is being done everywhere that it is quite the proper thing now for surgeons to visit each other and make comparisons which are bound to prove beneficial. We have always traveled East for surgical light, but at the present time there is a very respectable and rapidly increasing return current toward the West.

In conclusion: Let us continue to work so that it can not be said of us twenty years hence that we did good work for a few years and then folded our hands and said we have attained to such a height of perfection that farther advance is impossible. Let us continue to work, still admitting that there is room for advancement, not being content with our laurels until we shall have at-

tained the millennium when there will be no disease and no suffering.

Let us then be up and doing,
With a heart for any fate;
Still achieving, still pursuing,
Learn to labor and to wait.

Original Articles.

A MODIFICATION OF PANAS' OPERATION FOR PTOSIS.

FRANK ALLPORT, M.D.

Professor of Ophthalmology, Chicago Polyclinic; Ophthalmologist and Otologist to St. Luke's and Wesley Hospitals, etc.
CHICAGO.

Panas' operation for ptosis probably yields better results, on the whole, than any other operation, and yet its various steps are not always clearly understood by

Fig. 1.—Panas' operation for ptosis. (After Panas.)

operators, and it possesses some glaring defects that have served to render it rather unpopular. It is for the purpose of clearing up these misunderstandings and

Fig. 2.—Appearance after the operation is completed. (After Panas.)

remedying the defects that I have ventured to offer this communication.

I will first reproduce the drawings usually given to clarify the written description of Panas' operation, and it will be seen that they do not entirely clear up some of the points in the ordinary description.

I think that the addition of a plain schematic drawing would better demonstrate the different steps in the opera-

tion, and for this purpose at my clinics I draw a plan like Figure 3.

The line indicated by 1 represents the first incision, which should be in the eyebrow and extend to the bone. The line indicated by 2 represents the second incision, which should be just below the eyebrow, and should extend also to the bone. The bridge of tissue between these two incisions should not be more than 5 or 6 mm. in width. The line marked 7 represents the palpebral margin of the upper lid. The line marked 3 represents a short incision in the lid at about the upper margin of the tarsal cartilage, and should be from 8 to 10 mm. long, and should pass completely through the skin. The line marked 4 represents another similar incision in the opposite angle of the lid. The lines marked 5 and 6 represent two incisions passed completely through the skin and uniting 3 and 4 with 2. While incisions 3, 4, 5 and 6 are being made, the horn spatula should be placed underneath the lid, both to facilitate the making of the incisions and to avoid accidents to the underlying eyeball. The tissue between incisions 5 and 6 should be about 15 mm. in width.

Figure 1, as given by Panas, shows all the lines, etc.,

Fig. 3.—1. First incision. 2. Second incision. 3. Third incision. 4. Fourth incision. 5. Fifth incision. 6. Sixth incision. 7. Edge of upper lid. A. Bridge flap. B. Tongue flap. C. Nasal flap. D. Temporal flap. E. Palpebral flap (nasal side). E. Palpebral flap (temporal side).

to be beautifully curved in outline, but in practical work these curves are not feasible, and I believe straighter incisions can be much more accurately made, and will not detract from the efficiency of the operation.

The original drawing of Panas (Fig. 1) is misleading, as it shows the two threads in the five needle holes in the superior portion of the eyebrow incision, both coming out of the same needle hole, whereas they should be shown as coming out of separate needle holes, side by side and close together, as shown in Figure 4, which I would suggest as in every way a plainer drawing than Figure 1.

The description as given by Panas is also not explicit concerning the sutures 4 and 5, which are placed to prevent a tendency to ectropion induced by the traction effect of sutures 1, 2 and 3. These sutures should include the tissues of the lid except the skin and tarsal cartilage, and should be inserted in the spaces between the skin flaps. One of the needles of a double-needled thread should be inserted directly through the lid from within outwards, just above the upper edge of the tarsal cartilage, and pulled through, after which the other needle should be inserted through the tissues, also from within outwards, and brought out close to the first point of entrance of this suture, leaving the loop of the suture on the conjunctival surface of the lid. The lid should