On the elements of the orbits of Σ 228 and $O\Sigma$ 400.

By *W. Doberck*.

Σ 228. During the past twenty years there has been so marked an improvement both in the number and in the accuracy of observations of binary stars, that at the present time orbits can be determined both more easily and more accurately than formerly. When determining the following elements I began by projecting the distances, and I then projected the angles and the tangents to the interpolation curve. The two curves were then so altered as to agree with each other. It is usually easy to accomplish this after 1860, but before that year the uncertainty is greater. Mr. Gore calculated this orbit in 1889, when the arc described

was only 129° . His elements with the exception of the eccentricity and the periastron resemble the following orbit considerably. Mr. Gore's period was 89 years. The present elements were corrected by equations of condition as the plane of the orbit was badly determined by construction. They are referred to 1900.0.

$$\begin{aligned}\Omega &= 90^\circ 41' & e &= 0.309 \\ \gamma &= 66^\circ & P &= 123.71 \\ \lambda &= 4^\circ 52' & T &= 1905.19 \\ & & a &= 0.899\end{aligned}$$

Observer	Epoch	θ_o	ϱ_o	$\theta_o - \theta_c$	$\varrho_o - \varrho_c$
W. Struve	1831.46	262.91	1.08	- 1.9	- 0.03
Mädler	36.61	267.2	1.04	- 0.5	- 0.12
O. Struve	1841.68	270.8	1.23	- 0.3	+ 0.05
"	42.21	273.8	1.18	+ 2.4	0.00
Mädler	45.15	275.6	0.96	+ 2.1	- 0.21
"	1852.19	280.2	1.11	+ 1.8	0.00
Dembowski	56.76	281.1	1.0	- 0.9	- 0.04
Mädler	57.21	284.2	0.9	+ 1.8	- 0.13
Secchi	57.62	286.9	0.99	+ 4.2	- 0.04
Dembowski	1862.96	286.5	0.92	- 1.5	+ 0.01
"	66.07	291.6	0.95	- 0.2	+ 0.12
Dunér	69.48	299.5	0.71	+ 2.6	- 0.02
O. Struve	1870.18	297.5	0.81	- 0.6	+ 0.10
Wilson	72.04	308.7	—	+ 7.0	—
Dembowski	72.71	299.4	0.5	- 3.8	- 0.14
Wilson	73.93	309.0	0.77	+ 2.8	+ 0.17
Dembowski	74.90	309.4	0.71	+ 0.7	+ 0.14
Gledhill	74.94	307.2	0.6	- 1.6	+ 0.03
Dunér	75.20	311.4	0.54	+ 1.8	- 0.02
Dembowski	76.89	314.9	0.57	- 0.1	+ 0.06
Hall	77.12	313.6	0.54	- 2.2	+ 0.03
Doberck	78.90	327.7	0.47	+ 4.7	+ 0.02
Seabroke	1880.13	329.3	(0.4)	+ 0.3	- 0.02
Hall	81.96	336.2	0.41	- 3.4	+ 0.03
Seabroke	83.16	341.4	—	- 6.4	—
Engelmann	83.50	348.3	0.31	- 2.0	- 0.5

Observer	Epoch	θ_o	ϱ_o	$\theta_o - \theta_c$	$\varrho_o - \varrho_c$
Perrotin	1884.21	11.8	0.53	+ 16.2	+ 0.18
Seabroke	84.21	354.1	0.95	- 1.5	+ 0.60
"	85.23	7.5	—	+ 3.9	—
"	86.17	2.6	—	- 8.5	—
Engelmann	86.34	9.5	0.46	- 3.0	+ 0.12
Tarrant	88.44	20.3	0.41	- 8.6	+ 0.05
Hall	89.00	26.8	0.43	- 6.2	+ 0.06
Schiaparelli	89.02	30.8	0.35	- 2.4	- 0.02
Young	89.02	34.6	0.41	+ 1.4	+ 0.04
Seabroke	89.10	11.5	—	- 22.2	—
Tarrant	89.66	37.0	0.37	- 0.5	- 0.01
Hall	89.98	41.6	0.43	+ 2.0	+ 0.05
Tarrant	1890.10	42.8	0.35	+ 2.5	- 0.03
Hall	90.94	51.3	0.33	+ 5.9	- 0.07
Highton	91.14	40.5	—	- 6.1	—
Bigourdan	91.15	51.1	—	+ 4.5	—
Tarrant	91.80	62	—	+ 11.8	—
Bigourdan	92.05	52.0	—	+ 0.4	—
Highton	93.94	51.1	—	- 9.3	—
Bigourdan	93.99	56.6	—	- 4.0	—
Highton	94.13	61.2	—	0.0	—
Comstock	95.45	69.5	0.44	+ 3.3	- 0.06
Lewis	95.87	69.4	0.37	+ 1.7	- 0.14
Hussey	97.75	78.0	0.57	+ 4.2	+ 0.02
Aitken	98.00	76.5	0.64	+ 1.9	+ 0.09

$O\Sigma$ 400. In this case it was found that the distances were too incompletely observed to be of any help in the first instance. A very rough orbit was constructed by aid of distances derived from the angles. This orbit was corrected by successive trials according to the formula:

$$M_o + \frac{dM}{d\Omega} \Delta\Omega + \frac{dM}{d\lambda} \Delta\lambda + \frac{dM}{d\gamma} \Delta\gamma + \frac{dM}{de} \Delta e = M_c + \Delta M_c + (t - T) \Delta n$$

where M_o signifies a mean anomaly calculated from an observed position angle by aid of the elements Ω , λ , γ and e . Thus Elements II were obtained. I also compared the observations with the elements, which Mr. Burnham has published in M. N. Vol. LVIII page 89, but could find no agreement except by altering λ from 7° , as given by Mr. Burnham, to 353° , and the observations have been com-

pared with the elements thus amended ($\theta_o - \theta_\beta$, $\varrho_o - \varrho_\beta$) although the agreement with Mr. Burnham's calculations still remains general rather than absolute. As the period obtained by Mr. Burnham is about 80 years, instead of about 60 years, the value obtained by me, I give another set of elements (Elements I) with a period of about 100 years. Although the latter do not represent the observations quite

as well as Elements II, they still indicate that this orbit is very uncertain, although the arc described is about 294° , which proves that the accuracy of a set of elements of a double star does not only depend upon the size of the arc described by the companion but depends also upon, and depends mainly upon the number and accuracy of the observations, on which the calculations are based.

The elements are referred to 1900.0.

	Elements I	Burnham's Elements	Elements II
Ω	$= 148^\circ 24'$	$157^\circ 1$	$132^\circ 49'$
λ	$= 11\ 28$	353.0	$14\ 33$
γ	$= 60\ 33$	59.9	$69\ 47$
e	$= 0.5369$	0.46	0.3645
P	$= 1007.12$	817.04	607.92
T	$= 1886.67$	1888.23	1887.58
a	$= 0''.485$	$0''.47$	$0''.533$

Observer	Epoch	θ_0	ϱ_0	$\theta_0 - \theta_1$	$\varrho_0 - \varrho_1$	$\theta_0 - \theta_\beta$	$\varrho_0 - \varrho_\beta$	$\theta_0 - \theta_2$	$\varrho_0 - \varrho_2$
Mädler	1843.39	326.7	0.50	-3.4	$-0''.23$	-11.3	$-0''.18$	-17.0	$+0''.13$
O. Struve	44.83	337.1	0.71	$+7.9$	-0.02	$+0.4$	$+0.03$	-1.5	$+0.29$
»	45.69	335.5	0.72	$+6.9$	-0.01	-0.4	$+0.04$	-0.7	$+0.27$
»	46.68	333.9	0.65	$+6.0$	-0.07	-0.9	-0.03	$+0.3$	$+0.17$
»	1851.81	329.2	0.63	$+4.9$	-0.06	-0.7	-0.03	$+4.6$	0.00
Dawes	53.89	320.5	0.65	-2.2	-0.02	-7.3	0.00	-1.4	-0.02
O. Struve	54.69	321.6	0.59	-0.4	-0.07	-5.4	-0.05	$+0.6$	-0.09
»	58.59	320.8	0.59	$+2.2$	-0.02	-1.8	0.00	$+4.1$	-0.13
»	1861.62	317.7	0.60	$+2.2$	$+0.04$	-0.8	$+0.05$	$+4.2$	-0.12
Dembowski	65.73	310.8	—	$+0.6$	—	-0.7	—	$+1.8$	—
»	68.55	301.6	—	-3.6	—	-3.3	—	-3.6	—
»	1870.51	302.0	—	$+1.2$	—	$+2.0$	—	-0.1	—
»	71.57	300.4	—	$+2.6$	—	$+5.3$	—	$+0.2$	—
»	72.60	314.9	—	$+20.4$	—	$+24.3$	—	$+17.0$	—
»	73.88	274.1	—	-15.3	—	-9.9	—	-20.6	—
»	74.55	287.8	—	$+1.6$	—	$+7.9$	—	-4.8	—
Schiaparelli	75.67	267.9	0.33	-11.8	$+0.10$	-4.0	$+0.09$	-20.5	-0.02
Seabroke	79.82	150.2	0.50	(-85.3)	—	(-78.2)	—	(-100.5)	—
Schiaparelli	1883.62	185.3	0.35	$+4.5$	$+0.17$	-2.0	$+0.13$	$+16.0$	$+0.15$
Young	85.72	144.5	0.3	-16.8	$+0.09$	-26.3	$+0.05$	-4.3	$+0.02$
Smith	85.77	172.2	1.0	$+11.2$	—	$+1.7$	—	$+23.7$	—
Hough	87.71	122.7	0.3	-24.3	$+0.07$	-34.5	$+0.05$	-14.7	-0.03
Schiaparelli	87.92	140.1	0.25	-5.4	$+0.02$	-16.6	0.00	$+3.7$	-0.08
»	88.89	126.1	0.22	-12.8	-0.01	-23.0	-0.03	-5.7	-0.12
»	89.86	123.9	0.23	-8.3	$+0.01$	-18.0	-0.01	-3.4	-0.11
»	1890.93	115.4	0.25	-8.8	$+0.04$	-17.4	$+0.03$	-6.8	-0.08
»	91.93	114.6	0.17	-1.4	-0.03	-8.3	-0.03	-2.1	-0.13
Comstock	92.79	120	0.2	$+11.7$	$+0.01$	$+6.9$	$+0.01$	$+8.7$	-0.08
Schiaparelli	92.91	129.3	0.5	$+22.3$	—	$+17.8$	—	$+19.0$	—
»	93.96	57.7	0.15	-39.0	-0.03	-39.8	-0.02	-44.4	-0.10
See	95.73	346.8	0.23	(-91.0)	$+0.05$	(-84.4)	$+0.06$	(-95.3)	$+0.03$
Hussey	96.90	67.8	0.18	$+2.4$	0.00	$+12.5$	-0.01	$+3.9$	0.00
Aitken	96.92	63.5	0.18	-1.6	0.00	$+8.5$	-0.01	0.0	0.00
»	97.74	47.3	0.20	-9.7	$+0.01$	$+1.7$	0.00	$+1.7$	$+0.03$

Mr. *Plummer*, who has kindly helped me in these calculations, has calculated the following ephemerides:

t	θ_β	ϱ_β	θ_2	ϱ_2
1898.7	36.3	$0''.22$	32.6	$0''.18$
1899.7	28.5	0.25	17.9	0.20
1900.7	22.2	0.28	6.5	0.23
1901.7	17.1	0.31	357.8	0.27
1902.7	12.8	0.33	351.1	0.30
1903.7	9.1	0.36	346.0	0.34

From these ephemerides it appears that it will be possible to decide between these orbits in a few years.

Hongkong Observatory, 1898 Aug. 12.

W. Doberck.