of the left bronchus through the anterior mediastinum is anatomically impossible. 2. A foreign body in either bronchus well down near the lung can be reached, if other means fail and the nature of the case demands it, through the posterior mediastinum. With the arm hanging over the table a vertical incision midway between the base of the scapula and the vertebral column, and extending from the upper border of the third rib to the lower border of the seventh rib, should be made dividing everything right down to the ribs. A flap incision is a disadvantage as free drainage is advisable. The ribs, third to seventh inclusive, should be divided subperiosteally, and from two inches to two and a half inches removed from each. Median division of the pleura is an advantage as its reflection to each side protects the hand from the cut edge of the ribs. 3. By deliberately opening the pleural cavity and inserting one's hand the root of the lung could be easily manipulated, the bronchus incised, and a foreign body removed with greater certainty and certainly with less danger from hæmorrhage than by stripping the pleura from the chest wall. No attempt should be made to close the opening made in the bronchus. The danger of subsequent sepsis would be much lessened by only partially closing the wound and packing from the bottom.

Remarks by Mr. MAYLARD.—My comparatively short association with the case in its initial stages leaves me little to add to what has already been stated by Dr. Andrew. It had been my unfortunate experience to see two cases of death from gangrene of the lung due to the accidental lodgment of teeth within the bronchi; and the gravity of the symptoms which were beginning to show themselves in the case led me to agree with Dr. Andrew that we were fully justified in employing any measures that would reasonably be expected to lead to the extraction of the foreign body or favour its expulsion. Although the attempt to remove it failed it is not altogether improbable that the manipulation involved in the operation may have had something to do with the subsequent dislodgment. The operation was undertaken, as will be learned from Dr. Andrew's own statement, with the strictest appreciation of the gravity of the symptoms which apparently manifested themselves and the severity of the measure it was proposed to employ. Apart, however, from all considerations connected with the operation as employed for this particular case I am of opinion that it is not without its own teaching in regard to the method of approaching the root of the lung, the bronchi, and œsophagus by the posterior mediastinal route; and quite apart therefore from any intrinsic merits connected with the case itself the operation is one which from its great rarity is worthy of being recorded.

Glasgow.

A CASE OF HÆMATOSALPINX DUE TO TUBAL PREGNANCY COMPLICATED BY TORSION OF THE PEDICLE; OPERATION; RECOVERY.

BY FREDERICK J. McCANN, M.D., C.M. EDIN.,
F.R.C.S. ENG., M.R.C.P. LOND.,

PHYSICIAN TO THE SAMARITAN FREE HOSPITAL FOR WOMEN AND CHILDREN, MARYLEBONE-ROAD, N.W.

IN searching through the literature of hæmatosalpinx I have succeeded in finding only one recorded example where torsion of the pedicle complicated a hæmatosalpinx due to tubal foetation. Martin¹ of Berlin describes a case and gives a drawing of the specimen. The patient, a woman, aged 28 years, had a left-sided ampullary tubal pregnancy at about the second month. The peritoneal cavity contained a large amount of sanious ascites. The tumour was bluish-black in appearance and had two twists in its pedicle. When removed it was found to consist of the left tube and ovary. It had a length of 13 centimetres and a breadth of six centimetres. The Fallopian tube was S-shaped and the ampullary end was dilated to the size of a goose's egg. No foetus was found, but there was a placenta infiltrated with blood, and there were also blood, hæmatoidin crystals, and fatty detritus. The ovary was the seat of a large hæmatoma and the ovarian tissue was infiltrated with blood. This case presents certain points of resemblance to that recorded in

this paper, for in both a foetus was absent and the torsion of the pedicle was well marked.

In May, 1898, I was asked by my friend, Dr. D. T. Macleod of Sydenham, to see a patient, aged 34 years, who was then suffering from a purulent uterine discharge. Curetting was recommended and performed. No abnormal condition of the appendages was detected at this operation. As a result of the curetting her general health improved and this improvement was maintained. In October, 1900, she was suddenly seized one evening with very acute pain, said to be in the "kidney region," which extended round towards the costal margin in front. Rest and the application of heat for two or three hours relieved this attack. In March, 1901, this pain recurred and the same treatment was again successful. She did not consult a medical man until her third attack, which occurred in the following April. Dr. Macleod was then consulted and he has kindly furnished an account of this and a subsequent, attack, which I shall give in his own words:—

The patient was certainly suffering most acutely and I came to the conclusion that it was due to renal colic. I gave a hypodermic of morphia and applied warmth. The pain subsided in six or eight hours. The urine failed to reveal gravel and the pain, which was felt well over the ribs under the right breast, did not give me any clue to its exact origin—indeed, intercostal neuralgia was even thought of.

The next attack was the final and was accompanied by a rise of temperature (102° F.). It occurred on May 25th, 1901. The pain was distinctly abdominal, not pelvic, occupying the right lumbar region in front. It was most distressing in its acuteness and I felt certain that there was peritonitis (localised) from the general appearance of the patient and the condition of the pulse.

On May 27th Dr. Macleod detected on examination a swelling in the position of the right ovary which was very tender on pressure; and I saw the patient in consultation with him on June 11th. I had the advantage of anæsthesia

Tubal gestation with twisted pedicle. The figure represents the natural size of the tumour and shows the three twists in the pedicle.

during my examination. A smooth, rounded, slightly moveable swelling was felt occupying the right posterior quarter of the pelvis. As the patient had never been free from pain since the last attack abdominal section was recommended.

On June 15th, assisted by Dr. Macleod, I opened the abdomen. The anæsthetic (ether) was administered by Mr. Rickard W. Lloyd. After the escape of some ascitic fluid a swelling on the right side of the uterus was exposed. It was bluish black in colour, but at certain points a greenish-yellow shade was noted. The intestines adhered to it. The adhesions were carefully separated and the tumour, with its

pedicle, was exposed. Three turns were observed in the pedicle (*vide* illustration). A ligature was applied below the twist and the tumour was removed. The right ovary was not removed. The appendages on the opposite side were normal. The pelvic cavity was washed out with salt solution and the abdominal wall was closed. The patient convalesced rapidly and Dr. Macleod wrote to me that "recovery after operation was uninterrupted and the wound healed by first intention. She has remained absolutely well ever since. The menses are as usual."

The illustration above represents the parts removed in their natural size. The dilated tube contained blood clot and dark fluid blood. The tubal wall was infiltrated with blood. Sections made from the wall showed chorionic villi also infiltrated with blood. The tube was closed at its fimbriated extremity. Mr. S. G. Shattock, who kindly examined the specimens, confirmed this opinion. The patient, who had been married for nine years, was sterile. Her menstruation was always regular.

Although the case narrated together with that described by Martin of Berlin are the only two examples of torsion of a hæmatosalpinx due to tubal foetation which I have been able to find, still such torsion has been described where the hæmatosalpinx was due to other causes. Mr. J. Bland-Sutton² describes and figures a specimen of hydrosalpinx with axial rotation. This specimen, which is now preserved in the museum of the Royal College of Surgeons of England, was removed by Mr. Henry Morris. The Fallopian tube between the cyst and the uterus was tightly twisted three and a half times. The tumour was "strongly adherent." The contents were fluid and of a chocolate colour. Here was an example of hæmorrhage from the tubal wall due to torsion of the pedicle rendering the contents chocolate-coloured. This, as is well known, is what occurs when a similar change (torsion) occurs in the pedicle of an ovarian cyst. Hartman and Delbet³ have also recorded examples of axial rotation of distended tubes. More recently at the Gynæcological Congress held in Vienna von Herff⁴ demonstrated a specimen of hæmatosalpinx with twisted pedicle and Kleinhaus⁵ has also described and figured a specimen.

No doubt the publication of this case may be the means of bringing forth other cases but at the present time it may be said that torsion of the pedicle is one of the rare complications of hæmatosalpinx due to tubal pregnancy.

Curzon-street, W.

A NOTE ON THE SERUM REACTION OF TUBERCLE,

WITH SPECIAL REFERENCE TO THE INTIMATE NATURE OF AGGLUTINATION REACTIONS GENERALLY AND TO THE THERAPEUTIC INOCULATION OF THE NEW TUBERCULIN.¹

BY A. E. WRIGHT, M.D. DUB.,

PATHOLOGIST TO ST. MARY'S HOSPITAL; LATE PROFESSOR OF PATHOLOGY, ARMY MEDICAL SCHOOL, NETLEY.

Technique of the serum reaction.—The difficulties which have stood in the way of the general exploitation of the serum-sedimentation reaction in connexion with tubercle have been technical difficulties associated with the fact that cultures of the tubercle bacillus do not yield uniformly turbid suspensions such as are required for the institution of a sedimentation reaction. Arloing and Courmont showed a way out of this difficulty. They showed that it is possible by a process of careful selection and by frequent agitation of the growing cultures of selected strains of the tubercle bacillus to obtain a uniform bacterial suspension for the purposes of serum sedimentation. Koch introduced an improved technique, the essential feature of which is the grinding up of the tubercle bacilli after previous sterilisation of the cultures under alcohol. The tubercle powder employed—I speak of a specimen kindly placed at my disposal by Professor Dönitz of Berlin—appears

under the microscope as an irregular detritus possessing the typical staining reaction of tubercle and containing interspersed here and there an intact bacillus. This powder is employed in the preparation of the test fluid. As much of the powder as will lie on the point of a very small knife blade is taken, is introduced into an agate mortar, and is then triturated with normal salt solution—the solution being supplied drop by drop until about one cubic centimetre has been added. The suspension thus obtained is centrifuged in a hand centrifuge until the grosser particles have sunk to the bottom, leaving above a clear opalescent fluid. This is the test fluid. When there is mixed and digested for 24 hours with this test fluid a serum which gives the characteristic tuberculous reaction there is thrown down in the fluid a floccular precipitate which aggregates itself into a pellet at the bottom of the containing vessel. The operations of making graduated dilutions of the serum and of mixing with these the test fluid are conveniently undertaken in capillary sedimentation tubes. The technique to be employed is at all points the same as that which I have described in connexion with the typhoid fever and Malta fever reactions.

Question of the intimate nature of the serum reaction.—It will be manifest on consideration that the test fluid which is employed owes its characteristic opalescence to the suspension—probably in a state of semi-solution—in the fluid of a multitude of infinitely fine particles. The essential features of the reaction are possibly in the first instance a process of precipitation, and certainly an agglutination of the finely divided particles, followed by a sedimentation of the flocculated masses. Many convergent lines of work seem to indicate that reactions of this kind play a very large part in the economy of nature and that they are the result of forces derived from the presence of salts in solution. The work of Hardy² and Whetham,³ following upon that of Linder and Picton,⁴ showed that the formation of jellies and the coagulation and precipitation of albuminous substances are brought about by forces of this kind. The latter author in particular showed that the forces exerted are progressively more powerful according as the salts in solution are the salts of monads, diads, or triads. Following upon, though not brought into relationship with, the work of Hardy and Whetham came the work of Bordet.⁵ This author showed that the bacterial agglomerations formed under the influence of an agglutinating serum are (a) resolved into their component elements when these agglomerated masses are transferred to distilled water and (b) re-agglutinated by the addition of salt to the water. Lastly, Joly,⁶ working in connexion with the problem of the formation of sedimentary rocks and the deposition of silt where river water falls into the sea, has shown that more rapid agglutination and sedimentation of inorganic particles (finely powdered slate) are determined by the presence of salts, in particular by salts of diad and triad elements. It may be noted in this connexion that the clarification of muddy water by alum depends on a practical exploitation of the agglutinating forces derived from a salt of the trivalent element aluminium. These forces are inferred to be electrical forces. The interpretation is justified by the generalisation that electrical forces are generated in watery solutions by the breaking up of the dissolved salts into ions, carrying single, double, or treble charges of positive or negative electricity, according as the ions are monovalent, divalent, or trivalent. The competence of such forces to produce the movements of transposition which are associated with agglutination is established by observations which show that a solid body suspended in a fluid may be driven out or, as the case may be, drawn in, between electrodes, according as it offers greater or less resistance to the passage of electricity than the enveloping fluid. A series of simple diagrams—diagrams, let it be remarked, which I owe to suggestions gathered from my friend Professor J. Joly—will serve to bring more clearly before the mind's eye the possible mode of action of the forces which produce agglutination.

In Fig. 1 we have represented two electrodes distinguished respectively with the positive and negative sign. Intervening between these we have a fluid across which the current is conveyed. The lines of force are represented in the diagram by straight lines. In Fig. 2 we have the same

² Surgical Diseases of the Ovaries and Fallopian Tubes, p. 221.

³ Annales de Gynécologie et d'Obstétrique, 1874.

⁴ Frommel's Jahresbericht, 1896, p. 201.

⁵ Veit's Handbuch der Gynäkologie, p. 716.

¹ A communication read before the Pathological Society of London on April 21st, 1903.

² Proceedings of the Royal Society, vol. lxvi.

³ Philosophical Magazine, 1899.

⁴ Chemical Society's Journal, 1895.

⁵ Annales de l'Institut Pasteur.

⁶ Congrès Géologique International, 1900; Proceedings of the Royal Dublin Society, vol. ix. (new series).