evidence, and which has left him unrelieved. What then, surgically speaking, remains? Laparotomy. If, however, a displaced hernia should be found, it can only be relieved by a peritoneal section in a situation where its risks are greater and its management more difficult. It may be said against laparotomy that it precludes any attempt at radical cure; but this, under such circumstances, is surely a minor consideration; and it is interesting to notice that the patient whose case is described above has not yet been troubled by any reappearance of his hernia. If the gut should be gangrenous, opening the sac into the general peritoneal cavity would appear to mean certain death from septic peritonitis; but we have apparently not yet thoroughly explored the resources of antiseptics, peritoneal irrigation, and enterorrhaphy; and I am not aware that there is any case on record in which a patient suffering from displaced hernia in which gangrene had supervened has ever recovered under any method of treatment.

Leeds.

THE DISTRIBUTION OF PYÆMIC ABSCESSSES. By STEPHEN PAGET, F.R.C.S.

IN THE LANCET of April 17th I published sixty cases of that form of parotitis which is secondary to some abdominal or pelvic lesion. This parotitis is very seldom accompanied by any other similar lesion; it is not evidence of true pyæmia, and it usually ends in recovery. There is reason for thinking that its connexion with the abdomen and pelvis is due to the lymphatic system.

Wishing next to see whether, in true pyæmia, the distribution of the secondary lesions is in any way related to the site and nature of the primary lesion, I analysed more than 500 recorded cases of pyæmia. They show clearly that the distribution of abscesses in pyæmia is not merely a matter of chance. Thus, abscess of the liver is related to fracture or section of a bone; pleurisy and pericarditis are more common after injuries of the limbs than after injuries of the head or of the trunk; certain lesions are plainly grouped together in certain sets of cases. It is hard to see how these facts could be explained as mere results of embolism. In all embolism, the emboli, wherever they are first formed, must all start fair again from the heart, after passing through the lung-capillaries; all have exactly the same chance of being lodged in this or that particular organ. If, then, the distribution of pyæmic abscesses is not haphazard, but subject to certain rules and combinations, these abscesses must be due, not to septic embolism alone, but to something else as well. This "something else" is probably a certain state of the tissues themselves, or of the lymphatic system. Perhaps pyæmia may here be compared to syphilis; in syphilis there is the constitutional disease, and also a special liability of certain tissues to the disease, and a special state of the lymphatics, shown by the general glandular enlargement. So in pyæmia, the state of the tissues themselves, and of the lymphatics, may work together with septic embolism in producing suppuration.

The importance of the lymphatics in pyæmia is well shown in Dr. Tilt's paper on "Lymphangitis in Pelvic Pathology" (Obstetrical Society, xvi.). It is easy among my 500 cases to pick out many where the serous and synovial membranes, which are really parts of the lymphatic system, have borne the whole brunt of the disease. Perhaps their importance may show itself if we take the questions which most naturally suggest themselves as to the pathology of the disease, and consider them with the help of the tabulated cases.

I. In pyæmia, does previous disease or weakness of any organ make it more liable to abscess? Thus Guthrie says: "I conceive the viscera in each person most predisposed to disease will be the most likely to be affected." Travers says: "In a person of phthisical diathesis the weight of the injury would fall upon the lungs, whereas in one subject to hepatic congestion it would probably fall upon the organs included in the system of the vena porta." And Lawrence says much the same. Evidence is not easily found in support of these views, but there is some evidence that a simple fracture or subcutaneous injury is likely to suppurate if the poison of pyæmia gets at it. This was shown by the

experiments of Chauveau, Loeffler, and others, and by the following cases:—

1. Primary amputation of right thigh. Pyæmia; death. Post-mortem: "Une fracture de la côte, dont le malade ne s'était aucunement plaint. Le foyer de la fracture est rempli de pus louable et sans odeur." (Sédillot.)
2. Amputation of thigh. Pyæmia; death. Post-mortem: "The fourth rib was found fractured, and a small quantity of pus external to the pleura costalis." (Balfour.)
3. Fracture of both legs and of sternum: Amputation of right leg. Pyæmia; death. Post-mortem: "Left saphena vein full of pus; fracture of sternum bathed in dirty pus." (Path. Soc., xxx.)
4. Scalp wound with fractured skull; fracture of sternum. Pyæmia; death. Post-mortem: "Suppuration in portions of the fractured sternum." (Wilks.)
5. Compound fracture of right leg; fracture of humerus and of ribs. Amputation of leg; death. Post-mortem: "Suppuration round the fractured humerus; no visceral abscesses." (Wilks.)
6. Fracture of lower jaw. Pyæmia; death. Post-mortem: "Suppuration round simple fractures of thigh, humerus, and ribs." (Path. Soc., xxx.)
7. Venesection; phlebitis. Pyæmia; death. "A large abscess formed rapidly in the leg, at the spot where the patient had received a trivial injury just before the venesection." (Castelnau and Ducrest.)

II. Is there such a thing as a predisposition to pyæmia? Dr. Moxon says, "There must be something very special in the constitution of the person." But Braidwood says, "The majority of the victims are the healthiest of our race." Certainly, in some cases the rapidity and extent of the suppuration are astonishing: they seem to run to abscesses, to have a real "prédisposition à la purulence." Yet this may be due to some admixture of pyæmia with scrofula. Mr. Hewett's case is interesting here: A gentleman who recovered from pyæmia after tonsillitis died, four years later, from pyæmia after removal of a wart.

III. Is pyæmia from a slight wound worse than from a bad one? Mr. Bryant says: "After trivial accidents the pathological results were far more severe than after grave accidents. Suppuration involved a larger number of glands, as the liver, the kidneys, the spleen, and the heart." Dr. Chevers says much the same. My cases show a different result, as follows:—

		Liver.	Spleen.	Kidneys.
Severe injury of a limb, such as amputation, excision, compd. fracture (195 cases).	Marked disease	55	29	13
	Congestion &c.	15	19	14
Slight injury of a limb, such as venesection, or dissection wound, or whitlow (45 cases).	Marked disease	1	1	0
	Congestion &c.	0	2	1

This is roughly reckoned; but it is evidently opposed to the view taken by Mr. Bryant.

IV. The state of the heart in pyæmia. Dr. Wilks says of his large collection of cases: "In several cases it will be seen that the heart has been affected by purulent deposits; and it will be noticed that this occurred only in the very worst forms of the disease, and especially where a diffused abscess in a limb had been the source of the mischief." The heart is a muscle; and some muscles are more liable to pyæmic abscess than others. But perhaps it is not safe to argue from other muscles to the "musculus nobilissimus." The 500 cases appear to show that true abscess of the heart in pyæmia is most common after some lesion of the lower limb. Thus, in 97 cases of pyæmia from some genito-urinary lesion, not one had true abscess of the heart. One had purulent infiltration of the mitral valve; one had mottling of the heart, with softening patches; six had ecchymosis of the heart. But ecchymosis may proceed from very different causes—as from poisoning by strychnia, &c. In 77 cases of pyæmia from some lesion of the head or neck, 3 had true abscess, 3 had ulcerated valves, and 3 had ecchymosis. In 96 cases of pyæmia from lesion of the upper limb, 4 had true abscess, 1 had myocarditis, and 4 had ecchymosis. In 243 cases of pyæmia from lesion of the lower limb, no less than 21 had true abscess, 2 myocarditis, 4 ulcerated valves, and 9 ecchymosis. Thus abscess of the heart wall seems to be most common in pyæmia when the primary lesion is of the lower limb. The primary lesion was not, in all the cases, severe: in one it was a sore on the heel; in another it was a

scratch on the foot. It is certain that pyæmia which starts from the lower limb does, as a rule, have a severe course, affecting many organs and many joints. Perhaps this fact is related to the fact that pyæmia does, to begin with, choose wounds of the lower limbs for its primary seat more often than wounds of any other part. Thus, in the American war, the statistics were as follows:—

After wounds of the lower limb, pyæmia occurred in	18.1	per 1000
" " upper limb, "	10.4	"
" " trunk "	5.9	"
" " head and neck "	2.5	"

It seems natural that the poison which has entered the body at its favourite site should act with the full measure of its severity.

V. In pyæmia, is the inflammation of the serous membranes due merely to extension of inflammation from the viscera underneath? Thus Dr. Wilks says, "Inflammations of the serous membranes are altogether secondary to the inflammation of the tissues which the serous membrane covers." And Sédillot says much the same. It is hard to see why pleurisy and pericarditis in pyæmia should not be independent of pneumonia and myocarditis, just as the effusions into the joints are independent of the heads of the bones, or just as ordinary pleurisy and pericarditis are independent of pneumonia and myocarditis. Cases are not rare where there are large effusions into the pleuræ, but nothing wrong noticed in the lungs; and where there is pericarditis, but nothing wrong noticed in the heart. My 500 cases give the following figures:—

{ Cases of heart disease, no mention of pericardial disease ...	35
{ Cases of pericardial disease, no mention of heart disease ...	31
{ Cases where both heart and pericardium were diseased ...	25
{ Cases of lung disease, no mention of pleura ...	202
{ Cases of pleural disease, no mention of lungs ...	21
{ Cases where both lungs and pleuræ were diseased ...	191

Of course there are cases where the serous inflammations do result only from underlying inflammation, and are proportionate to it. But the following cases show that serous inflammations in pyæmia may occur separately:—

1. Wrist disease (male, aged forty-three). Post-mortem: Lungs healthy. In the right pleura two pints and a half of semi-purulent fluid; in the left pleura three pints. Pus in right knee. Spleen dark and soft. Veins healthy. (Path. Soc., xxx.)
2. Chilblains on fingers, phlebitis (female, aged thirty-six). Post-mortem: Pleurisy same side; bloody fluid in pericardium; serous meningitis. (Ribes.)
3. Injury to hip-joint at birth; death in seven days. Post-mortem: Pleura on same side filled with pus; abscess over wrist. (*Med. Times and Gaz.*, 1863.)
4. Purulent otitis. Post-mortem: A pound of puriform fluid in the opposite pleura; the lung collapsed. (Abercrombie.)
5. Purulent otitis (male, aged thirty). Post-mortem: Pleurisy on same side; pericarditis; peritonitis; abscess in opposite lung; abscesses in liver. (R. Reynolds, "System of Medicine.")
6. Dissection wound of finger. Post-mortem: Lungs congested; eight ounces of bloody fluid in each pleura; serum in pericardium; serous meningitis. Pus on knee in same side; ecchymoses of kidneys and intestines. (Travers.)

There are many more cases like these. Pleurisy, pericarditis, and meningitis seem to group themselves together, and to occur in association with effusions into the joints and into the synovial sheaths of the tendons, rather than with visceral abscesses.

VI. The favourite sites of abscess in pyæmia. First, of course, the lungs. Next, among the muscles, the pectorals and the calf muscles suffer most often—about one case in thirty. Abscess of the pectoral muscle appears in some cases to be due to the spread of suppuration from the axilla or neck, or from around the shoulder-joint or the sterno-clavicular joint. But in others it seems to begin in the muscle, or between the major and the minor, or beneath both. The following cases show this:—

1. Gonorrhœa (male, aged twenty-two). Effusions into various joints; pericarditis; large abscess in pectoral region, involving shoulder-joint. (St. Barthol. Hosp. Reports, vol. iii.)
2. Fractured base (male, aged fifty-four). Abscesses in liver and kidneys; an enormous abscess under the pectoral muscle. (Lebert.)

3. Venesection of right arm; purulent clot in vein. Lungs hepatized; pleurisy; meningitis; thick greenish pus in interfibrillar tissue of right pectoral muscle. (Arnott.)
4. Gangrene of left leg after old gunshot wound; pus in veins. Abscess, apex of left lung; meningitis; large abscess under pectoralis major. (Ribes.)
5. Necrosis of left femur. Large abscess in course of pectoral muscles. (Guthrie.)
6. Necrosis of left femur. Lungs, heart, and kidney all diseased. Abscesses in the pectoral muscles. Path. Soc. xxx.)
7. Amputation of left great toe (male, aged forty-one). Pus in veins. Abscesses in lungs; an enormous abscess under left pectoral, passing down into axilla. (Sédillot.)

I have notes of fourteen or fifteen cases of this kind of abscess.

Abscess of the calf seems to begin in or near the soleus. It is to be observed that in three-fourths of the cases where it occurred there was also suppuration on in one or both of the upper limbs. Among sixteen or eighteen cases are the following:—

1. Puerperal fever. Abscess of left axilla; abscess of left calf; destructive suppuration of left eye. (Marshall Hall.)
2. Injury of vagina; erysipelas. Abscesses over shoulder, in elbow, and in each calf; superficial abscesses elsewhere. (*Med. Times and Gaz.*, 1863.)
3. Scalp-wound with fracture. Abscesses of liver, lungs, and abdominal wall; large abscess in arm; large abscess in leg. (Wilks.)
4. Amputation of right thigh. Abscess under left soleus, close to interosseous ligament. (Guthrie.)
5. Amputation of left foot. Abscesses of left lung; abscess near right elbow; abscess in right calf. (Wilks.)
6. Excision of varicose vein, right leg. Abscess of right lung; meningitis; inflammation of both eyes; intra-muscular abscesses in both forearms; deep abscess in left leg. (Arnott.)

Another favourite site of abscess in pyæmia is the back of the hand. This abscess is not due to suppuration in the wrist-joint or in the finger-joints; it seems to begin in or around the synovial sheaths of the extensor tendons. I have notes of 16 cases, and it is interesting that 12 were on the left hand and only 4 on the right. Of these 16, one or two were not "abscesses," but "pustules" or "vesicles"; still, I venture to include them. There is some reason for thinking that this abscess at the back of the hand is related to the state of the lymphatic system. It is in relation to a synovial membrane, and in a part of the body where the lymph spaces are extraordinarily large; it is found along with other synovial and serous effusions rather than with visceral abscesses; it seems, by its preference for the left hand, to be not merely due to haphazard embolism; and, lastly, it seems, like parotitis, to be in some way influenced by the site of the primary lesion. In one very interesting case of chorea, ending fatally, in which I made a post-mortem, I found double parotid abscess, and one at the back of the left hand, and no other evidence of pyæmia. The following are also interesting examples of this lesion:—

1. Delivery; endometritis. Abscess at back of left hand, with sloughing; pus up the forearm; no trace of disease in bloodvessels. (Lee.)
2. Delivery; abscess in lung; meningitis. Abscess in extensor sheaths of both hands. A case of lymphangitis. (Tilt.)
3. Cystotomy for a foreign body. Both knees full of pus; pus in left elbow and in sheaths of extensor tendons of left hand. (Path. Soc., xxx.)
4. Removal of part of right clavicle; pleurisy; pericarditis. Pus in left sterno-clavicular joint; pus in sheaths of left extensor tendons. Veins normal. (THE LANCET, 1864.)
5. Fracture of right ulna. Abscesses in lungs and in liver; pericarditis; meningitis; pus in right elbow and shoulder; abscess at back of left hand. (Amer. War, vol. iii.)
6. Wound of right finger; purulent infiltration of right arm; veins diseased. Abscess of left lung; pericarditis; abscess at back of left hand. (Amer. War.)

There is one more special lesion of great interest—inflammation of the eyeball. I have notes of 18 cases. Like the

inflammation of the back of the hand, it shows a marked preference for the left side over the right; like it, a very large proportion of the cases have followed uterine disease. These facts, with what we know of parotitis, may make us inclined to think that septic embolism alone does not fully explain this lesion. This ophthalmitis must be distinguished both from phlegmonous inflammation of the orbital cellular tissue and from inflammation extending into the orbit from the base of the brain. It is true inflammation of the globe itself, with subretinal effusion and purulent infiltration of the vitreous. It is beautifully described in Bowman's "Diseases of the Eye." The best examples are Marshall Hall's six cases after delivery, all of the left eye; also the following:—

1. Mrs. E— (aged thirty-five). Three weeks after delivery, right phlegmasia dolens; three weeks later, pyæmia, with destructive inflammation of both eyes. Post-mortem: Pus in left lung; two pints of purulent fluid in left pleura; plugging of veins of limb.
2. Mrs. — (aged thirty). Delivery; peritonitis. Inflammation of left eye. No post-mortem. (Mackenzie.)
3. Mrs. —. Delivery: adherent placenta. Meningitis; inflammation of right eye. (Mackenzie.)
4. Arnott's case, quoted above.
5. Venesection for pleurisy (male, aged thirty-five). Meningitis; destructive inflammation of both eyes. (Mackenzie.)
6. Abscess in right leg (male, aged eighteen). Swollen and painful joints; pus in left knee. Destructive left ophthalmia; pus round both globes, especially left. Sinuses healthy. (Wilks.)

Admirable descriptions of this lesion are to be read in the *Archives of Ophthalmology*, vols. ix. and xi. There is abundant evidence that septic embolism may cause it, but I venture to think that the state of the eye itself, or of the lymphatic system, may in some of the cases co-operate with the work of embolism and predispose the eye to suppuration.

VII. Section of the bones, as influencing the production of abscess of the liver. I was struck by finding how much more common abscess in the liver is, after division of the bones of limbs, than after other injuries. Sédillot says: "Nous avons été frappés d'une sorte de coïncidence entre les suppurations du foie et les altérations du système osseux chez beaucoup de nos pyoémiques. Un vénérien, atteint d'une ostéite du pubis avec infection purulente, avait des abcès du foie: plusieurs amputés, dont les os étaient malades, nous offrirent les mêmes lésions." My tables run as follows:—

		Abscess (not counting congestion &c.)			
		Liver.	Spleen.	Kidneys.	
Injury or disease of medulla.	Upper limb (52 cases).	20	...	8	2
	Lower limb (165 cases).	36	...	20	4
Necrosis, in most cases acute.	Upper limb (4 cases).	0	...	1	3
	Lower limb (34 cases).	2	...	4	14
Suppurative arthritis.	Upper limb (2 cases).	0	...	0	0
	Lower limb (17 cases).	4	...	3	5
Injury or disease of soft parts only.	Upper limb (40 cases).	4	...	0	3
	Lower limb (22 cases).	0	...	3	5

Putting the two limbs together, we have—

		Abscess (not counting congestion &c.)			
		Liver.	Spleen.	Kidneys.	
Injury or disease of medulla: amputation, compound fracture, caries (217 cases) ...		56	...	28	6
Necrosis (38 cases) ...		2	...	5	17
Suppurative arthritis (19 cases) ...		4	...	3	5
Injury or disease of soft parts only (68 cases) ...		4	...	3	8

This seems to prove that abscess of the liver and of the spleen is promoted, in pyæmia, by injury or disease of the medulla of the bones. Now, the medulla of bones is part and parcel of the lymphatic system, and so is the spleen. Do we not see here another instance of the influence of the lymphatic system in the distribution of pyæmic abscesses? The association of abscess of the kidneys with acute necrosis is also important.

ON TINNITUS AURIUM AND ITS TREATMENT BY ELECTRICITY.

By JULIUS ALTHAUS, M.D., M.R.C.P.,

SENIOR PHYSICIAN TO THE HOSPITAL FOR EPILEPSY AND PARALYSIS, REGENT'S PARK.

TINNITUS AURIUM, when loud and continuous, is one of the most distressing affections, and apt to greatly depress the tone of the mind and of the whole nervous system of the patient. St. John Roosa¹ mentions the case of a gentleman who consulted him for it, and received an unfavourable opinion as to recovery; and who thereupon said that, if he thought it correct, he would put an end to his existence which he did a few months afterwards by blowing out his brains! The same author speaks of another patient who committed suicide on account of this symptom; and most practitioners must have met with cases in which all enjoyment of life has been destroyed, the patient's temper soured, and his power of work greatly reduced, through tinnitus. This symptom is probably always produced by increased pressure on the labyrinth, and therefore occurs in a variety of affections of the external, middle, and internal ear, and in the neighbourhood, giving rise to this undue pressure. Sir William Dalby² says that, when it is due to some curable local cause, it rapidly disappears with its removal; but that when it accompanies deafness in nervous affections, it is often the more troublesome symptom of the two, and less amenable to treatment.

The fact that we are able, by a judicious and careful application of electricity, to cure or relieve a considerable proportion of this latter class of cases, where there is no coarse lesion giving rise to the undue pressure, but where this latter is probably owing to disturbed vaso-motor influence, is either ignored or doubted in most of the best works on aural diseases. This unfavourable opinion of the authorities is undoubtedly owing to the circumstance that there is perhaps no other complaint the successful treatment of which by means of electricity requires so much special knowledge of the physiological effects of that agent, such delicate adjustment of apparatus, and such technical skill in manipulating the same, as tinnitus of nervous origin; while, on the other hand, a haphazard application of electricity, even by those who are tolerably well acquainted with the other more common uses of the battery, is likely to do rather harm than good, and the chance of thereby accomplishing a cure must be considered as exceedingly remote. I do not intend in this paper to give an account of the researches of Brenner, Hagen, Moos, Erb, and others, which have opened up this subject, but proceed at once to the narration of a well-marked case of tinnitus of vaso-motor origin which was recently under my care, and in which the beneficial influence of a carefully adjusted electric treatment was strikingly displayed.

S. N—, a merchant, aged thirty-five, married, and father of two children, consulted me in October last, and informed me that about two years ago he had, in consequence of unusually hard work and great anxiety in business, got into a low state of health, with dyspepsia, constipation of the bowels, and considerable mental depression. While in this condition he heard one morning, on awakening, a curious noise in his left ear as if a kettle was boiling there, and had a feeling of giddiness on changing the position of his head. The noise continued, and in the course of a few days became much louder, so as to resemble the breaking of the waves on the sea-shore, and the hearing of the same ear became much impaired. The patient was treated with alteratives, strychnia, iodide and bromide of potassium, &c., and was eventually advised to take a long holiday. From this he returned rather better, but after having been in business again for a few days the trouble returned worse than before. Further treatment did not improve the condition, and matters at last began to assume a serious aspect, inasmuch as the patient's power of application was very much diminished, his confidence in himself almost gone, and the irritability of his temper so extreme that his family as well as his associates in business found the greatest difficulty in bearing with him. Under these circumstances the question was mooted whether he should not give up business, at least for

¹ The Diseases of the Ear (London, 1885), p. 344.

² Quain's Dictionary of Medicine (London, 1882), p. 1636.