negative conjunctival reaction 67 days previous to death. One gave positive reactions to both tests 73 days before death.

From the above we conclude that in known tuberculosis the kind of reaction has no bearing on the prognosis of the case. Positive reactions have occurred with us as early as one day before death; both reactions have been present in variable percentage as long as three months previous to death. The negative reactions have practically equaled the positive reactions in occurrence two or three months previous to death.

UNTOWARD EFFECTS OF THE CONJUNCTIVAL REACTION

The evidence of accumulating cases has unquestionably proven that the conjunctival instillation of tuberculin is not entirely free from danger. On the contrary, from the following report one must conclude that there is a liability of producing severe inflammatory processes in normal eyes, of aggravating non-tuberculous affections of the eye, and of producing an extension, and in some cases a rapidly destructive process, in tuberculous affections of the eye, from its use. McKay⁶¹ reports a severe conjunctivitis lasting over ten weeks in an eye that was normal previous to the instillation of a 1 per cent. solution. Eppenstein⁶² reports three cases of conjunctival phlyctenular keratitis which were greatly exaggerated, and says that chronic hyperemia, blepharitis or conjunctivitis of any kind contraindicates its use.

Ferrien⁶³ reports a conjunctivitis lasting two months, which terminated by having the aspects of tuberculosis of the conjunctiva. Babier and Renon⁶⁴ have observed conjunctivitis persist for one or two months. Stevenson reported three cases of follicular conjunctivitis which existed from 20 to 40 days. Kalt⁶⁵ reports serious results in iritis, sclerokeratitis and iridochorioiditis. Trousseau⁶⁶ reports a marked extension of a tubercle of the conjunctiva immediately following the instillation of a 0.5 per cent. of tuberculin. In another case a tuberculous iritis was greatly aggravated, although the reaction was negative. Babier⁶⁴ reports keratitis and ulceration of the cornea from a conjunctival reaction. A large majority of other authors, not oculists, have observed more or less unfavorable results varying in degree and severity from this test.

Besides these local untoward reactions Wolff²³ observed a very marked febrile reaction in the conjunctival test. Wolff-Eisner⁶⁷ and many others obtained general reactions. Proff noted that the tuberculo-opsonic index was modified by these reactions, and that in some cases the negative phase was marked and unusually prolonged.

We wish to take this opportunity of expressing to senior house physicians, Dr. R. Bunch and Dr. F. Esselbruegge our appreciation for services rendered us in the preparation of this article.

CONCLUSIONS

The diagnostic value of these reactions in adults is restricted to many limitations:

1. The specificity of these reactions have as yet not been demonstrated.
2. The activity or extent of the lesion bears no definite relation to the degree or kind of reaction.
3. The conjunctival reaction is not free from danger.

61. Boston Med. and Surg. Jour., March 12, 1908.

62. Med. Klin., Sept. 8, 1907.

63. Bull. Soc. d'Ophth. de Paris, Oct. 8, 1907.

64. Bull. et mém. Soc. d. hôpitaux de Paris, Dec. 12, 1907.

65. Bull. Soc. d'Ophth. de Paris, Oct. 8, 1907.

66. Jour. de méd. et de Chir., Jan. 10, 1908.

THE CONDITION OF THE PERIPHERAL BLOOD VESSELS IN SHOCK*

M. G. SEELIG, M.D. AND E. P. LYON, Ph.D.

ST. LOUIS

From a purely historical point of view the literature devoted to shock may be divided into three periods: 1, From 1795, when the word shock was first used as we now use it, down to the beginning of the era of experimentation in 1870; 2, the period from 1870 to 1885, devoted to careful experimental investigations; and, 3, from the reopening of the question by Crile in 1889 down to the present time. From 1885 to 1889 practically no papers other than purely casuistic ones appeared on the subject.

THEORIES IN REGARD TO SHOCK

The excellent summary published by Mansell Moullin¹ in 1882, and the encyclopedic volume contributed to the subject by Groeningen² in 1885 render it unnecessary to restate the historical development of the theme. It is of more than passing interest, however, to note that those views regarding shock which are most commonly accepted to-day differ but little from opinions published nearly half a century ago. It does not seem to be known generally that as early as 1864 W. W. Keen, S. Weir Mitchell and G. R. Morehouse³ enunciated practically the same theory as that put forward by Crile nearly forty years later. I have been able to find only one reference to this work by Keen, Mitchell and Morehouse, and that reference was in a German monograph. The experimental work of Fischer⁴ in 1870 confirmed the theory proposed by Keen, Mitchell and Morehouse, and set on a fairly firm basis the idea that shock was due to vasomotor exhaustion. Leyden,⁵ however, took issue with Fischer and announced his belief to be that shock was due to a reflex inhibition of the activity of the centers of the cord. Meltzer,⁶ in the most recent paper contributed to the shock question, practically coincides with the theory of Leyden. The theory recently expounded by Boise,⁷ that cardiac exhaustion is the main underlying factor in shock, is a reiteration of the conclusion of Blum⁸ published in the seventies.

It is in no disparaging sense that we emphasize the fashion in which old views have been rehabilitated. Indeed, we feel that every contribution that we have mentioned has been of importance. Crile's⁹ work, for example, furnishes the most comprehensive data ever contributed to the shock problem and has been instrumental in starting anew investigations as to the cause of one of the most dreaded surgical complications. Our purpose is rather to express agreement with Meltzer, when he says that despite all theorizing and experimentation, "the bringing forward of new experimental facts capable of shedding some more light on our puzzling problem is not yet a superfluous luxury."

Meltzer, in his paper, points out the fact that Crile fails to draw a distinction between surgical shock, with

* An Experimental Study from the Physiological Laboratory of St. Louis University Medical School.

1. Mansell Moullin, C. W.: International Encyclopedia of Surgery, London, 1882.

2. Groeningen, G. H.: Ueber den Shock. Weisbaden, 1885.

3. Circular No. 6. Surgeon-General's Office, 1864.

4. Fischer, H.: Samml. klin. Vortr. (Volkman's), 1870, No. 10.

5. Leyden, E.: Samml. klin. Vortr. (Volkman's), 1870, No. 2.

6. Meltzer, S. J.: The Nature of Shock, Arch. Int. Med., July, 1908, i, 571.

7. Boise, E.: Am. Jour. Obst., 1907; Tr. Am. Gynec. Assn., 1908.

8. Blum, A.: Arch. gén. de méd. de Paris, January, 1876.

9. Crile, G. W.: Blood Pressure in Surgery, Philadelphia, 1903; An Experimental Inquiry into Surgical Shock, Philadelphia, 1899.

rapid onset of symptoms, and traumatic shock, with slow onset of symptoms. This criticism makes it incumbent on anyone who deals with this problem to define his terms at the very outset. The term "shock" has assumed a medical significance so peculiarly its own that the primitive idea expressed by the word has almost been lost sight of. Originally "shock" signified a concussion or a blow, but toward the middle of the eighteenth century English medical writers began to use the word in explanation of the subtle and unexplained effects resulting from a severe blow or concussion. This latter use became so common that the word finally stood, not for the force itself, but for its resultant destructive tendencies.¹⁰ The essence of this idea is so inherently bound up in the word that the French and German schools have been forced to adopt the word bodily, after numerous and futile attempts at translation by such terms as *impressionnabilité extraordinaire*, *léthargie des blessés*, *épuisement par la douleur*, *résolution général des forces*, *syncope*, *Wundstupor*, *traumatische Torpor*, *Schuss-Erschütterung*, *Wundschreck*, *Neuroparalyse*, etc.

This difficulty in translation emphasizes a fundamentally important fact, namely, that the word represents an intricate symptom-complex and does not stand for a specific clinical actuality. The elder Gross¹¹ furnished the most graphic and satisfying description of this symptom-complex when he said that it was the manifestation of a rude unhinging of the machinery of life. It is confusing to adopt Meltzer's views regarding traumatic and surgical shock, for, after all, we assume that every type of shock results from trauma. Scientific accuracy will not admit the use of Howell's¹² classification of "cardiac" and "vascular" shock, for other vital functions besides the cardiac and vascular ones have been shown to be involved. So we shall use the word in the sense of an unhinging of vital machinery, bearing in mind that the onset of symptoms may be sudden or rapid, or at times insidious, and that either erethism or torpor may be characteristic symptoms.

As the problem stands to-day there are several contradictory theories. Crile and his followers maintain that vasomotor exhaustion is the primary cause of shock, Boise states that cardiac exhaustion is the prime cause, whereas Howell believes that both these factors (cardiac and vascular) must be reckoned with. Meltzer, after a critical survey of the literature, concludes that shock is consequent on the development of the inhibitory side of all the functions of the body. Kinnaman¹³ argues for a disturbance of the thermogenic function, Jaboulay¹⁴ for the formation of an irreducible oxyhemoglobin, Vale¹⁵ for an increased specific gravity of the blood, Henderson¹⁶ for the reduction of the carbon dioxide content of the blood and tissues (acapnia), Bainbridge and Parkinson¹⁷ for a loss of chromaffin tissue following

trauma, and Schur and Weisel¹⁸ for a loss of this same tissue following the administration of volatile anesthetics.

Our object is neither to criticize these various theories nor to add a new theory to the already formidably long list. Cardiac excitation and inhibition, vascular dilatation and contraction, the *modus operandi* of the sympathetic nervous system, the complex phenomena of exhaustion and inhibition, would all have to be taken into account by a theory, despite the fact that not one of the factors that we have mentioned is understood fully.

This much is certain, however: that low blood pressure is a constant accompaniment of a pronounced degree of shock—so constant, indeed, that investigators practically always express the state of shock in terms of manometric pressure. We have based our experiments on the fact that the blood pressure is always low in shock, and we have set as our problem the determination whether the peripheral vessels are contracted or dilated during the time that the pressure is low in the larger, more centrally situated vessels, such as the carotids.

Mitchell, Morehouse and Keen, Groeningen and others assert that the peripheral vessels are not contracted, because, in the first place, no one has ever shown that they are, and, in the second place, because it is not probable that the peripheral vessels can be contracted tightly for any great length of time.

Crile argues that, consequent on vasomotor exhaustion, the peripheral vascular system is paralytic and, therefore, relaxed. Porter,¹⁹ on the other hand, denies Crile's contentions *in toto*, and seems to show that the vasomotor center is not exhausted. Neither Crile nor Porter, however, directed their attention to the vessels themselves.

Malcolm²⁰ of London, dealing with the problem from a purely clinical point of view, makes two assertions: 1, that the peripheral vascular system is contracted in shock; and, 2, that a state of low blood pressure is thoroughly consistent with contracted peripheral arterioles. Malcolm makes an uncommonly strong clinical argument against the views of Crile in a fashion that refreshingly demonstrates the invaluable aid of clinical medicine to laboratory experimentation. If the theory of vasomotor exhaustion is correct, says Malcolm, then the network of smaller peripheral arterioles, being cut off from central control, must be relaxed. This relaxation should result in an overfilling of the vessels, but, as a matter of fact, all clinical evidence speaks against such an overfilling: the body surface is cold and pale, the mucous membranes are anemic, incisions made during the state of shock are almost bloodless, the radial pulse is often small to the degree of being impalpable, and the application of surface warmth tends to lessen the symptoms of shock, despite the fact that warmth usually relaxes vessels.

There are three possible ways of determining whether the peripheral vessels are dilated or contracted: 1, the vessels may be inspected visually; 2, the outflow of blood from an organ or extremity may be measured; 3, coincident variations in venous and arterial pressure may be measured. In our work, which has been done on dogs, we have used only the first two or three methods. We studied visually the vessels of the retina and measured the outflow from the femoral vein. We selected

10. It is interesting to note the beginning of a tendency to modify the character of the word again, so that in addition to signifying the destructive results of external force, it will also express the idea of constructive results. For example, Wetherill has published an article entitled "The Conservative Agency of Shock," *THE JOURNAL A. M. A.*, March 12, 1908, xxx, 578. Furthermore, we find in recent biologic contributions, statements such as: "Mutation in species may be considered to result from a shock to the organism, the shock being stronger than normal hereditary impulses." In other words, shock enables the organism to construct something new, and possibly useful.

11. Gross, S. D.: *System of Surg.*, Philadelphia, 1872, i, 426.
12. Howell, W. H.: *Vaughan Festschrift*, Ann Arbor, 1903, p. 51; *Am. Med.*, 1904, p. 482.

13. Kinnaman, R. C.: *Internat. Jour. Surg.*, 1904, p. 378.

14. Jaboulay, L.: *Lyon méd.*, 1904, p. 378.

15. Vale, F. P.: *Med. Rec.*, Aug. 27, 1904.

16. Henderson, Y.: *Am. Jour. Physiol.*, 1908, p. 126.

17. *Lancet*, 1907, p. 1269.

18. *Wien. klin. Wchnschr.*, Feb. 20, 1908.

19. Porter, W. T.: *Am. Jour. Physiol.*, 1907-08, pp. 399, 444, 500.

20. Malcolm, J. D.: *Lancet*, 1905, i, ii, 573, 618, 737, 922; *Med. Chir. Tr.*, 1908.

the eye as the site of visual inspection of the caliber of the vessels, because this is the only situation where the vessels can be studied without the necessity of exposing them by dissection. It is unnecessary to say that the caliber of a vessel that has been traumatized by dissection affords no reliable index of the caliber of that same vessel in an undisturbed state. We selected the femoral vein from which to measure the outflow because the circulation in the extremity is, in the strictest sense of the word, peripheral, and because the femoral vein is more accessible than is the axillary.

METHOD OF EXPERIMENTATION

Our procedure was as follows: Under Grehant anesthesia the carotid artery was connected to a recording manometer and the blood tracing started. Both femoral veins were dissected free, ligated where they leave the thigh to enter the abdomen, clamped just below this, and divided between ligature and clamp. Paraffined canulas were tied into the distal ends of the veins. Both sciatic nerves were dissected free at the posterior aspect of both thighs, and a bridle ligature was placed under each nerve so that it could be drawn out and divided with ease and rapidity. Only a minimal amount of fall of blood pressure resulted from this preliminary, as a rule.

Fig. 1.—Tracing indicating outflow of blood in experiment on Dog 3. Moderate blood pressure. No evidence of shock. Division of sciatic (at X) followed by more rapid outflow of blood from femoral vein. Time markings, 1 second.

The clamp was now removed from the left femoral and the blood allowed to flow through the canula into a tall glass graduate of moderately small caliber. After about 25 or 30 c.c. had flowed out and the current of blood was running smoothly and evenly, readings were begun in order to note the rapidity of the flow. The method of making these readings was as follows: At the moment the reading was begun an electric button connected with the second-time marker was pushed. After 5 c.c. of blood had flowed the button was pushed again, and this was repeated at the completion of the flow of each fifth cubic centimeter of blood until the time record on the drum showed four or five periods, each one representing the time flow for 5 c.c. of blood. Then the left sciatic nerve was severed and, after waiting a short while for the flow to steady, the time flow for 5 c.c. was recorded several times. The animal was then reduced by a state of shock, by manipulation and chilling of the abdominal viscera, and after the blood pressure showed a permanent fall, the blood flow was measured from the right femoral vein, just as it had been from the left one, before and after section of the right sciatic nerve.

The vast majority of vasomotor impulses that reach the blood vessels of the hind leg of a dog travel through the sciatic nerve; therefore, as is well known, severance of this nerve results in vasodilation in the extremity, and consequently in a more rapid flow of blood from the femoral vein. This point is illustrated by the tracing shown in Figure 1. Before the sciatic was cut it took seven seconds for 5 c.c. of blood to flow out, whereas after the sciatic was divided the same quantity of blood flowed out in five seconds. The crux of these experiments lay in the rate of outflow after the animal was reduced to a state of shock, for if in profound shock the vessels are beyond the control of the vasomotor centers, then severance of the sciatic nerve ought not to make any appreciable difference in the rate of flow. All our experiments showed that the proportional rate of flow from the vein was more rapid after division of the sciatic in a shocked animal than it has been from the opposite veins before the animal had entered the state of shock. Figure 2 is a continuation of the tracing shown in Figure 1, taken from Dog 3 of our series. This portion of the tracing represents a period when the animal was in a profound degree of shock. It will be noted that before the sciatic was divided it took nineteen seconds for 5 c.c. of blood to flow, whereas after section of the nerve it required only eleven seconds. This shows that the intact sciatic must have been transmitting impulses that were maintaining vascular tone. More than this, the proportionately more rapid flow²¹ can not be due to an actually greater dilatation of the vessels in one instance than in the other, because in both instances we have a maximal dilatation consequent

Fig. 2.—Tracing indicating outflow of blood in experiment on Dog 3. Very low blood pressure. Animal in profound shock approaching *critus lethalis*. Division of sciatic (X) followed by a proportionately more rapid outflow from femoral vein than in Fig. 1. Time markings, 1 second.

on severance of the nerve. It must be due to the fact that during shock the vessels (arterioles and capillaries) are more contracted than they are before the animal shows the effects of trauma. It must be borne in mind clearly that these experiments do not show anything more than that the division of the sciatic nerve of a shocked animal causes dilatation of the vessels of the leg, just as it does in a normal animal. This dilatation, however, is proportionately greater than that which occurs in the normal animal, and this proportional difference must be referred to the fact that in the shocked animal, before division of the sciatic, the vessels are in a state of greater contraction than they are in the normal animal.

Two reasonable objections may be made to our results, on the ground, first, that the time measurements were not accurate in the strictest scientific sense, and, second, that the anterior crural nerve was not severed so as to cut off all possible vasomotor impulses to the leg. As

21. In the normal animal, the rate of flow before and after division of the sciatic was 7 to 5, whereas in the shocked animal the rate was 19 to 11. These two sets of figures demonstrate a proportionately more rapid outflow in the shocked animal.

regards time measurements, they were not taken automatically and were, therefore, not true to a fraction of a second. Repeated experiments, however, demonstrated that this source of error was minimal, and, furthermore, our object was not to measure the exact time rate. We did not sever the anterior crural, because we found this procedure to be unnecessary. Our object was not to cut off all impulses, but to block out most of them.

The task now remains to confirm by visual inspection the statement that the peripheral vessels are contracted. For reasons already stated, we selected the retina as our field for this part of our work, and we are indebted to Dr. John Green, Jr., for our ophthalmic reports. Unfortunately, we failed in efforts to devise a scheme for measuring, with absolute accuracy, the diameters of the retinal vessels. In order that Dr. Green might not be prejudiced by the facts already secured by us, he was not told about them, being merely requested to examine the fundi of the dogs' eyes, before and after shock was induced, and to note carefully the size of the vessels. He took the size of the optic disc as his standard and compared the size of the vessels with it. In every instance he reported that the vessels showed a marked degree of contraction after the animal was in shock. The vessels contracted down to from one-third to one-half of their size before shock.

Two observations were particularly interesting: In one case the dog died immediately after an examination of the fundus which had disclosed marked contraction of the vessels, thereby demonstrating that vasoconstriction persisted up to the very point of *exitus lethalis*. Unfortunately, we neglected to examine any of the fundi after death in order to see if vascular relaxation occurred. The second observation of especial interest was a case in which the retinal vessels were contracted so tightly as to make them resemble mere strands. In this connection it is interesting to note that ophthalmic literature contains at least two reports²² of sudden blindness occurring during shock, lasting from two to five hours, and then passing off without leaving any demonstrable eye changes or disturbances of vision. Roberts, who makes these reports, does not give the fundus findings during the state of shock, nor does he attempt to explain the cause of the blindness; but it seems fairly reasonable to assume that the occurrence of a sudden acute anemia of the retina, resulting from pronounced contraction of the retinal vessels, might cause transient blindness.

CONCLUSION

In conclusion it is important again to emphasize the fact that the experimental work detailed in this paper is in no sense of the word to be taken as an explanation of the causative factor or factors of shock. Just as Porter claims to show explicitly that in shock the vasomotor center is not exhausted and, therefore, implicitly that the peripheral vessels are not paralyzed, so we try to show explicitly that the peripheral vessels are contracted, and implicitly that not all the vasomotor centers can be exhausted. The complexity of the vasomotor apparatus with its multiplicity of centers, and the seemingly independent action of many of these centers, render it impossible to frame a satisfactory theory based on our results. This much may be said with assurance: If the work herein detailed stands the test of confirmation, then the doctrine that shock is due to vasomotor exhaustion must be revised.

Clinical Note

A TYPHOID BACILLUS CARRIER

HISTORY AND AUTOPSY

F. S. HAMMOND, M.D.

Assistant Physician and Pathologist, New Jersey State Hospital at Trenton; Pathologist to the Mercer Hospital
TRENTON, N. J.

Since the comparatively recent discovery of what may be termed latent typhoid infection and its bearing on the spread of the disease, as well as its pathologic and bacteriologic significance, additional examples of this condition have been reported repeatedly.

As yet, however, the number of postmortem examinations of bacillus carriers has not been large, and the present instance is cited, not only as an example of the possibilities of transmission of infection under certain circumstances, but as an additional confirmation of the generally accepted belief in the origin of the bacilli in such cases.

HISTORY OF A BACILLUS CARRIER

The individual in question was a colored man admitted as a patient to the New Jersey State Hospital at Trenton in 1902. He was at that time sixty-seven years old, and presented then, or subsequently, nothing beyond the gradual mental and physical failure incident to his age, and was able to perform such light work as was assigned him.

During July and August, 1907, the institution was visited by an epidemic of typhoid fever, due to the contamination of a spring from a leaking city sewer, over one hundred cases resulting, with the usual mortality.

The patient (C.C.) then seventy-two years of age, became ill early in August, and responded positively to the agglutination test on the 13th. The course of the disease was exceptionally mild; the temperature was at all times low, rarely above 100 F.; and there were few, if any, subjective or objective symptoms, so that, in spite of his age and comparative weakness, the patient was able to be about in a little over three weeks from the time of onset. Indeed, so mild was the disease that the diagnosis was made entirely on the Widal reaction, this test being made at this time on all patients showing any signs of ill health whatever. In the absence of an epidemic it is probable that the few symptoms observed would have given no indications of the condition.

He made an uneventful recovery, and very quickly regained his usual health.

During the fall and winter (1907), following the epidemic, bacteriologic examinations of the urine of all convalescent patients (including C.C.) were made at the hospital laboratory, three examinations each of fifty-two convalescents being performed. Digalski-Conradi agar medium was used for this purpose, and in no instance were suspicious organisms isolated.

All patients were at this time kept segregated after having had typhoid and hexamethylamin (urotropin) systematically administered over a period of months. Subsequent to the urine examinations, in the spring of 1908, the Research Laboratory of the New York Health Department, under the direction of Dr. W. H. Park, undertook the examination of the feces of the same individuals, samples being sent to New York for this purpose. Three or more examinations of specimens from each patient were made, and, as already reported by Dr. Park,¹ two bacillus carriers were discovered, one being the patient C. C. Four consecutive fecal examinations of this patient during March and April each showed typhoid bacilli in large numbers. He was at once isolated and rigid disinfection of all discharges carried out. No further cases occurred. It is here of interest to note what would have been the inevitable consequences, had not these individuals been recognized.

1. Park, William: Typhoid Bacilli Carriers, THE JOURNAL A. M. A., Sept. 19, 1908, II, 981.

22. Roberts, W. O.: Am. Pract. and News, 1907, p. 24.