The aneurism in the present case was situated in the subclavian, in the usual site of subclavian aneurism—namely, between the scaleni muscles, and to us seemed almost to invite some modification of the Dublin surgeons' plan by compression on the first rib. The plan by compression, we need hardly observe, is in general applied to the artery above the aneurism, between the latter and the heart. Crampton, however, in 1816, showed that the obliteration of an artery can be effected without rupture or ligature of its coats, as generally conceived, simply by this blocking-up process. The early volumes of THE LANCET contain cases also cured by Brasdor's operation; it seems, however, more applicable to carotid than subclavian aneurism.

Mr. Fergusson related to his class on the 11th, at some length, the details of a previous case of subclavian aneurism, of the same character as the present, in which his ideas on this subject were first matured. In both cases the method of cure by deligation at the tracheal side of the scaleni, as well as Brasdor's operation at the distal end of the aneurism, were inadmissible; yet it was gratifying to find the present plan, by firm pressure of the thumb on the aneurism, so as to displace some of the fibrinous clots, followed up by local pressure, succeeded in obtaining most striking and in many respects curious but satisfactory results. Intimately associated as the subclavian is at the right side with the vertebrals and carotid, the method of displacing fibrinous coagula is not without danger. A patient under such circumstances will fall down perhaps in a fit from want of circulation in one side of the "circle of Willis," formed by these arteries; yet, as the cause is so apparent, the danger may not be very alarming. Some instances of cure of aneurism of even the innominate have been given by American surgeons, in which recourse was had to ligature on Brasdor's plan of the subclavian; the result here ought to be equally dangerous. Hodgson gives us cases also in which a plug of effused lymph had nearly obliterated the subclavian; while Gendrin has imitated all the phenomena of arteritis and blocking up of aneurisms by injecting irritant substances into a portion of artery contained between two ligatures. In Mr. Fergusson's new mode of operation, we believe an entirely novel idea is acted on—namely, the displacement of the lamellated fibrin of the aneurism, on which no operation has been performed, and so directing the clots of fibrin that they shall block up the distal end of the artery so diseased. As Mr. Fergusson has expressed an intention of bringing the entire subject under the notice of the Medico-Chirurgical Society, we purposely abstain from giving the cases in detail. The method of treating aneurism by compression, originating with Desault and Hunter, and recently revived with such excellent results by the Dublin surgeons, will gain an immense accession of interest, if it should prove that the fibrinous deposit of the sac of the aneurism may be thus as it were utilized in bringing about the results hitherto gained in a different mode by Brasdor's operation at the distal end of the aneurism. Considerable caution will be at first necessary, as observed by Mr. Fergusson, in selecting cases which are fitted for the present method, as premature or ill-judged experiments in the shape of direct pressure or manipulation on the sac of aneurism not requiring it, one of which we mentioned recently as brought into Guy's, ("Mirror," *ante* p. 5.) where direct and prolonged pressure had been made in the popliteal space before the patient came into hospital would be certain to be followed by severe inflammation of the sac and other dangerous results. The spontaneous cure of aneurism is not unknown in practice; it may take place, it must not be forgotten, by a coagulation of the contents or increase of the quantity of lamellated blood in the sac, the cavity becoming filled, and the circulation conveyed to the parts beyond the disease by the collateral vessels; or, again, in some rare cases the aneurismal tumour may be doubled up and press upon the portion of artery leading directly to the aneurism; or in a third fashion, as in a remarkable case given by Mr. Liston, where the patient had well-marked subclavian aneurism, which subsided and disappeared—an aneurism of the innominate pressing on and obliterating the aneurism of the subclavian!

Whatever may prove to be the correct pathological explanation of the phenomena in Mr. Fergusson's present cases, we deem it our duty to state here briefly that the cure seems complete and unequivocal without any ligature of vessels, nor is there any reason to believe the case was one of spontaneous cure of subclavian aneurism, as in the case given by Mr. Liston. It is now two years since the man came first under observation; he has been, on and off, under treatment all that time in King's College Hospital and at home in the country; but happening to be in town within the last fortnight, Mr. Fergusson took advantage of the opportunity to exhibit the case to his class.

THE LONDON HOSPITAL.

INJURY OF THE FEMORAL ARTERY, CONSECUTIVE TO OPERATION; DELIGATION; RECOVERY.

(Under the care of Mr. CRITCHETT and Mr. ADAMS.)

The following case bears very significantly on the affections of arteries which surgeons are called on to remedy, and may find a place here. A man, aged about forty, otherwise apparently healthy, was admitted a short time ago into the London Hospital, with a large tumour of the lower extremity, which subsequently proved to be of fibro-plastic origin, situated on the inner aspect of the right thigh. Mr. Critchett and Mr. Luke, in consultation with Mr. Adams and Mr. Curling, advised removal of the tumour, which was accordingly undertaken by the first-named surgeon. During the operation, the sheath of the femoral vessels could be perceived quite exposed for some distance under Mr. Critchett's knife, so completely had the tumour involved all the deeper parts of the thigh in this situation, yet with care Mr. Critchett managed to avoid in any way endangering the sheath of the vessels thus included. The man lost very little blood, considering the gravity of the operation, and was placed again in bed, quite overjoyed at getting rid of such an incumbrance. The case went on very well until the eleventh or twelfth day, when the separation of the ligatures might have been expected. A sudden and profuse gush of arterial blood then took place, which Mr. Adams, who happened to be in the hospital at the time, believed had issued from the femoral, and on which view of the matter he at once proceeded to place a ligature on this vessel, observing the precaution, as considerable inosculation of vessels had taken place round the parts, to place a ligature on both ends of the wounded artery. In this case it was quite clear that, though in the operation by Mr. Critchett, the external coats of the femoral artery had not been injured, yet that the separation or insulation of this vessel for such a distance from the adjacent parts, or, in other words, the cutting off of the supply of small capillaries and vessels by which the sheath and its contents were previously nourished, had been more than the reparative power of the man's system was equal to, and sloughing had taken place, with consequent hæmorrhage. The ligatures have since all come away, and the man has left the hospital well.

POPLITEAL ANEURISM SUCCESSFULLY TREATED BY COMPRESSION.

A case of popliteal aneurism, treated by compression, has also been very successful this year, under the care of Mr. Adams and Mr. Ward. The patient, a healthy-looking, muscular man, ascribed his ailment to a kick from a horse. In the lower part of the right popliteal region, a smooth, round, characteristic pulsating tumour was readily to be detected. If the femoral was compressed over the pubes, the pulsation in the ear was quite commanded. The ear or stethoscope detected a bruit de soufflet. About a week after the patient's admission, the thigh was shaved and dusted with flour,—a very useful preliminary,—and compression used with a common 4lb. weight, and small compresses applied on the femoral as it passes near the groin. Two days after, a 7lb. weight was substituted for the 4lb., as the latter did not seem to control the pulsation sufficiently. Shortly after this the patient felt a pain over the skin of the knee and inner side of the thigh, resembling that produced by the pouring of hot water over a part, no doubt from some interference with the nerve. He was ordered infusion of digitalis, and on the 25th of January, the compression having been commenced on the 19th, the clamp tourniquet was tried on the lower part of Scarpa's triangle. This quite succeeded, and in less than a month from his admission he was dismissed cured, the pulsation in the tumour having entirely ceased. The integuments were not at all inflamed, and no necessity appeared for any further treatment, except placing a bandage on the limb, which having been done he left the hospital.

UNIVERSITY COLLEGE HOSPITAL.

CASE OF PUNCTURED WOUND OF THE PERICARDIUM; RECOVERY.

(Under the care of Mr. ERICHSEN.)

THE following is a short abstract of a case of wound of the pericardium, or, at least, wound in the situation of the heart in the fourth intercostal space, in which were well shown the various disturbances of the great central organs of the circulation likely to follow such an accident. We believe that in this case if the treatment had not been so active, there might have