Periscope

Brain

(Vol. 36, Part 2)

1. Experimental and Pathologico-Anatomical Researches on the Corpus Callosum. C. T. VAN VALKENBURG.
2. Sensory Changes in Friedreich's Disease. P. W. SAUNDERS.
3. The Wassermann Reaction and Its Application to Neurology. PAUL FILDES and JAMES MCINTOSH.
4. On the Bulbar Nuclei, with Special Reference to the Existence of a Salivary Center in Man. ANTHONY FEILING.

I. *Researches on the Corpus Callosum.*—On separating the two fore-brain hemispheres of any mammal, the corpus callosum becomes visible; its position between the two halves of the brain at once suggests that it serves as a connecting link between the hemispheres. There is no doubt that certain bundles exist in this mass of fibers, the origin and termination of which lie on the same side of the middle line. These are the fibers which belong, in the wide sense of the term, to the olfactory system, *i. e.*, the fornix longus, tænia tecta, and the nervi Lancisii. These, however, the author says, do not come within the province of his research, which is confined to what is commonly understood as the corpus callosum, that is to say, the chief hemispherical commissure of the neopallia. The term commissure may be understood to mean a connection of corresponding regions of the opposite hemispheres; or, taking it in a broader sense, a connection in general of the one half of the brain with the other, through which both homotopical and heterotopical cortical regions are associated with each other. To give the word its full meaning it connotes an interhemispherical association system. Both interpretations are found in the literature, though it is not possible to state that they always rest upon conclusive anatomical or pathological evidence. The more modern writers embrace the latter opinion on the whole, chiefly on the strength of Cajal's histological investigations; in small mammals, as the mouse, with the aid of his silver methods, Cajal found that cortical axones arise, which either directly or as collaterals of projection and longer association neurons, pass to homo- and heterotopical points of the opposite hemisphere through the corpus callosum.

The author, however, states his conclusion that the ending of the callosal fibers cannot be positively proved from normal silver preparations. Moreover, were normal histological investigations sufficient to place Cajal's theory beyond all question, they afford very little evidence of the "associative" nature of the commissure. Cajal believes that the callosal fibers give off collaterals before ending, but these can be followed with certainty over very insignificant distances only in one section. Though strictly speaking the term association may be applicable to such a connection, it approaches more nearly to what one may consider as a commissure. In any case it is evident that Cajal's theories, although they may be perfectly correct, by no means justify the frequent assumption that heterotopical

cortex regions of both hemispheres stand in connection by callosal fibers with each other.

The author believes he is justified in stating that in the posterior central convolution (1) The relation of the origin and termination of the callosal fibers is not definitely local. (2) The ending of these fibers is relatively diffuse. (3) Fibers end here which come from the contra-lateral gyrus centralis anterior, and to some extent at least the origin and termination both in front and behind the sulcus centralis are situated in the same horizontal level.

On the other hand, the anterior central convolutions are:

(1) Connected homotopically, in the sense that the origin of corticofugal fibers is more or less surrounded by a radiation of fibers from the homologous spot on the opposite side; it is doubtful if there exists a heterotopical connection between the gyri centrales anteriores; the author has no argument in favor of it.

(2) They are connected with the contra-lateral gyrus centralis posterior. The fibers which form this heterotopical connection are apparently far more numerous.

Commenting on clinical pathology, the author says that it is remarkable that of all the symptoms described in cases of callosal lesions very few may be ascribed, with any degree of certainty, to destruction of the commissural connections. Perfectly isolated callosal lesions rarely, if ever, occur, and the total absence of this great commissure does not necessarily cause any typical symptom. The symptomatology is very variable. Practically only those cases come under consideration where either an isolated, or more or less local, callosal softening existed, or in which by a softening in one hemisphere, a function is disturbed, which is not under the control of that hemisphere alone. Such disturbance, when remote influence by a tumor, etc., may be excluded, can only be explained through the commissural fibers.

The author believes that in the first place his investigation offers an anatomical explanation of the well-known fact that in cases of cortical epilepsy the excitation of a motor center in the affected hemisphere can cross through the corpus callosum to the homologous center in the healthy hemisphere. If the progression of the convulsions from the one side to the other is not accurately homotopical, as is indeed often the case, it may be assumed that the excitation of the center first affected was so strong that the irritation spread directly from here to the homologous region of the opposite hemisphere.

Indirect apraxia (apraxia callosa) of the left extremities which Liepmann has described, is assumed to be due to a lesion of the commissural connection between the left and right sensori-motor centers. This view, put forward by Liepmann, is closely connected with his statement that the left sensori-motor centers have a primary rôle in evoking more complicated actions in general; even the movements necessary for given action of the left limb are innervated from the right hemisphere. The result, the author says, of his anatomical investigations, that there exists a connection between the gyrus centralis anterior and the two contralateral central convolutions, may assist in the explanation of the fact that this direct apraxia itself can be considered as a "cortical" disturbance ("gliedkinetische, innervatorische Apraxie"). The movements of the left arm then are not disturbed in consequence of a deficient formulation of the "psychic plan," nor by the ineffective transmission of the "plan" to the region, whence the execution takes place finally, but by the inadequate execution of the

action itself, as this depends on the adequacy of the immediately preceding intra-cortical activity; but this must not be confused with the regulating impulses from the immediate proximity, the disturbance of which produces cortical ataxia. In the reported case in which only two handed movements were disturbed, it was because not only did these need the coöperation of the apraxic right hand, but the functions of the left hand were primarily affected. All remaining actions of the left hand, as far as they could be examined in the aphasic patient, were quite normal.

In addition to the complete apraxia, there was a strong disinclination to make any movement of the right hand or arm. Whether this innervation was affected by the extensive centro-parietal, or by the frontal softening, as Förster and others seem to believe possible, it is certain that the initiative of the left arm and hand was not lessened by these large lesions. There can be, in any case, no question of a unilateral "eupraxic center."

Finally, there is the question as to the causes of mind blindness. A lesion of the posterior part of the corpus callosum is generally considered, at least partly, responsible for this. If this assumes that it is due to the commissural connection between two primary visual fields (area striata) being disturbed, the view is untenable, as it is an anatomical fact that no such connection exists. If the chief cause of mind blindness is to be sought in a lesion of the callosal fibers, it must be of those that connect other parts of the two occipital lobes, and it is certain that fields 18 and 19 (Brodmann) receive and give rise to parastriate radiations. These regions of the cortex comprise the "visuo-psychic area" of the English authors. It is certain that these fields coöperate with the area striata in the higher visual functions. They are, moreover, phylogenetically more recent than field 17, and increase relatively in the higher mammals. Parallel with this the occipital callosal fibers become more abundant. The exactness of the hypothesis of Niessl v. Mayendorf, that in visual function the left hemisphere predominates over the right, and localized softenings of the pole and ventral part of the occipital lobe ("left macula center") may cause mind blindness, can be determined only by fuller clinical and anatomical investigations. One will then have to reckon with the fact that the area striata has no commissural connections.

2. *Sensory Changes in Friedreich's Disease.*—The sensory changes, says this author, have been studied by several observers. Noica, in two cases of the disease that he examined carefully, found very considerable loss to different forms of sensibility. Egger, in his investigations of "osseous sensibility," considered that this is regularly affected in Friedreich's disease. Other investigators have from time to time reported cases in which a certain amount of sensory loss was present, and cases indeed with some sensory loss have been mentioned incidentally from the time of the early differentiation of the disease from tabes dorsalis, and its occurrence is referred to in most neurological text-books.

The classical description of the disease, however, while admitting that loss to some or all forms of sensations may occur, regards it as altogether an unusual and exceptional feature, notwithstanding the fact that one of the great sensory plants of the cord is the chief site of the pathological changes that are present in the disease. A series of systematic observations, therefore, on a considerable number of cases of Friedreich's disease would seem justified, in order to see how far the classical description of the disease holds in the light of our present knowledge concerning the different forms of sensibility.

The author's chief aim in undertaking his work was the important

question of sensory conduction in the cord, as the more recent work of Petré, Rothmann, Head and his collaborators, and other investigators make it probable that impulses underlying the sense of position and appreciation of passive movement, the recognition of two compass points simultaneously applied to the surface of the body, the appreciation of vibration, and the recognition of size, shape, form and weight, and possibly of roughness and texture, are conveyed through the dorsal columns, tracts of which there is systematic degeneration in Friedreich's disease.

Most investigators now agree with the view that the dorsal columns furnish the principal pathway in the cord for the conduction of impulses that subserve muscle sense, meaning by that term chiefly the sense of position in space and the recognition of passive movement. Petré concludes from his examination of cases of stab wounds involving the spinal cord that these impulses pass up the cord uncrossed in the dorsal columns and in the dorsal spino-cerebellar tracts, but the view of Head and Thompson, based on their investigation of the grouping of sensory elements in Brown-Séquard lesions, that the dorsal columns subserve the conduction of the impulses underlying the recognition of position and movement that pass to consciousness, while the spino-cerebellar tract is concerned with those impulses that are responsible for the unconscious maintenance of equilibrium, seems likely to be the more correct, and Thompson has adduced later clinical and pathological evidence from a case of combined degeneration to show that the sense of position and recognition of passive movement may be grossly affected by a lesion limited to the long fibers of the dorsal columns.

Head and Thompson have shown also that the recognition of compass points simultaneously applied is dissociated in the cord from tactile sensibility, and passes up uncrossed in the dorsal columns, while Bing has brought evidences that the impulses that subserve the appreciation of vibration also pass up the homolateral side of the cord, and therefore must be conducted by the dorsal columns.

The investigations of Head and Holmes on Brown-Séquard lesions, cited in their paper go to show that not only do the knowledge of position and passive movement, the recognition of two points, and the appreciation of vibration depend on the integrity of the dorsal columns, but that the recognition of size, shape, form and weight, and the appreciation of consistence and texture are affected in common with these other elements of sensibility, and the impulses which underlie them would seem, therefore, also to be conducted by the dorsal tracts. Thompson also, in his case, noted that the recognition of weight was grossly affected, as well as the appreciation of position, movement and double contacts. Moreover, the author speaks of an unpublished case of cervical Brown-Séquard lesion, studied by Head and Holmes, which goes to confirm the view that the impulses which underlie recognition of size, shape, form, weight, consistence and texture, as well as the appreciation of position, movement, simultaneous contacts and vibration, are conducted through the homolateral side of the cord.

The impulses of light touch and pressure touch are, according to most of these observers, associated together in the cord. These impulses are conducted partly by an uncrossed path in the dorsal columns, mostly by a crossed path in the lateral columns (Petré), or in the ventral columns (Rothmann). The localization of tactile sensibility is associated with touch and is mainly the crossed pathway. All impulses of whatever kind that are painful, all heat impulses and all cold, cross almost at once after

a relay in the dorsal horns, and pass up the cord in the opposite lateral column, though possibly some of them pass without crossing by the lateral column of the same side. Systemic degenerations of the dorsal columns are not numerous. *Tabes dorsalis*, it is generally recognized, is, at least predominantly, a disease of the dorsal roots. In combined degeneration the process is probably not a true system degeneration but a funicular myelitis (Nonne). In Friedreich's disease, however, we have a condition which is usually regarded as essentially a systemic degeneration of the dorsal columns, even although the pathological changes are not limited to them. The changes in the dorsal column tracts are the most constant and most pronounced features of the pathological anatomy of the disease, and are present in all cases. In the sacral and lumbar regions these tracts may be almost entirely degenerated, but higher in the cord the degeneration in the column as a whole is not so marked. In the dorsal and cervical regions the root-entry zone seems considerably less affected than the rest of the column; and the cornu-commissural zone, the areas immediately next the grey matter of the dorsal horns, and Lissauer's tract may be scarcely at all affected. The degeneration in the other parts of the column, however, remains very severe throughout the whole cord. The long and medium fibers of the dorsal columns are therefore the ones chiefly affected, and the affection of these may be so intense that scarcely a fiber may remain visible in cross section.

In the other tracts involved in the disease, the pyramidal and the spino-cerebellar, the degenerative changes are rarely so great as in the dorsal columns. The affection of the pyramidal tracts is most intense in the sacral and lumbar regions and diminishes as we pass up the cord, though in some cases it may be traced as far as the pons or the lower part of the crus. The degeneration in the spino-cerebellar tracts, on the other hand, increases in extent and degree, from below upwards and becomes most pronounced in the higher cervical segments.

As contrasted with the great affection of the fibers in these different tracts, and especially in the dorsal columns, the changes in the cells are not so frequent or so pronounced except, perhaps, in Clarke's column. In some cases the spinal ganglion-cells are slightly atrophied and the dorsal spinal roots, and even the peripheral nerves may be involved. There is, however, no reason to consider that the affection of the dorsal columns is secondary to a primary degeneration of the ganglion-cells or of the roots, for the changes in the roots are not nearly so definite, so frequent, or so intense as in the dorsal columns, and the affection of the ganglion cells is slight and inconstant.

The widespread nature of the pathological changes in this disease and its variable incidence on different systems of fibers in the cord are in harmony with the view that it is a true systemic degeneration of the long spinal tracts, and especially of the dorsal column system. But even if the degeneration in the dorsal columns be regarded, not as a primary systemic one, but as due to an elective or chance degeneration in the dorsal roots, the existence of profound changes in these columns is by far the most outstanding feature of the pathology, and we should expect, therefore, to find in Friedreich's disease those forms of sensibility disturbed which have been assigned to the dorsal columns, and disturbed in proportion to the intensity of the dorsal column affection.

For the purpose of investigating this, the author examined twenty cases of Friedreich's disease, and found that all the patients examined presented the usual signs of Friedreich's disease, the skeletal deformities, the

absent deep reflexes, the extensor plantar responses, the nystagmus and the ataxia so characteristic of the disease.

The methods which he used were mainly those employed by Head and Holmes in their work on sensory changes in cerebral lesions in as far as these were applicable. They had to be modified somewhat, because the symptoms in Friedreich's disease being bilateral, there was no sound side which could be used as a standard of comparison to estimate the amount of loss present as in Head and Holmes's work, where the disturbance was unilateral.

He concludes that, in this disorder, the appreciation of touch, pain and temperature are very irregularly affected in the upper extremities, never more than very slightly and often not at all. When loss occurs it is almost always a slight distal blunting to touch, and very rarely to pin prick, or to heat and cold as well. In the lower extremities these cutaneous elements are more frequently involved, and there is often some distal hypoaesthesia to light touch or even to pressure touch, and in many cases also to pin prick and to heat and cold. It is the other elements of sensation, however, chiefly those that are commonly included in the term "deep sensibility," but also the appreciation of simultaneous contacts, and of size, shape, and form, that are most severely affected. The sense of position and the recognition of passive movement, the appreciation of double contacts and of vibration are almost always more or less affected, especially in the distal parts of the lower limbs, and not infrequently, though to a less degree, in the upper limbs as well. The frequency and the constancy with which these three elements of sensation are disturbed are indeed characteristic of the disease.

The recognition of shape and form, and the appreciation of size and weight are also affected, but much less frequently and less seriously than the sense of position, the appreciation of compass points, or of vibration. This is probably due to the fact that these forms of sensibility can be examined in the upper extremities only where the sensory loss as a whole in the early stages of the disease is usually not great, and possibly also to the fact that they are more elaborate forms of sensibility. They would seem, however, to be associated in their disturbance rather with the loss of the sense of position, the appreciation of double contacts and of vibration than with tactile loss. The recognition of shape, and that of size and of weight are affected with the same frequency, and usually in the same patients, while the recognition of form is less often disturbed.

The elements of sensibility, therefore, which it is assumed are conducted through the dorsal columns of the cord, would seem to be disturbed in Friedreich's disease more or less constantly and characteristically. The loss, especially in the early stages of the disease, falls most heavily on the lower extremities, that is on the regions innervated by the spinal segments that are earliest affected, and tends to spread higher and higher over the body and to become more profound as the disease advances.

Those elements of sensibility that are conducted through the dorsal columns have been divided into two groups. The recognition of two points of the compass, and the appreciation of size, shape and form which are dependent at the periphery on the integrity of the cutaneous nerve-endings would seem to run in one group, and the sense of position and of movement, the recognition of vibration and the appreciation of weight, which are dependent peripherally on nerve endings in joints, tendons and muscles, apparently compose another group. As the disturbance of the sense of position, of double contacts and of vibration was always so great

in every case in which the recognition of size, shape, form and weight was affected he found it impossible to correlate the disturbance in recognition of size, shape, and form with that in the recognition of compass points particularly; of the loss in the appreciation of weight with that of position. In the description of the fate of these elements of peripheral sensibility, therefore, he preferred for convenience to follow the order in which they were tested clinically, and to consider together the disturbance of the sense of position and of movement, the loss of appreciation of double contacts and the diminished recognition of vibration which constitute a more primary group; the recognition of size, form, shape and weight, would seem to compose a group of most secondary and elaborate sensations.

In a disease such as Friedreich's ataxia where there is a very slow progressive degeneration of the dorsal column of the cord, it is inevitable that though the sensory loss may conform to a definite type there must be much variability in the relative intensity of the affection of the different elements of sensibility in different cases. The frequency, however, with which a characteristic sensory loss does occur would seem to justify its inclusion in the clinical picture of the disease as an integral, not as an exceptional, feature; and further, the constancy and regularity with which this loss involves the sensory impulses assigned to the dorsal columns of the cord goes to confirm the conclusions regarding sensory conduction in the cord that have been obtained by the clinical and anatomical study of local spinal lesions.

3. *Wassermann Reaction in Neurology.*—The authors' intentions in writing this paper were to review the present state of knowledge of the Wassermann reaction in all particulars which bear directly or indirectly upon neurology, but in these particulars only. They have also described fully the methods utilized by them as being the most satisfactory according to their deductions. These deductions are based upon researches into the subject of syphilis generally, and also upon the results of a long collaboration with Head and Fearnside upon syphilis of the nervous system in particular.

The work is treated largely from a technical point of view, and thus assumes a general knowledge of the principles, and even of the technique, of the reaction on the part of the reader. The theory of the subject is not discussed, since that has no application to neurology, neither have the modified tests been reviewed.

The paper is intended to be read in conjunction with works published or forthcoming by Drs. Head and Fearnside and the authors themselves, and consequently constant references are made by means of index numbers, which remain the same throughout all these papers.

The conclusions drawn as a result of this extensive study are as follows:

1. In general diagnostic work, exclusive of nervous conditions, a positive reaction is so constant in active syphilis that such a diagnosis, made in the face of a negative result, will require very special support on other grounds before it can be considered probable.

2. A positive reaction in the cerebrospinal fluid indicates a syphilitic lesion of the central nervous system.

3. In active, untreated cases of dementia paralytica, tabes dorsalis, and cerebrospinal syphilis, the reaction is positive both in the serum and cerebrospinal fluid. In cerebral syphilis, without involvement of the spinal cord, the cerebrospinal fluid is usually negative. Except in the latter cases the strength of the reaction has no certain diagnostic value.

4. The reaction is apt to be negative in the serum or cerebrospinal fluid in cases of hemiplegia, non-progressive tabes dorsalis, and old lesions of the nervous system, especially in cases of stationary congenital syphilis. It is also usually negative in the serum particularly in very recent nerve lesions of the late secondary period, and in the cerebrospinal fluid particularly in treated cerebrospinal cases.

5. The occurrence of a positive reaction in one of the test-fluids when the other is negative is so common that a reaction found negative in one fluid only is insufficient to exclude syphilis.

6. Pure "parasyphilitic" lesions do not respond satisfactorily to treatment as shown by the slight effect upon the reaction in the cerebrospinal fluid. Cerebrospinal lesions react very readily to treatment. The celerity with which the reaction in the cerebrospinal fluid responds to treatment is an index of the "parasyphilitic" or "syphilitic" nature of the case.

7. The positive reaction in the serum is rapidly made negative by treatment in acute syphilis; but in chronic syphilis this effect is produced very slowly. In the cerebrospinal fluid, on the other hand, the rate of reduction of the reaction does not depend upon the duration of the infection. Thus, in gummatous lesions of the central nervous system, as also in secondary cerebrospinal syphilis, the effect of treatment upon the reaction in the cerebrospinal fluid is rapid.

8. The "provocative" injection of salvarsan for the purpose of exciting an exacerbation of the reaction has some, but probably an exaggerated diagnostic value.

4. *Bulbar Nuclei*.—Anatomical observations in man and experimental investigations in the lower animals have provided us with an extensive knowledge of the position and relations of the nuclei of the cranial nerves in the medulla oblongata. The author of the paper, which is based upon a case which afforded unusual facilities for investigation, does not claim to do more than confirm our previous knowledge of these nuclei. He does attempt, however, to go further and establish the existence of a salivary center in the medulla oblongata.

The patient, a miner, received a severe injury on the upper part of the neck on April 18, 1913. When a careful examination was made a few days later, the following conditions were observed: (1) Paralysis of the left side of the tongue. (2) Paralysis of the left sterno-mastoid and upper part of the left trapezius. (3) Loss of taste in the whole of the back of the tongue, more marked, however, on the left side. (4) Paralysis of the left side of the palate. No anesthesia of the palate could be detected. (5) Aphonic speech. By laryngoscopic examination the left vocal cord was seen to be immobile. Loss of power of swallowing, even for liquids. In addition to the above symptoms, a slight left facial paralysis was observed on the first day after the accident, but this rapidly disappeared. It is also stated that immediately after the accident some question arose as to whether the phrenic nerve was injured or not, but on the following day no paralysis of the diaphragm could be detected. At no time was there any paralysis of the extremities, nor with the exception of the paralysis of the left sterno-mastoid and trapezius muscles was there any limitation of the movements of the neck.

The author gives a full account of the autopsy findings and results of the microscopical investigations, and summarizes the anatomical changes as follows: (1) Degeneration in the cells of origin of the spinal accessory nerve in the ventro-lateral part of the grey matter of the upper cervical region of the spinal cord. (2) Degeneration throughout the whole region

of the nucleus ambiguus. (3) Partial degeneration of the dorsal nucleus of the vagus. (4) Partial degeneration of the hypoglossal nucleus. (5) Marked fatty changes in the solitary bundle. (6) Scattered degenerated cells in the grey matter of the upper part of the medulla just dorsal to the inferior olive.

There was only a small amount of fatty change present, it being practically confined to the solitary bundle. The length of time which elapsed between the injury and the death of the patient (over two months) is no doubt in part a reason for the limited amount of fatty changes present. Since the dorsal nucleus of the vagus, which is largely a sensory nucleus, showed only partial degeneration, it may be surmised that perhaps the injury to the nerve took place on the distal side of the ganglion, the degenerated cells representing the motor part of this nucleus. In this case, the fatty change in the solitary bundle would represent degenerated fibers from the glosso-pharyngeal nerve. It was, however, impossible to say at exactly what point in their course through the jugular foramen the nerves were injured. From the fact too that the hypoglossal nucleus showed only partial degeneration, we may assume that the injury to the nerve was not complete, a hypothesis borne out by an examination of the skull, for it was found that the fracture did not directly involve the anterior condyloid foramen.

JELLIFFE.

MISCELLANY

A STUDY OF THE METABOLISM OF CALCIUM, MAGNESIUM, SULPHUR, PHOSPHORUS, AND NITROGEN IN ACROMEGALY. Olah Bergeim, F. T. Stewart, P. B. Hawk. (The Journal of Experimental Medicine, Vol. XX, No. 3.)

The mechanisms underlying the metabolism of the mineral constituents of the human body are almost unknown. The present study is a tentative approach to certain aspects of the problems involved. The authors find that in acromegaly a distinct retention of calcium, magnesium and phosphorus was noted, which was not accompanied by corresponding changes in general metabolism, as evidenced by a practical balance of nitrogen and sulphur. It is concluded that there is a primary disturbance in the metabolism of the mineral elements mentioned, with the probable formation of new bony tissue. The distribution of elements between urine and feces shows no abnormal variations. A suggestion is made as to the reasons for variable calcium output in the urine on a uniform diet.

JELLIFFE.

LIVER FUNCTION AS INFLUENCED BY DUCTLESS GLANDS. G. H. Whipple and W. P. Christman. (Jour. of Exp. Medicine, Vol. 20, No. 3, p. 297.)

Physico-chemical metabolism mechanisms are slowly being analyzed with the result that vegetative neurology is opening up as a specialty almost of its own as a fundamental stratum of neurological knowledge. The relation of liver function to certain vegetative neurological structures is here discussed.

The authors find that when phenoltetrachlorophthalein is injected intravenously, it is eliminated from the body in the bile through the activity of the hepatic epithelium. The feces may be collected after purgation and the phthalein extracted and estimated against a standard solution. The