Annals and Magazine of Natural History: Series 6

Publication details, including instructions
for authors and subscription information:
<http://www.tandfonline.com/loi/tnah12>

LXI.—A new genus of aglossal batrachians

G.A. Boulenger F.R.S.

Published online: 02 Oct 2009.

To cite this article: G.A. Boulenger F.R.S. (1896) LXI.—A new genus of
aglossal batrachians , Annals and Magazine of Natural History: Series 6,
18:107, 420-420, DOI: [10.1080/00222939608680478](https://doi.org/10.1080/00222939608680478)

To link to this article: <http://dx.doi.org/10.1080/00222939608680478>

PLEASE SCROLL DOWN FOR ARTICLE

Taylor & Francis makes every effort to ensure the accuracy of
all the information (the "Content") contained in the publications
on our platform. However, Taylor & Francis, our agents, and our
licensors make no representations or warranties whatsoever as
to the accuracy, completeness, or suitability for any purpose of
the Content. Any opinions and views expressed in this publication
are the opinions and views of the authors, and are not the views
of or endorsed by Taylor & Francis. The accuracy of the Content
should not be relied upon and should be independently verified
with primary sources of information. Taylor and Francis shall not be
liable for any losses, actions, claims, proceedings, demands, costs,

expenses, damages, and other liabilities whatsoever or howsoever caused arising directly or indirectly in connection with, in relation to or arising out of the use of the Content.

This article may be used for research, teaching, and private study purposes. Any substantial or systematic reproduction, redistribution, reselling, loan, sub-licensing, systematic supply, or distribution in any form to anyone is expressly forbidden. Terms & Conditions of access and use can be found at <http://www.tandfonline.com/page/terms-and-conditions>

sexual maturity. He also describes an individual which, though internally it exhibited fully developed male sexual organs, as far as its exterior was concerned bore quite the character of a female, even possessing, indeed, rudiments of brood-lamellæ.

I incline towards the view that Isopods were originally altogether hermaphrodite. Such conditions, however, persisted only in parasitic forms (Cymothoidæ), for which hermaphroditism must be universally advantageous. In the case of free-living forms this state of affairs disappeared. In them we find that only scattered and scanty hermaphrodite remnants or rudiments are preserved.

LXI.—*A new Genus of Aglossal Batrachians.*

By G. A. BOULENGER, F.R.S.

AMONG the rich herpetological collections made by Stuhlmann in East Africa the frog described by Tornier as *Xenopus Boettgeri* (Thierw. Ost-Afr., Rept. Amph. p. 163, fig., 1896) is of exceptional interest in adding a second genus to the Dactylethridæ. As Tornier has not realized the systematic importance of the characters which differentiate the new species from *Xenopus laevis* and its allies, it is thought desirable to emphasize it by raising *Xenopus Boettgeri* to generic rank under the name of *Hymenochirus Boettgeri*. The principal characters that distinguish *Hymenochirus* from *Xenopus* are, so far as can be judged from Tornier's description and figure, the half-webbed fingers, the incompletely webbed toes, the third of which considerably exceeds the fourth in length, and, above all, the absence of lines of sensory muciferous canals on the body. No doubt a careful examination of the type specimen, unfortunately still unique, would reveal further differences, and the Röntgen rays might be usefully applied to obtain some information on the osteological characters.

LXII.—*Descriptions of Two new Frogs obtained in Madagascar by Dr. Forsyth Major.* By G. A. BOULENGER, F.R.S.

Mantidactylus Majori.

Closely allied to *M. curtus*, Blgr., but snout much longer, acutely pointed, and very strongly projecting beyond the mouth. Vomerine teeth in two small, oblique, oval groups close together behind the level of the choanæ. Head longer